



Cite this article: de Vries J, Woehle C, Christa G, Wägele H, Tielens AGM, Jahns P, Gould SB. 2015 Comparison of sister species identifies factors underpinning plastid compatibility in green sea slugs. *Proc. R. Soc. B* **282**: 20142519. <http://dx.doi.org/10.1098/rsob.2014.2519>

Received: 13 October 2014
Accepted: 7 January 2015

Subject Areas:

cellular biology, physiology, plant science

Keywords:

invertebrates, sacoglossa, kleptoplasty, reactive oxygen species, starvation, photosynthetic slugs

Author for correspondence:

Sven B. Gould
e-mail: gould@hhu.de

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsob.2014.2519> or via <http://rsob.royalsocietypublishing.org>.

Comparison of sister species identifies factors underpinning plastid compatibility in green sea slugs

Jan de Vries¹, Christian Woehle¹, Gregor Christa¹, Heike Wägele², Aloysius G. M. Tielens^{3,4}, Peter Jahns⁵ and Sven B. Gould¹

¹Institute of Molecular Evolution, Heinrich-Heine-University Düsseldorf, Universitätsstrasse 1, 40225 Düsseldorf, Germany

²Zoologisches Forschungsmuseum Alexander Koenig, Adenauerallee 160, 53113 Bonn, Germany

³Department of Biochemistry and Cell Biology, Faculty of Veterinary Medicine, Utrecht University, Utrecht, The Netherlands

⁴Department of Medical Microbiology and Infectious Diseases, Erasmus University Medical Center, Rotterdam, The Netherlands

⁵Plant Biochemistry and Stress Physiology, Heinrich-Heine-University Düsseldorf, 40225 Düsseldorf, Germany

The only animal cells known that can maintain functional plastids (kleptoplasts) in their cytosol occur in the digestive gland epithelia of sacoglossan slugs. Only a few species of the many hundred known can profit from kleptoplasty during starvation long-term, but why is not understood. The two sister taxa *Elysia cornigera* and *Elysia timida* sequester plastids from the same algal species, but with a very different outcome: while *E. cornigera* usually dies within the first two weeks when deprived of food, *E. timida* can survive for many months to come. Here we compare the responses of the two slugs to starvation, blocked photosynthesis and light stress. The two species respond differently, but in both starvation is the main denominator that alters global gene expression profiles. The kleptoplasts' ability to fix CO₂ decreases at a similar rate in both slugs during starvation, but only *E. cornigera* individuals die in the presence of functional kleptoplasts, concomitant with the accumulation of reactive oxygen species (ROS) in the digestive tract. We show that profiting from the acquisition of robust plastids, and key to *E. timida*'s longer survival, is determined by an increased starvation tolerance that keeps ROS levels at bay.

1. Introduction

Some sacoglossan slugs can house functional plastids for months in the cytosol of cells that line the animals' digestive tubules. The slugs steal the plastids (kleptoplasts) from siphonaceous algae upon which they feed. Theory has it that the presence of functional kleptoplasts [1,2] allows some of the sea slug species to survive starvation periods that can last almost a year [3,4]. Recent reports question the categorical importance of photosynthesis during starvation for several species [5,6] and while the animals fix CO₂ in a light-dependent manner [5,7–10], for how long and how much during starvation is not well documented. Animals induce autophagy to reallocate resources through the recycling of tissue when facing starvation [11] and mitochondria-generated reactive oxygen species (ROS) signalling has been found to be a key mediator of autophagy triggered by nutrient deprivation [12,13]. Over the past 20 years, research on 'photosynthetic (green) slugs' has focused on understanding how the stolen plastids can remain functional in a foreign cytosol [14], but the molecular response to starvation has not, to our knowledge, been assessed in sacoglossan slugs until now.

The kleptoplasts' stability outside of the algae disagrees with our knowledge about the sensitivity of land plant plastids, whose photosystem II is highly

susceptible to photosynthesis-associated damage [2,15,16]. Recent works [16,17] are reviving old observations that demonstrated some algal plastids are naturally more robust than land plant plastids [18]. Kleptoplast longevity therefore appears an intrinsic property of the stolen organelles bring along, and depending on the slug species this results in different modes of kleptoplast retention and lengths of survival during starvation [2,16]. It is generally suspicious that among 300 sacoglossan species that feed on siphonaceous algae, only seven species are known to maintain functional kleptoplasts and survive long term [2,19,20].

The two congeneric species *Elysia cornigera* and *Elysia timida* both feed on the ulvophycean alga *Acetabularia*, but only *E. timida* tolerates prolonged starvation [2,21]. Why? We compared the two sister taxa by monitoring their kleptoplasts' photosynthetic capacity and characterizing the slugs' physiological response to starvation focusing on gene expression modulation and ROS development. Our results indicate that algal cytosol are the slug's main food source and furthermore demonstrate that the plastid-bearing slugs' capacity to endure prolonged periods of starvation is not determined by the photosynthetic activity of their kleptoplasts, but by how they have evolved to respond to starvation, in particular the ability to cope with ROS.

2. Results and discussion

(a) Performance of stolen plastids does not depend on the slug species

We fed *E. timida* and *E. cornigera* solely on *Acetabularia acetabulum* D11 from the day they hatched in our laboratory. After eight and six weeks, respectively, and at which time the animals had reached maturity, the slugs were separated from the alga and routinely kept at $25 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ for 12 h every day. All sacoglossans that starve are observed to shrink [10,22] and the same was true for our two species. Yet, *E. cornigera* decreased in size more rapidly than *E. timida* (figure 1a) and rarely survived more than two weeks without food, which is consistent with previous reports [19,21]. Loss in size was accompanied by a slow fading of the animals' greenish coloration owing to the loss of the pigments chlorophyll (Chl) *b* and, most prominently, Chl *a* (electronic supplementary material, table S1). In *E. cornigera*, pigment concentration declined from starvation day 2 to 4 by 32.9%, but then increased again for a few days in relation to the dry weight of the animals (figure 1b). At day 10, the Chl *a* and *b* concentration had dropped from 3109 to 1544 pmol mg^{-1} animal dry weight (49.7%). These values correlate with the rapid decrease observed in body size between day 4 and 7 of starvation (figure 1a). In *E. timida*, we observed a similar, yet less steep, decline in the concentration of pigments in relation to body mass over the 10 days of starvation measured (from 2193 to 1606 at day 4 (73.2%) and 1638 pmol Chl *a* + *b*/mg animal dry weight at day 10 (74.7%); figure 1b). The increase of photosynthesis-associated pigments around day 6 demonstrates that the animals we measured metabolize their own tissue at an apparently faster rate than the kleptoplasts degrade, which was also observed for starving juveniles of *E. chlorotica* [23].

Photosynthetic capacity was determined through measurements of photosystem II quantum efficiency (derived from Chl

fluorescence analyses determined by pulse-amplitude modulation (PAM) fluorometry) and the amount of incorporated $^{14}\text{CO}_2$. Freshly fed slugs showed a similar initial maximum quantum yield (F_v/F_m of 0.83 ± 0.03 for *E. cornigera* and 0.85 ± 0.03 for *E. timida*). Measured F_v/F_m of *E. timida* declined slowly ($-0.001 F_v/F_m$ per day starved) over the one-month period, reaching 0.77 ± 0.07 at day 30. In *E. cornigera*, the decline of F_v/F_m was more prominent ($-0.008 F_v/F_m$ per day starved) over the 10 days measured, dropping to 0.75 ± 0.1 , at which point the species began to die (figure 1c). At the same time, the amount of $^{14}\text{CO}_2$ incorporation had dropped to 34% ($\pm 15\%$) and 36% ($\pm 12\%$) in *E. cornigera* and *E. timida*, respectively, after 10 days of starvation (figure 1c). At that point, the F_v/F_m had dropped to 0.84 ± 0.03 and 0.75 ± 0.1 for *E. timida* and *E. cornigera*, respectively. After 30 days of starvation, the amount of $^{14}\text{CO}_2$ fixed in *E. timida* had dropped to $4 \pm 1\%$ in regard to the initial amount. In any case, after 10 days of food deprivation *E. cornigera* died in the presence of intact (electronic supplementary material, figure S1) and functional kleptoplast. The difference of profiting from kleptoplasty must depend on the animals' tolerance to starvation, not the stolen organelles' performance. This observation challenges the idea that kleptoplasts could act as food depot [5,10], and raised the question, why in particular *E. cornigera* dies in the presence of functional kleptoplasts.

(b) Global gene expression response is predominantly governed by starvation

To uncover the differences of the two species in the response to starvation, and test whether the slugs can to some respect sense the kleptoplasts' status, we analysed gene expression changes throughout starvation and under different environmental stimuli. We performed comparative transcriptomics of the slugs: (i) under starvation (S), (ii) under starvation and in the presence of the photosynthesis inhibitor drug monolinuron (S + M), and (iii) under starvation and including a daily bleaching pulse with $1000 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ for 1 h d^{-1} (S + B). A total of 10 857 contigs for *E. cornigera* and 11 152 contigs for *E. timida* were assembled that were supported by at least 100 reads and homologous to eumetazoan sequences (electronic supplementary material, figure S2 and table S2). Global gene expression trends were confirmed for all conditions on six individual genes using quantitative reverse-transcription PCR (qRT-PCR) (electronic supplementary material, figure S3).

If the slugs have evolved to depend on the photosynthetic capacity of their kleptoplasts, we would expect to see a strong difference in the regulation of global gene expression profiles when kleptoplast electron transport is blocked (S + M) or when the kleptoplasts are under high light stress (S + B). Yet, when analysing the co-regulation of gene expression we found that starvation (S) in comparison to feeding (F) was the main denominator for the transcription response in *E. cornigera* (figure 2). The additionally applied treatments (S + B, S + M) only had marginal effects on the global expression profile of *E. cornigera* compared to starvation alone (S). In contrast to this, the response of *E. timida* throughout the first two weeks of starvation is more differentiated in regard to the individual stimuli and in comparison to *E. cornigera* (figure 2). After 4 d of S + M and 7 d S + B, the most differing expression was detected, indicating that *E. timida* indeed responded to the status of their kleptoplasts. Intriguingly, for *E. timida* a similar yet delayed transcriptional




Figure 1. *Elysia cornigera* dies while showing comparable photosynthetic activity to *E. timida* during starvation. Adult slugs were fed on *Acetabularia acetabulum* and the effects of starvation at $25 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ on the photosynthetic capacity of both species evaluated. (a) Representative images of one single specimen of each species. *Elysia cornigera* shrinks more rapidly than *E. timida*, and while the *E. cornigera* specimens never survived starvation for more than two weeks, all *E. timida* specimens survived and retained an almost equal body length during the 30 d of starvation analysed. (b) Analysis of pigment concentration in relation to dry weight of the slugs during the first 10 days of starvation. Quantification was performed on three biological replicates per time point and normalized to an external standard; each pigment extract was derived from three to seven *E. cornigera* (in total 143 individuals) and two *E. timida* individuals (in total 66 individuals). (c) Assessment of the maximum quantum yield (F_v/F_m ; measured using a PAM fluorometer) of photosystem II and net $^{14}\text{CO}_2$ fixation relative to freshly fed animals (t_0), during starvation within the same group of individuals. Each single F_v/F_m curve represents measurements on a group of at least 12 *E. cornigera* and nine *E. timida* specimens; from each group four *E. cornigera* (in total 48 individuals) and three *E. timida* (in total 54 individuals) were used for determining the $^{14}\text{CO}_2$ incorporation after 0, 4, 7, 10, 20 and 30 days of starvation (latter two only for *E. timida*).




Figure 2. Global co-expression analysis of gene expression changes. Pearson correlations were calculated by comparing the FPKM (fragments per kilobase of exon per million reads mapped) values of filtered genes of the different samples. Note the clear separation of the control and the remaining conditions and for *E. timida* the shift between the time points of 4 and 7 days in comparison to 30 days. The colour key indicates the degree of co-regulation. The bold numbers represent time of treatment in days. No values for *E. cornigera* were obtained for 30 days, because all individuals had succumbed to starvation (indicated by a cross). F, freshly fed; S, starving; M, monolinuron; B, bleach.

response to starvation was observed, as transcriptional profiles at day 30 began to reflect those observed for *E. cornigera* earlier during starvation based on the correlation of co-regulated genes within each species (figure 2). The slugs predominantly respond to starvation and not to the photosynthetic capacity of the kleptoplasts.

(c) Host response underpins kleptoplast compatibility

When mapping the transcriptomes to the Kyoto Encyclopedia of Genes and Genomes (KEGG) database (figure 3), two main trends were observed: first, downregulation of carbohydrate, energy, and lipid metabolism associated pathways (mean \log_2FC of -0.32 versus an overall mean \log_2FC of -0.10). On average, 81 of the 89 metabolism pathways detected for *E. timida* were downregulated after 30 days of starvation. Monolinuron, blocking the quinon-binding site of the D1 protein [24], enforced the downregulation of core metabolic genes in both species (at 7 d S + M \log_2FC -0.44 in both). Second, in contrast to *E. timida* (mean \log_2FC of -0.11), starvation in *E. cornigera* triggered an elevated expression of environmental information processes and cellular and organismal systems including stress response and autophagy (mean \log_2FC 0.13). Light stress further promoted this reaction in *E. cornigera* (mean \log_2FC of 0.20 at 4 d/7 d S + B) compared to *E. timida* (mean \log_2FC of -0.03 at 4d/7d S + B). Starvation imposes a metabolic dormancy and gene expression profiles to a minor degree reflect kleptoplast manipulation. The activity of the kleptoplasts, however, does not explain the survival during starvation.

Ubiquitination is a key mechanism to tag organelles and misfolded proteins—accumulating under stress conditions such as starvation—for degradation [25,26]. Expression levels of polyubiquitin remained the same in both species, except for a single polyubiquitin homolog of *E. timida* (Eti000028, 4.5-fold upregulation, $p < 0.05$). For the ubiquitin binding autophagic receptor p62/SQSTM1 [27], an average of only 28 FPKMs (fragments per kilobase of exon per million reads

mapped) were identified for *E. cornigera*, but 659 FPKMs for *E. timida*. And only for the latter species an upregulation of p62/SQSTM1 was observed during starvation (electronic supplementary material, table S3). *Elysia timida* might be better equipped to detect damaged proteins and organelles than *E. cornigera*, allowing the former to better recognize and remove damaged organelles and misfolded hazardous proteins during starvation. The ability to recognize damaged kleptoplasts for their subsequent degradation could explain how the slugs profit from stored kleptoplasts. Only malfunctioning and ROS-leaking plastids would be detected by the autophagosomal machinery and removed; the profit of kleptoplasts would be an indirect one. This would also explain the more rapid decline of F_v/F_m values in *E. cornigera* in comparison to *E. timida*, while retaining the same CO_2 fixation rate (figure 1c). Accumulation of damaged plastids that fix no CO_2 would lead to a lower F_v/F_m ratio in *E. cornigera*. If *E. timida* specifically removes damaged plastids, then all malfunctioning photosystems are excluded from that equation.

We found that the upregulation of genes encoding ROS scavengers differed between the two sister taxa. With progressing starvation, *E. cornigera* elevated the expression of superoxide-dismutase homologs regardless of the treatment (e.g. Eco000152; on average 3.1-fold up at day 7, $p < 0.05$). Apparent changes in the expression of catalases were only detected for *E. timida* in the first days of starvation (Eti001444; up more than twofold on average). It was recently noted that a crucial part of the slug-plastid symbiosis is evolving mechanisms to cope with kleptoplast-associated toxins [2]. A difference in stress response and ROS evolution could explain the different survival rates of the two species observed during starvation.

(d) *Elysia cornigera* accumulates reactive oxygen species, not *Elysia timida*

ROS are known as signalling molecules [13] but also as deleterious cytotoxins [28]. To monitor hydrogen peroxide




Figure 3. Gene expression changes of animal KEGG pathways. Transcriptomes were sequenced for both animals after 4 and 7 days of starvation (bold numbers), and additionally after 30 days for *E. timida*. *Elysia cornigera* and *E. timida* were subjected to three different conditions: starvation alone (S), starvation plus monolinuron (S + M; 2 $\mu\text{g ml}^{-1}$) and starvation plus bleaching (S + B; 1000 $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$ for 1 h each day). Upregulation is shown from red to black and downregulation from black to blue. The numbers at the bottom represents the amount of K-numbers (KEGG orthology group identifier) found within each reaction module (modules in alternating light and grey flags). The bars on the right show the total number of regulated genes (minimum of twofold) for the individual conditions analysed (blue bars, downregulation; red bars, upregulation).

(H_2O_2) production on tissue level we used 2',7'-dichlorofluorescein (DCF). Freshly fed slugs showed no detectable amount of H_2O_2 (electronic supplementary material, figure S4), but at day 10 of starvation H_2O_2 was most prominently detected in *E. cornigera* within the lumen of the digestive tubules (figure 4a,e). Both the lumen and the epithelium of the digestive tubules—the localization of the kleptoplasts—showed intensive DCF staining (figure 5). By contrast, DCF staining in *E. timida* occurred only in some globular structures of 10–20 μm in diameter, but of otherwise unknown nature, within individual digestive tubules (figure 4g, arrowhead). After 30 d of starvation, the digestive glandular tubules of *E. timida* still showed no apparent increase in H_2O_2 levels (electronic supplementary material, figure S4), but DCF now additionally stained globular structures of approximately 100 μm in diameter that were situated between the epidermis and the digestive tubules, and independent from the latter. These structures could represent one of the secondary metabolite-harboring vacuoles typical for these kind of slugs [29]. Stroma of the *A. acetabulum* kleptoplasts resting within the digestive tubule cells of both *E. cornigera* and *E. timida* showed equal intensity in regard to H_2O_2 levels (figure 4c, f insert). Mitochondria, a common source of ROS [13], were found to accumulate H_2O_2 as proved through the co-localization of the DCF signal with MitoTracker Red (figure 4d). Hydrogen peroxide, a critical ROS in terms of cytotoxicity [28], accumulates only in *E. cornigera*.

Animals respond to nutrient depletion by autophagy, and superoxide ($\text{O}_2^{\cdot-}$) is a key signal that triggers the autophagosomal machinery [12,13] and mammalian tissues cultures show strong increase in ROS after a few hours of starvation [12]. We conducted nine separate experiments, each time with different individuals, and monitored $\text{O}_2^{\cdot-}$ evolution using dihydroethidium (DHE) which stains the nuclei of cells that experience elevated levels of $\text{O}_2^{\cdot-}$ [30]. Chl autofluorescence was observed to decrease, but not completely fade, in both species throughout the 10 d monitored (figure 6a). This concurs with the pigment measurements and the ability of the remaining kleptoplasts to incorporate $^{14}\text{CO}_2$ (figure 1c). Feeding slugs showed only faint $\text{O}_2^{\cdot-}$ signals towards the edges of the parapodia, especially the epidermis (figure 6a). Staining did not predominantly dye nuclei of the plastid-bearing digestive gland cells, but instead those of surrounding and connective tissue of the parapodia. Levels of $\text{O}_2^{\cdot-}$, apparent through the number of nuclei stained and the intensity of the staining itself, steadily increased throughout starvation and predominantly in *E. cornigera* (figure 6a). While both slug species harboured functional kleptoplasts stemming from the same source, only *E. cornigera* exhibited increasing levels of $\text{O}_2^{\cdot-}$ that were significant, but not *E. timida* ($p < 0.001$). *Elysia cornigera* accumulates ROS to a much higher degree than its sister taxon (figure 6b), reflecting the different capacities of the two species with regard to managing starvation in the presence of functional kleptoplasts.




Figure 4. Hydrogen peroxide accumulates in the digestive tubules of starving *E. cornigera*, but not *E. timida*. Confocal laser scanning micrographs of the kleptoplast-bearing digestive tubules that pervade the parapodia of 10 days starved, whole mount *E. cornigera* and *E. timida* stained with 100 μM dichlorofluorescein (DCF; blue) and MitoTracker Red (green); chlorophyll autofluorescence in red to yellow (*a,e*) or red only (*b–d, f, g*). Digestive tubules of *E. cornigera* (lined with autofluorescent kleptoplasts) show well-defined DCF staining within the lumen and epithelium (*a–d*). Details of single digestive tubules revealing readily H_2O_2 accumulators of equal size as kleptoplasts (*b* arrowheads), kleptoplasts displaying weak DCF staining (*c*; see arrowheads) and MitoTracker Red co-localizing with DCF staining (*d*). Digestive tubules of *E. timida* displaying no distinguishable DCF signal in epithelium or lumen (*e*). Details of single digestive tubules revealing accumulating weak levels of H_2O_2 (*f*, arrowheads), comparable to those seen in *E. cornigera* (*c*, arrowheads) and showing globular structures of unknown nature accumulating H_2O_2 (*g*, arrowhead).




Figure 5. Hydrogen peroxide is present in the lumen and epithelium of digestive tubules of starving *E. cornigera*. Orthogonal view of a confocal laser scanning z-stack micrograph reconstructed from 58 individual optical sections (21.35 μm of depth in total) of a single digestive tubule (outlined by dashed lines in the bottom panel) of 10d starved *E. cornigera* stained with 100 μM dichlorofluorescein (DCF; purple-blue to green gradient); chlorophyll autofluorescence in red. At the top, a transverse section through the stack is shown to illustrate at which level the three coronal sections (indicated by dashed lines) were captured.

3. Conclusion

Contrary to the current view our study demonstrates that the presence of functional kleptoplasts alone does not underpin the survival of green photosynthetic slugs. It is first and foremost the ability of some species to tolerate starvation. The best hints we currently have that underpin starvation

tolerance include the upregulation of a set of canonical ROS quenchers such as catalases, and/or slug-specific polypropionate metabolites such as Elysiapyrone or Elysiene [31], and/or the efficient and specific degradation of damaged kleptoplasts. Literature has divided slugs into non-, short- and long-term retention species, depending on for how




Figure 6. Superoxide accumulation in starving slugs. (a) Confocal laser scanning micrographs of the kleptoplast-bearing digestive tubules from similar positions on the parapodia of freshly fed and 4, 7 and 10 days starved, whole mount *E. cornigera* and *E. timida* (optical sections; pinhole set to 1 airy unit) stained with 100 μM DHE (staining $O_2^{\cdot-}$), shown in blue–yellow gradient; chlorophyll autofluorescence of the kleptoplasts is shown as red to yellow glow. Dashed lines mark the outer rim of the parapodia. DHE signal increases over time in *E. cornigera*, while the diameter of the digestive tubules decreases, visible due to their lining with kleptoplasts. Superoxide in the parapodia of *E. timida* does not significantly increase over time, while kleptoplasts lining the digestive tubules appear less densely packed. (b) Quantification of the mean DHE fluorescence intensity in the parapodial tissue; significance groups (for (a,b) $p < 0.01$; (c) $p < 0.05$) were determined using a Mann–Whitney U -test; measurement of fluorescence intensity was performed on greater than or equal to six individual specimens for each time point; error bars indicate standard deviation.

long kleptoplasts are functionally retained [19]. Our results demonstrate that it is equally important to divide the slugs into starvation-intolerant and starvation-tolerant species. That tolerance might be associated with the habitat and availability of the food source. *Elysia cornigera* was collected from the Caribbean Sea, but *E. timida* from the Mediterranean. Maybe only *E. timida* experiences food shortage due to seasonal differences and a difference in the calcification of individual algal species, and had to evolve accordingly. In any case, sacoglossan research has focused on how the animals chaperone their kleptoplasts, but it should just as much focus on how they manage starvation, because it will teach us on how they suppress ROS stress.

4. Experimental procedures

Elysia timida was collected on Giglio, Italy (42°22' N, 10°52' E and 42°21' N, 10°52' E) between 3 and 6 m depth. *Elysia cornigera* was collected at the Florida Keys (24°38' N, 81°18' W) at about 1 m depth. Both species were kept at a 12 L:12 D rhythm at 25 μmol quanta $m^{-2} s^{-1}$ in 3.7% salt water (Tropic Marine) and at 21°C. *Acetabularia acetabulum* DI1 (isolate of Professor Diedrik Menzel, Bonn University, Germany) was grown at the same conditions as the animals, but in algal f/2 medium. For starvation experiments, animals were moved to Petri dishes (Ø200 mm) and 50% of the water was changed every other day. Monolinuron (JBL GmbH) was added from a stock solution of 4000 mg l^{-1} to a final concentration of 2 μg ml^{-1} . Bleaching was performed by illuminating the slugs with 1000 μmol quanta $m^{-2} s^{-1}$ for 1 h every day at the same time. For each pigment extraction, pooled slugs were transferred to 90% acetone for each time point, homogenized and kept at –20°C for 1 day. Extractions for every time point were carried out as biological triplicates, each replicate consisting of at least three *E. cornigera* and two *E. timida*. Crude extracts were filtered through a 200 nM polytetrafluoroethylene

membrane and then analysed by reversed-phases high pressure liquid chromatography with ultraviolet/visible spectroscopy detection (Hitachi/Merk) as described earlier [32]. Pigment concentrations were determined using external pigment standards isolated from spinach thylakoids. Dry weight was determined on dried slug homogenate after extraction.

Photosystem II maximum quantum yield (F_v/F_m) was determined using a WALZ MINI-PAM as described previously [33], on the same group of individuals that were used for $^{14}CO_2$ incorporation measurements. For every time point for which $^{14}CO_2$ incorporation was determined, an individual group of slugs (at least 12 *E. cornigera* or nine *E. timida*) was kept in isolated dishes. For each of these groups, separate F_v/F_m measurements were performed, and each individual slug of a group was measured at least three times. For each data point, the mean of all individual triplicate measurements was calculated. The light-driven incorporation of $^{14}CO_2$ was determined as described earlier [5]. Briefly, for each triplicate measurement four individuals of *E. cornigera* and three of *E. timida* were used. Slugs were incubated in 1.2 ml artificial seawater supplemented with 0.40 mM [^{14}C]-NaHCO₂ (25 μCi per incubation, NEN-radiochemicals, MA, USA) for 2 h at room temperature and 72 μmol quanta $m^{-2} s^{-1}$ illumination. After rinsing, homogenization and acidification of the material with 150 μl 1 M HCl, all the substrate was expelled from the homogenate. Incorporation of ^{14}C by the slugs was determined in a scintillation counter after the addition of 12 ml LUMA-Gel scintillation cocktail (LUMAC, The Netherlands).

For ROS-imaging, slugs were incubated for 45 min with 100 μM DHE (DHE; excitation/emission HeNe 543/610 nm; Sigma) or 100 μM DCF (2',7'-dichlorofluorescein; excitation/emission Argon 488/529 nm) plus 2 μM MitoTracker Red CMXRos (excitation/emission HeNe 543/599 nm; LifeTechnologies) in artificial seawater. Slugs were rinsed twice with seawater and decapitated before mounting. Confocal laser scanning microscopy was carried out using a Zeiss LSM 710. Imaging was always performed with the same settings and

at a similar position along the parapodial rim at the same depth relative to the epidermis. Images were processed using Fiji/IMAGEJ 1.48f [34] and statistics were performed using R [35]. Normality was tested via a Shapiro–Wilk test [36] and significance using a Mann–Whitney *U*-test [37].

Total RNA was extracted three times from seven *E. cornigera* (a total of 21) and three times from five *E. timida* (a total of 15) feeding on *A. acetabulum* (t_0 , feeding control group), 28 individuals of *E. cornigera* that were starved for 4 days, of which nine individuals had only starved (S), nine had starved and were treated with monolinuron (S + M), and 10 of which had starved and were treated with 1000 μmol quanta $\text{m}^{-2} \text{s}^{-1}$ for 1 h each day (S + B). The RNA of 20 *E. timida*'s (7 S, 7 S + M, 6 S + B) was isolated after four days of starvation (t_1), of 30 *E. cornigera* (10 S, 9 S + M, 11 S + B) and of 20 *E. timida* (6 S, 8 S + M, 6 S + B) after 7 days of starvation (t_2), and finally of 32 *E. timida* (11 S, 10 S + M, 11 S + B) after 30 days of starvation (t_3). RNA was isolated using TRIzol (Life Technologies) according to the manufacturer's protocol and an additional DNase treatment (Thermo Fisher Scientific). Poly(A) mRNA enrichment, library preparation using the TruSeq kit (Illumina) and 100 bp paired end sequencing using the Illumina HiSeq2000 system was performed by the Beijing Genome Institute (Hong Kong). Reads with remaining adapter sequences, more than 5% of unknown nucleotides or more than 20% of nucleotides with quality scores less than 10 (Illumina 1.5) were removed as part of the raw data extraction.

A total of 1 186 405 486 reads (a minimum of 52 million reads/library) were obtained (electronic supplementary material, table S2). Reads were inspected by FASTQC v. 0.10.1 [38] and filtered and trimmed using TRIMMOMATIC 0.32 [39] (parameters: ILLUMINACLIP:TruSeq3-PE.fa:2:30:10; HEADCROP:10; TRAILING:3; SLIDINGWINDOW:4:20; MINLEN:36). Reads of all samples were assembled using TRINITY r20131110 [40], which yielded 249 855 and 150 314 transcripts for *E. cornigera* and *E. timida*, respectively. Expression values for transcript clusters (or unigenes) were extracted and analysed using the TRINITY pipeline [41] via RSEM 1.2.11 [42], BOWTIE 1.0.1 [43] and EDGER 3.4.2 [44]. Only unigenes with raw read counts of greater than or equal to 100 over at least two samples were analysed, resulting in 36 380 and 32 897 unigenes, respectively. Differential expression was determined by \log_2 fc of at least ± 1 compared to the control set of unigenes (with raw read counts of greater than or equal to 100). Expression change significance was assessed based upon the dispersion of available replicate information in EDGER. The longest isoform of each cluster was defined as the representative sequence for a unigene.

To plot the overall taxonomic distribution of the assembled transcripts, unigenes were subjected to a BLASTx-based [45] search (*e*-value cut-off 10^{-10}) against a database consisting of protein sequences of REFSEQ version 64 [46] plus those of the genomes of *Crassostrea gigas* [47] and *Lottia gigantea* [48] (electronic supplementary material, figure S2). For all downstream analyses, only those genes were included for which top hits to the mentioned organisms or other metazoans were retrieved. Excluded were those with best blast hits to plants, protists, prokaryotes and viruses. Protein annotations for the 14 848 *E. cornigera* and 13 875 *E. timida* unigenes were extracted from mentioned BLAST hits and from a second BLASTx search to the UniProtKB/Swiss-Prot database [49]. KOG/NOG (EuKaryotic/Non-supervised Orthologous Groups [50–52]) categories were determined based on the best BLASTx hit to protein sequences of EGGNOG v. 4 [53]. KEGG [54] accessions were obtained using the KAAS 1.6a webservice [55] against metazoa.

For qRT-PCR, 200 ng RNA was reverse transcribed using random hexamers and the iScript Select cDNA Synthesis Kit (Bio-Rad). Two-step qRT-PCR was performed on biological triplicates for each time point and treatment (each containing pooled RNA from greater than or equal to five slugs) using the SsoAdvanced Universal SYBR Green Supermix (Bio-Rad) according to the manufacturer's instructions on a StepOne Plus (Applied Biosystems) real-time PCR system. Primers were designed using Primer-BLAST [56] (for primer sequences, see the electronic supplementary material, table S4) and data were analysed according to Pfaffl [57], using the *EcRPL38* (Eco000149), *EtRPL19* (Eti000121) and *EtSETMAR* (Eti000317) as reference genes.

Data accessibility. All sequencing data was deposited at the NCBI Single Read Archive (SRP047359) and the Transcriptome Shotgun Assembly Sequence Database (*E. cornigera*: GBRW000000000; *E. timida*: GBRM000000000).

Acknowledgements. We thank Margarete Stracke for assistance in culturing, Steffen Köhler (CAi, HHU) for help with slug photography and Sophie Rommel (Population Genetics, HHU) for help with the statistical analysis. We thank the 'Zentrum für Informations- und Medientechnologie' (ZIM, HHU Düsseldorf) for computational support and infrastructure.

Author contributions. J.d.V., A.G.M.T., P.J. and S.B.G. carried out the main experiments. G.C., J.d.V. and H.W. additionally collected animals and C.W. processed the RNA-seq data. S.B.G. designed the experiments and wrote the manuscript together with J.d.V. All authors discussed the results, read and approved the final manuscript.

Funding statement. Funding through the Deutsche Forschungsgemeinschaft to S.B.G. (GO1825/4-1) is gratefully acknowledged.

References

1. Trench RK. 1969 Chloroplasts as functional endosymbionts in the mollusc *Tridachna crispata* (Bérgh), (Opisthobranchia, Sacoglossa). *Nature* **222**, 1071–1072. (doi:10.1038/2221071a0)
2. de Vries J, Christa G, Gould SB. 2014 Plastid survival in the cytosol of animal cells. *Trends Plant Sci.* **19**, 347–350. (doi:10.1016/j.tplants.2014.03.010)
3. Rumpho ME, Worful JM, Lee J, Kannan K, Tyler MS, Bhattacharya D, Moustafa A, Manhart JR. 2008 Horizontal gene transfer of the algal nuclear gene *psbO* to the photosynthetic sea slug *Elysia chlorotica*. *Proc. Natl Acad. Sci. USA* **105**, 17 867–17 871. (doi:10.1073/pnas.0804968105)
4. Cruz S, Calado R, Serôdio J, Cartaxana P. 2013 Crawling leaves: photosynthesis in sacoglossan sea slugs. *J. Exp. Bot.* **64**, 3999–4009. (doi:10.1093/jxb/ert197)
5. Christa G, Zimorski V, Woehle C, Tielsen AGM, Wägele H, Martin WF, Gould SB. 2014 Plastid-bearing sea slugs fix CO_2 in the light but do not require photosynthesis to survive. *Proc. R. Soc. B* **281**, 20132493. (doi:10.1098/rspb.2013.2493)
6. Klochkova TA, Han JW, Chah K-H, Kim RW, Kim J-H, Kim YK, Kim GH. 2013 Morphology, molecular phylogeny and photosynthetic activity of the sacoglossan mollusc, *Elysia nigrocapitata*, from Korea. *Mar. Biol.* **160**, 155–168. (doi:10.1007/s00227-012-2074-7)
7. Trench RK, Boyle JE, Smith DC. 1973 The association between chloroplasts of *Codium fragile* and the

- mollusc *Elysia viridis*. *Proc. R. Soc. Lond. B* **184**, 63–81. (doi:10.1098/rspb.1973.0031)
8. Clark KB, Jensen KR, Stirts HM, Fermin C. 1981 Chloroplast symbiosis in a non-elysiid mollusc *Costasiella liliana* Marcus (Hermaeidae: Ascoglossa (=Sacoglossa): effects of temperature, light intensity, and starvation on carbon fixation rate. *Biol. Bull.* **160**, 43–54. (doi:10.2307/1540899)
 9. Marín A, Ros M. 1989 The chloroplast-animal association in four Iberian sacoglossan Opisthobranchs: *Elysia timida*, *Elysia translucens*, *Thuridilla hopei* and *Bosellia mimetica*. *Scient. Mar.* **53**, 429–440.
 10. Christa G, de Vries J, Jahns P, Gould SB. 2014 Switching off photosynthesis: the dark side of sacoglossan slugs. *Commun. Integr. Biol.* **7**, e28029. (doi:10.4161/cib.28029)
 11. Singh R, Cuervo AM. 2011 Autophagy in the cellular energetic balance. *Cell. Metab.* **13**, 495–504. (doi:10.1016/j.cmet.2011.04.004)
 12. Scherz-Shouval R, Shvets E, Fass E, Shorer H, Gil L, Elazar Z. 2007 Reactive oxygen species are essential for autophagy and specifically regulate the activity of Atg4. *EMBO J.* **26**, 1749–1760. (doi:10.1038/sj.emboj.7601623)
 13. Li L, Chen Y, Gibson SB. 2013 Starvation-induced autophagy is regulated by mitochondrial reactive oxygen species leading to AMPK activation. *Cell. Signal.* **25**, 50–65. (doi:10.1016/j.cellsig.2012.09.020)
 14. Pelletreau KN, Bhattacharya D, Price DC, Worful JM, Moustafa A, Rumpho ME. 2011 Sea slug kleptoplasty and plastid maintenance in a metazoan. *Plant Phys.* **155**, 1561–1565. (doi:10.1104/pp.111.174078)
 15. Nickelsen J, Rengstl B. 2013 Photosystem II assembly: from cyanobacteria to plants. *Annu. Rev. Plant Biol.* **64**, 609–635. (doi:10.1146/annurev-arplant-050312-120124)
 16. de Vries J, Habicht J, Woehle C, Huang C, Christa G, Wägele H, Nickelsen J, Martin WF, Gould SB. 2013 Is *ftsH* the key to plastid longevity in sacoglossan slugs? *Genome Biol. Evol.* **5**, 2540–2548. (doi:10.1093/gbe/evt205)
 17. Jesus B, Ventura P, Calado G. 2010 Behaviour and a functional xanthophyll cycle enhance photo-regulation mechanisms in the solar-powered sea slug *Elysia timida* (Risso, 1818). *J. Exp. Mar. Biol. Ecol.* **395**, 98–105. (doi:10.1016/j.jembe.2010.08.021)
 18. Giles KL, Sarafis V. 1972 Chloroplast survival and division *in vitro*. *Nat. New Biol.* **236**, 56–58. (doi:10.1038/newbio236056a0)
 19. Händeler K, Grzybowski YP, Krug PJ, Wägele H. 2009 Functional chloroplasts in metazoan cells: a unique evolutionary strategy in animal life. *Front. Zool.* **6**, 28. (doi:10.1186/1742-9994-6-28)
 20. Christa G, Gould SB, Franken J, Vleugels M, Karameinski D, Händeler K, Martin WF, Wägele H. 2014 Functional kleptoplasty in a limapontioidean genus: phylogeny, food preferences and photosynthesis in *Costasiella* with a focus on *C. ocellifera* (Gastropoda: Sacoglossa). *J. Mollusc. Stud.* **80**, 499–507. (doi:10.1093/mollus/eyu026)
 21. Krug PJ, Händeler K, Vendetti J. 2011 Genes, morphology, development and photosynthetic ability support the resurrection of *Elysia cornigera* (Heterobranchia: Plakobranchoidea) as distinct from the 'solar-powered' sea slug, *E. timida*. *Invertebr. Syst.* **25**, 477–489. (doi:10.1071/IS11026)
 22. Mujer CV, Andrews DL, Manhart JR, Pierce SK, Rumpho ME. 1996 Chloroplast genes are expressed during intracellular symbiotic association of *Vaucheria litorea* plastids with the sea slug *Elysia chlorotica*. *Proc. Natl Acad. Sci. USA* **93**, 12 333–12 338. (doi:10.1073/pnas.93.22.12333)
 23. Pelletreau KN, Weber APM, Weber KL, Rumpho ME. 2014 Lipid accumulation in the establishment of kleptoplasty in *Elysia chlorotica*. *PLoS ONE* **9**, e97477. (doi:10.1371/journal.pone.0097477.g012)
 24. Arrhenius A, Grönvall F, Scholze M, Backhaus T, Blanck H. 2004 Predictability of the mixture toxicity of 12 similarly acting congeneric inhibitors of photosystem II in marine periphyton and epipsammon communities. *Aquat. Toxicol.* **68**, 351–367. (doi:10.1016/j.aquatox.2004.04.002)
 25. Goldberg AL. 2003 Protein degradation and protection against misfolded or damaged proteins. *Nature* **426**, 895–899. (doi:10.1038/nature02263)
 26. Kirkin V, McEwan DG, Novak I, Dikic I. 2009 A role for ubiquitin in selective autophagy. *Mol. Cell* **34**, 259–269. (doi:10.1016/j.molcel.2009.04.026)
 27. Wooten MW, Geetha T, Babu JR, Seibenhener ML, Peng J, Cox N, Diaz-Meco M-T, Moscat J. 2008 Essential role of sequestosome 1/p62 in regulating accumulation of Lys⁶³-ubiquitinated proteins. *J. Biol. Chem.* **283**, 6783–6789. (doi:10.1074/jbc.M709496200)
 28. Balaban RS, Nemoto S, Finkel T. 2005 Mitochondria, oxidants, and aging. *Cell* **120**, 483–495. (doi:10.1016/j.cell.2005.02.001)
 29. Wägele H, Ballesteros M, Avila C. 2006 Defensive glandular structures in opisthobranch molluscs: from histology to ecology. *Oceanogr. Mar. Biol.* **44**, 197–276. (doi:10.1201/9781420006391.ch5)
 30. Carter WO, Narayanan PK, Robinson JP. 1994 Intracellular hydrogen peroxide and superoxide anion detection in endothelial cells. *J. Leukoc. Biol.* **55**, 253–258.
 31. Cueto M, D'Croz L, Maté JL, San-Martín A, Darias J. 2005 Elysiapyrones from *Elysia diomedea*. Do such metabolites evidence an enzymatically assisted electrocyclization cascade for the biosynthesis of their bicycle[4.2.0]octane core? *Org. Lett.* **7**, 415–418. (doi:10.1021/ol0477428)
 32. Färber A, Young AJ, Ruban AV, Horton P, Jahns P. 1997 Dynamics of xanthophyll-cycle activity in different antenna subcomplexes in the photosynthetic membranes of higher plants: (the relationship between zeaxanthin conversion and nonphotochemical fluorescence quenching). *Plant Phys.* **115**, 1609–1618. (doi:10.1104/pp.115.4.1609)
 33. Wägele H, Johnsen G. 2001 Observations on the histology and photosynthetic performance of 'solar-powered' opisthobranchs (Mollusca, Gastropoda, Opisthobranchia) containing symbiotic chloroplasts or zooxanthellae. *Org. Divers. Evol.* **1**, 193–210. (doi:10.1078/1439-6092-00016)
 34. Schindelin J *et al.* 2012 Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682. (doi:10.1038/nmeth.2019)
 35. R Core Development Team 2014 R: a language and environment for statistical computing. See <http://www.R-project.org>.
 36. Shapiro SS, Wilk MB. 1965 An analysis of variance test for normality (complete samples). *Biometrika* **52**, 591–611. (doi:10.1093/biomet/52.3-4.591)
 37. Mann HB, Whitney DR. 1947 On a test of whether one of two random variables is stochastically larger than the other. *Ann. Math. Stat.* **1**, 50–60. (doi:10.1214/aoms/1177730491)
 38. FastQC 2014 A quality control tool for high throughput sequence data. See <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
 39. Bolger AM, Lohse M, Usadel B. 2014 TRIMMOMATIC: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120. (doi:10.1093/bioinformatics/btu170)
 40. Grabherr MG *et al.* 2011 Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* **29**, 644–652. (doi:10.1038/nbt.1883)
 41. Haas BJ *et al.* 2013 *De novo* transcript sequence reconstruction from RNA-seq using the TRINITY platform for reference generation and analysis. *Nat. Protoc.* **8**, 1494–1512. (doi:10.1038/nprot.2013.084)
 42. Li B, Dewey CN. 2011 RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinform.* **4**, 323. (doi:10.1186/1471-2105-12-323)
 43. Langmead B, Trapnell C, Pop M, Salzberg SL. 2009 Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25. (doi:10.1186/gb-2009-10-3-r25)
 44. Robinson MD, McCarthy DJ, Smyth GK. 2010 edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140. (doi:10.1093/bioinformatics/btp616)
 45. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997 Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389–402. (doi:10.1093/nar/25.17.3389)
 46. Pruitt KD, Tatusova T, Brown GR, Maglott DR. 2010 NCBI Reference Sequences (RefSeq): current status, new features and genome annotation policy. *Nucleic Acids Res.* **40**, D130–D135. (doi:10.1093/nar/gkr1079)
 47. Zhang G *et al.* 2012 The oyster genome reveals stress adaptation and complexity of shell formation. *Nature* **490**, 49–54. (doi:10.1038/nature11413)
 48. Simakov O *et al.* 2013 Insights into bilaterian evolution from three spiralian genomes. *Nature* **493**, 526–531. (doi:10.1038/nature11696)

49. Boeckmann B *et al.* 2003 The SWISS-PROT protein knowledgebase and its supplement TrEMBL in 2003. *Nucleic Acids Res.* **31**, 365–370. (doi:10.1093/nar/gkg095)
50. Tatusov RL, Koonin EV, Lipman DJ. 1997 A genomic perspective on protein families. *Science* **278**, 631–637. (doi:10.1126/science.278.5338.631)
51. Koonin EV *et al.* 2004 A comprehensive evolutionary classification of proteins encoded in complete eukaryotic genomes. *Genome Biol.* **5**, R7. (doi:10.1186/gb-2004-5-2-r7)
52. Jensen LJ, Julien P, Kuhn M, von Mering C, Muller J, Doerks T, Bork P. 2008 eggNOG: automated construction and annotation of orthologous groups of genes. *Nucleic Acids Res.* **36**, D250–D254. (doi:10.1093/nar/gkm796)
53. Powell S *et al.* 2014 eggNOG v4.0: nested orthology inference across 3686 organisms. *Nucleic Acids Res.* **42**, D231–D239. (doi:10.1093/nar/gkt1253)
54. Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M. 1999 KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **27**, 29–34. (doi:10.1093/nar/28.1.27)
55. Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M. 2007 KAAS: an automatic genome annotation and pathway reconstruction server. *Nucleic Acids Res.* **35**, W182–W185. (doi:10.1093/nar/gkm321)
56. Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. 2012 Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinform.* **13**, 134. (doi:10.1186/1471-2105-13-134)
57. Pfaffl MW. 2001 A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **29**, e45. (doi:10.1093/nar/29.9.e45)