

POSTER PRESENTATION

Open Access

Multifunctional central pattern generator controlling walking and paw shaking

Brian Bondy¹, Alexander Klishko², Boris Prilutsky², Gennady Cymbalyuk^{1*}

From The Twenty Third Annual Computational Neuroscience Meeting: CNS*2014
Québec City, Canada. 26-31 July 2014

Central pattern generators (CPGs) are oscillatory neuronal networks controlling rhythmic motor tasks such as breathing and walking. A multifunctional CPG can produce multiple patterns, e.g. patterns with different periods [1-5]. Here, we investigate whether a pair of cat behaviors – walking and paw shaking – could be controlled by a single multifunctional CPG exhibiting multi-stability of oscillatory regimes. In experiments, both behaviors can be elicited in a spinalized cat, and there is evidence that the same circuitry is used for both rhythms [2,3]. We present a parsimonious model of a half-center oscillator composed of two mutually inhibitory neurons. These cells contains two slowly inactivating inward

currents, a persistent Na^+ current (I_{NaP}) and a low voltage activated Ca^{++} current (I_{CaLVA}). The dynamics of the multifunctional CPG is based on that the I_{CaLVA} inactivates much slower than I_{NaP} and at the more hyperpolarized membrane potentials. Here, we demonstrate the co-existence of two rhythms (Figure 1). At first, the model demonstrates walking pattern. A switch from a slow, 1-2 Hz walking rhythm to fast, 7-10 Hz paw shake rhythm was elicited by a pulse of conductance of excitatory current delivered to extensor and flexor neurons. Then, a switch back to walking was triggered by a shorter pulse of conductance of inhibitory current delivered to the extensor neuron.




Figure 1 Two mutually inhibitory interneurons, IntE (Extensor Interneuron) and IntF (Flexor Interneuron) produce alternating bursting activity at approximately 1.6 Hz representing walking pattern. A switch to paw shaking is executed by a pulse of excitatory conductance delivered to both cells for 1 second. The paw shake rhythm is represented by a 9 Hz bursting regime. An inhibitory conductance activated for .1 second in IntF causes a large rebound burst and a fast transition back to the walking rhythm.

* Correspondence: gcymbalyuk@gsu.edu

¹Neuroscience Institute, Georgia State University, Atlanta, Georgia, 30302, USA

Full list of author information is available at the end of the article

The CPG model was also incorporated into a neuro-mechanical model of a cat hindlimb in the AnimatLab environment [6]. The model provides a cellular mechanism of multifunctional CPG operation.

Acknowledgments

The authors acknowledge support by the NSF PHY-0750456 to Gennady Cymbalyuk and by NIH P01 HD32571, R01 EB012855, and R01 NS048844 and by the Center for Human Movement Studies at GATech to Boris Prilutsky.

Authors' details

¹Neuroscience Institute, Georgia State University, Atlanta, Georgia, 30302, USA. ²School of Applied Physiology, Georgia Institute of Technology, Atlanta, Georgia, 30332, USA.

Published: 21 July 2014

References

1. Berkowitz A: Both shared and specialized spinal circuitry for scratching and swimming in turtles. *J Comp Physiol A* 2002, **188**(3):225-234.
2. Carter MC, Smith JL: Simultaneous control of two rhythmical behaviors. II. Hindlimb walking with paw-shake response in spinal cat. *J of Neurophysiol* 1986, **56**(1):184-195.
3. Frigon A, Gossard JP: Evidence for specialized rhythm-generating mechanisms in the adult mammalian spinal cord. *J of Neurosci* 2010, **30**(20):7061-7071.
4. Venugopal S, Travers J, Terman D: A computational model for motor pattern switching between taste-induced ingestion and rejection oromotor behaviors. *J Comput Neurosci* 2007, **22**(2):223-238.
5. Manor Y, Nadim F: Synaptic depression mediates bistability in neuronal networks with recurrent inhibitory connectivity. *J Neurosci* 2001, **21**(23):9460-9470.
6. Klishko AN, Cofer D, Cymbalyuk G, Edwards DH, Prilutsky BI: Paw-shake response and locomotion: can one CPG generate two different rhythmic behaviors? *BMC Neuroscience* 2012, **13**(Suppl 1):P70.

doi:10.1186/1471-2202-15-S1-P181

Cite this article as: Bondy et al.: Multifunctional central pattern generator controlling walking and paw shaking. *BMC Neuroscience* 2014 **15**(Suppl 1):P181.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

