

CASE REPORT

Open Access



A case of episcleral tattooing – an emerging body modification trend

James Brodie*, Husam El Galhud and Adam Bates

Abstract

Background: In 2007 an article was published describing the first forays into the practice of episcleral tattooing. Currently only a handful of people worldwide have undergone this procedure, whereby a needle is used to inject dye under the bulbar conjunctiva. To date there have been no previous reports of the risks and complications of this emerging practice in the medical literature.

We present a case involving a complication that arose in one of the few people in Britain to have undergone episcleral tattooing for cosmetic purposes.

Case presentation: A 43-year-old Caucasian man presented to the eye casualty clinic with red, lumpy conjunctivae bilaterally, having undergone episcleral tattooing 7 weeks previously. On examination there were 3 distinct areas of conjunctival swelling in each eye, representing a total of 6 injection sites. No other gross abnormalities were identified. The clinical picture remained unchanged 6 months on, apart from a degree of fading of the conjunctival dye. He will remain under our care to ensure that any further complications such as granulomatous inflammation are managed and documented.

Conclusion: Episcleral tattooing is carried out by individuals with no medical training. The short-term complications reported so far include: headaches, severe photophobia, persistent foreign body sensation, and migration of ink staining. More serious short-term risks such as infection, globe penetration, and peri-ocular haemorrhage could occur. For now we can only speculate as to the long-term consequences, but these may include carcinogenic change or granulomatous inflammation. We feel that the potential risks of the procedure should be communicated more widely to those body modification practitioners undertaking it. This practice could result in more serious presentations to acute eye services in the future.

Background

In 2007 the Body Modification Ezine published an article describing the first instances of episcleral tattooing [1]. At present only a handful of people worldwide have undergone episcleral tattooing, whereby dye is injected under the bulbar conjunctiva, for purely cosmetic purposes. There have been no previous reports of the risks and complications of this emerging practice in the medical literature.

We present a case involving a complication in one of the few people in Britain to have undergone episcleral tattooing.

Case presentation

A 43-year-old man presented with bright red, lumpy conjunctivae bilaterally (see Fig. 1). 7 weeks previously he had undergone episcleral tattooing and his complaint was that the conjunctival lumps had not subsided. In this case a mixture of two dermal tattoo dyes (C.I Pigment Red 210 and C.I Pigment Blue 15) was injected under the conjunctiva at 3 sites in each eye. Visual acuity was 6/4 in both eyes and he reported no visual symptoms or discomfort.

On slit lamp examination there were 3 distinct areas of conjunctival swelling in each eye, representing a total of 6 injection sites (Fig. 2). Apart from this aesthetically poor, persistent lumpy appearance, no other gross abnormalities were identified. Fundal examination and intraocular pressure measurements were also normal. The clinical picture remained unchanged 6 months on, apart from a degree of fading of the conjunctival dye (Fig. 3).

* Correspondence: jtbrodie@doctors.org.uk
Ophthalmology Department, Maidstone Hospital, Hermitage Lane,
Maidstone, Kent ME16 9QQ, UK



Fig. 1 Bilateral episcleral tattoos

He will remain under our care to ensure that any further complications such as granulomatous inflammation are managed and documented.

Conclusion

The number of body modification procedures carried out annually is increasing [2] and new trends are pushing the boundaries of “extreme” in many directions. Procedures such as episcleral tattooing are carried out by individuals with no medical training, who are taking it upon themselves to pioneer an untested practice through trial and error.

The short-term complications reported so far have included: headaches, severe photophobia, persistent foreign body sensation, and migration of ink staining to nearby tissues [3, 4]. Episcleral tattooing by a non-ophthalmic trained user without the use of a surgical microscope carries the risk of globe penetration, traumatic cataract, retinal detachment and endophthalmitis. In addition, tattoos carry the risk of blindness from tattoo-induced uveitis [5–8]. If distant skin tattoos can induce uveal sensitization to the dyes




Fig. 3 Left eye showing fading of conjunctival dye at 6 months follow up

in the tattoo, placing the dye under the conjunctiva may result in more severe or more frequent uveitis. Finally, the presence of chronic granuloma can lead in the long term to scleral thinning or malignancy [9].

We feel that the potential risks of the procedure should be communicated more widely to those body modification practitioners undertaking it. This practice could result in more serious presentations to acute eye services in the coming years.

Consent

Written informed consent was obtained from the patient for publication of this Case report and all of the accompanying images. A copy of the written consent is available for review by the Editor of this journal.

Competing interests

The authors declare that they have no competing interests.

Author’s contributions

JB clinical review of patient, drafting and editing of manuscript. HEG clinical review of patient, photos. AB editing of manuscript. All authors read and approved the final manuscript.

Received: 1 September 2014 Accepted: 31 July 2015

Published online: 08 August 2015

References

1. Larratt S. Three Blind Mice. <http://news.bme.com/2007/07/02/three-blind-mice/> (accessed 3 November 2013).
2. Braverman PK. Body art: piercing, tattooing, and scarification. *Adolesc Med Clin.* 2006;17(3):505–19.
3. Positano R. Don’t Turn Blind Eye to Eye-Tattoo Risks. <http://nypost.com/2007/07/31/dont-turn-blind-eye-to-eye-tattoo-risks/> (accessed 5 November 2013).
4. Carmina L. A Look at Eyeball Tattoos and Extreme Body Modifications. http://www.huffingtonpost.com/la-carmina/eyeball-tattoos-extreme-b_b_3031043.html (accessed 5 November 2013).
5. Mansour AM, Chan CC. Tattoo associated uveitis (letter). *Am J Ophthalmol.* 2015;159:408–9.
6. Ostheimer TA, Burkholder BM, Leung TG, Butler NJ, Dunn JP, Thorne JE. Tattoo-associated uveitis. *Am J Ophthalmol.* 2014;158:637–43.




Fig. 2 Left eye showing 3 distinct lumps at injection sites

7. Mansour AM, Chan CC. Recurrent uveitis preceded by swelling of skin tattoos. *Am J Ophthalmol.* 1991;111:515–6.
8. Jacob J, Van Calster J. Skin tattoos and the development of uveitis. *Acta Ophthalmol.* 2013;91:650–2.
9. Multhoff G, Molls M, Radons J. Chronic inflammation in cancer development. *Front Immunol.* 2012;2:98.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

