

Letter to the editor:

METABOLIC DYSHOMEOSTASIS BY ORGANOPHOSPHATE INSECTICIDES: INSIGHTS FROM EXPERIMENTAL AND HUMAN STUDIES

Apurva Kumar Ramesh Joshi*, Bindhu Omana Sukumaran

Department of Biochemistry, School of Sciences, Jain University, Bangalore, Karnataka, India 560041

* Corresponding author: Department of Biochemistry, School of Sciences, Jain University, Jayanagar 3rd Block, Bangalore, Karnataka, India 560041, Tel: +918019236873; E-mail: r.apurvakumar@jainuniversity.ac.in

<http://dx.doi.org/10.17179/excli2019-1492>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

Dear Editor,

Organophosphate insecticides (OPI), derived from phosphoric, phosphonic or phosphinic acids, find application as agents for controlling insect pest populations. OPIs elicit their characteristic toxicity by phosphorylating and inhibiting the enzyme, acetylcholinesterase (AChE). Cholinergic stress resulting from overstimulation of nicotinic- and muscarinic acetylcholine receptors is the chief mechanism of acute toxicity of OPI (Fukuto, 1990; Sogorb and Vilanova, 2002; Abou-Donia, 2003; Costa, 2006). The ubiquitous nature of AChE and its conserved physiological role in the regulation of neurotransmission means that non-target animals (including humans) are at risk of adverse outcomes in the event of exposure to OPI. Neurotoxicity, characterized by cholinergic and non-cholinergic outcomes, is the most studied aspect of OPI toxicity. However, it is now unequivocally recognized that the toxicity of OPIs goes beyond the realm of neurotoxicity.

A large number of animal studies explicitly demonstrate that OPIs have the propensity to cause hyperglycemia, perturbations in carbohydrate metabolism and endocrine dysregulations. Repeated exposure to OPI precipitates insulin resistance (studies listed in Table 1), a key component of metabolic syndrome. Extrapolating the outcomes of animal experimentation to the human situation is a challenging task. Experimental studies often employ doses much higher than doses of environmental relevance. However, several studies clearly (listed in Table 2) establish that OPI exposure elicits metabolic dysregulation in human subjects. A recent study demonstrates that the incidence of diabetes among Thai farmers positively correlates with OPIs such as chlorpyrifos, dicofol, dichlorvos, ethyl-p-nitrophenyl, mevinphos, monocrotophos and methamidophos (Juntarawijit and Juntarawijit, 2018). Thus, it is evident that OPI exposure is a clear risk factor for metabolic dysregulations among those who are occupationally exposed. One has to take cognizance of the fact that levels of exposure to OPI among occupationally exposed people are likely to be higher than the general population. However, a recent study reveals that glycated hemoglobin levels correlate with plasma levels of OPI (due to environmental exposure), but not with AChE activity (Velmurugan et al., 2017). This indicates that low-level OPI exposure may cause metabolic dysregulations.

Hence, we believe that further studies are needed to evaluate the effects of low-level, chronic non-occupational exposure to OPI on metabolic health.

Table 1: Experimental studies reporting metabolic dysregulations caused by organophosphate insecticides in rodent models

Key findings	Reference
Long term exposure to chlorpyrifos reduces insulin sensitivity and causes fat gain in rats. Chlorpyrifos-treated rats also suffered from disruption of the gut barrier, alteration in gut microflora profile and low-grade inflammation.	Liang et al., 2019
Diazinon treatment was associated with an increase in blood glucose, insulin and vaspin levels, and glucose intolerance in rats. Increased expression of PTEN and FOXO1 was observed in the livers diazinon treated rats.	Salek-Maghsoudi et al., 2019
Repeated administration of diazinon for 4 weeks caused increase in blood glucose, creatinine, urea, percentage of lymphocytes, dyslipidemia, increased circulating aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase and lactate dehydrogenase activities in the adult and aged rats compared to respective controls.	Yousefizadeh et al., 2019
Methyl-parathion administration to rats causes elevated plasma glucose and creatinine levels along with diminished urinary flow rate and increased urinary excretion of glucose, phosphate, and albumin.	Fuentes-Delgado et al., 2018
Chronic exposure to monocrotophos causes hyperglycemia and glucose intolerance in mice. Intestinal metatranscriptomic and host metabolomic analyses revealed that short-chain fatty acids such as acetate produced by microbial degradation of organophosphates may induce gluconeogenesis.	Velmurugan et al., 2017
Exposure to acephate during pregnancy and lactation causes glucose intolerance in the offspring in rats.	Ribeiro et al., 2016
Augmentation of pancreatic beta cell functions precipitate as a way to counter glucose intolerance in rats subjected to chronic exposure to monocrotophos.	Nagaraju and Rajini, 2016
Repeated exposure to malathion caused increase in plasma glucose, insulin and glycated hemoglobin levels along with dyslipidemia in rats.	Lasram et al., 2015
Chronic exposure to monocrotophos causes elevated fasting glucose, hyperinsulinemia, hypercorticosteronemia, glucose intolerance, insulin resistance and increase in circulating triglyceride levels in rats. Further, tyrosine aminotransferase, glucose-6-phosphatase and glycogen phosphorylase were elevated in livers of treated rats along with loss of glycogen.	Nagaraju et al., 2015
Repeated exposure (8 weeks) to chlorpyrifos and profenofos causes hyperglycemia in rats.	Hamza et al., 2014
Repeated exposure to monocrotophos causes increase in blood glucose, dyslipidemia and cardiac toxicity in rats.	Velmurugan et al., 2013
Repeated exposure to malathion causes fasting hyperglycemia along with increase in phosphoenolpyruvate carboxykinase, glucose 6-phosphatase activity and tumor necrosis factor α levels in the liver.	Mostafalou et al., 2012
Hyperglycemia in rats subjected to a single dose of monocrotophos is attributable to adrenergic mechanisms, while hypercorticosteronemia is responsible for upregulation of liver tyrosine aminotransferase activity.	Joshi et al., 2012
Cholinergic antagonism prevents the hyperglycemic potential of a single dose of monocrotophos.	Joshi and Rajini, 2012
Acute exposure to chlorpyrifos causes hyperglycemia, hypercorticosteronemia and dyslipidemia in rats.	Acker and Nogueira, 2012
Acute exposure to malathion causes hyperglycemia and hepatic glycogen depletion in rats.	Lasram et al., 2008
Repeated exposure to dimethoate causes alteration in glucose homeostasis and oxidative impairments in the pancreas in rats.	Kamath et al., 2008

Key findings	Reference
A single dose of acephate causes reversible hyperglycemia in rats, with concomitant hypercorticosteronemia and increased activity of hepatic gluconeogenesis enzymes (tyrosine aminotransferase and glucose-6-phosphatase).	Joshi and Rajini, 2009
Sub-chronic exposure to malathion leads to hyperglycemia and up-regulation of hepatic glycogen phosphorylase and phosphoenolpyruvate carboxykinase activity in rats.	Abdollahi et al., 2004

Table 2: Studies that demonstrate the link between exposure to organophosphate insecticides and metabolic dysregulations in human subjects

Key findings	Reference
A prospective study of cases involving pesticide self-poisoning conducted in parallel in Sri Lanka and Bangladesh showed that OPI poisoning is associated with acute dysregulations in glucose homeostasis that are related to changes in insulin action and secretion.	Gifford et al., 2019
A survey performed in villages of Vadapalanji Panchayat (a southern part of India) revealed that OPI constituted nearly 50 % of used pesticides. A survey of over 3000 participants revealed a prevalence of diabetes (18 %). Plasma organophosphate residues correlated positively with HbA1c. HbA1c values did not correlate with plasma cholinesterase levels.	Velmurugan et al., 2017
A retrospective observational study of 184 non-diabetic patients with organophosphate poisoning suggests that hyperglycemia at presentation is associated with in-hospital mortality rates.	Moon et al., 2016
A study of 50 individuals admitted to Osmania hospital in Hyderabad (India) with OPI poisoning revealed that 38 % of patients had hyperglycemia. Hyperglycemia at admission correlated with suppression of pseudocholinesterase activity. Random blood sugar levels of > 200 mg/dl at admission and pseudocholinesterase levels below 1000 U/L are reported as reliable parameters to predict mortality and ventilator requirement in OPI poisoning.	Rao and Raju, 2016
A study of 71 cases of OPI poisoning revealed that blood glucose levels in subjects were higher (144.7 ± 85.3 mg/dL) than the normal range (70–110 mg/dl). More importantly, a subpopulation of patients exhibiting intermediate syndrome (n=11) exhibited much higher blood glucose levels (186.63 ± 57.31 v/s 138.55 ± 71.34 mg/dL) and severe suppression of serum cholinesterase activity (465.11 ± 302.63 v/s 2651.72 ± 1266.68 U/L; normal range – 3,600–12,900 U/L) than the patients without intermediate syndrome (n=56).	Çolak et al., 2014
Transient glycosuria with or without hyperglycemia was reported in a significant number of patients admitted due to OPI poisoning	Sudhir et al., 2013
A comparative cross-sectional study involving a non-diabetic farmer cohort (n=98) and age-matched control group (n=90) revealed that blood malathion (0.0746 ± 0.01404 v/s 0.0031 ± 0.0006 mg/L), fasting blood glucose (109.1 ± 20.6 v/s 89.8 ± 10.2 mg/dL) and insulin (12.3 ± 4.6 v/s 5.2 ± 1.6 µU/mL) levels were markedly higher in farmer group as compared to the control group. A positive correlation was observed between malathion blood concentration, waist circumference and insulin resistance.	Raafat et al., 2012
91 cases of OPI poisoning were classified into grade 0, 1, 2, and 3. Blood glucose was elevated and serum cholinesterase activity was suppressed in grade 1, 2 and 3 patients. The severity of hyperglycemia was related to cholinesterase suppression.	Amanvermez et al., 2010
A study, involving more than 33000 pesticide applicators, designed to investigate the relationship between lifetime exposure to specific agricultural pesticides and diabetes incidence among the applicators, revealed that	Montgomery et al., 2008

Key findings	Reference
chlorpyrifos, coumaphos, diazinon, dichlorvos, phorate, terbufos, and trichlorfon were associated with increased odds of diabetes, with chlorpyrifos, diazinon, and trichlorfon being dose-dependent.	
Hyperglycemia was observed in 982 out of 2708 OPI poisoning cases studied in Pakistan between 2001 and 2007.	Ather et al., 2008
A retrospective study of 220 patients hospitalized in Turkey between the years 1995 and 2004 revealed elevated blood glucose levels in patients (145 ±68 mg/dL) compared to the normal range (70-110 mg/dL).	Yurumez et al., 2007
Elevated levels of glucose, amylase and immunoreactive trypsin levels were observed in children exhibiting acute pancreatitis due to OPI and carbamate poisoning.	Weizman and Sofer, 1992
A case report of 2 fatal OPI poisoning revealed that subjects were associated with severe hyperglycemia, metabolic acidaemia and moderate hyperkalemia.	Hui, 1983
A 32 year old male patient with OPI poisoning developed hyperglycemia, glycosuria and ketonuria soon after hospitalization. Intervention with insulin was required to treat hyperglycemia.	Moore and James, 1981

Acknowledgements

Authors are thankful to Jain University, Bangalore for the support.

Conflict of interest

The authors declare no conflict of interest.

REFERENCES

- Abdollahi M, Donyavi M, Pournourmohammadi S, Saadat M. Hyperglycemia associated with increased hepatic glycogen phosphorylase and phosphoenolpyruvate carboxykinase in rats following subchronic exposure to malathion. *Comp Biochem Physiol Part C Toxicol Pharmacol*. 2004;137:343–7.
- Abou-Donia MB. Organophosphorus ester-induced chronic neurotoxicity. *Arch Environ Health*. 2003;58:484–97.
- Acker CI, Nogueira CW. Chlorpyrifos acute exposure induces hyperglycemia and hyperlipidemia in rats. *Chemosphere*. 2012;89:602–8.
- Amanvermez R, Baydın A, Yardan T, Başol N, Günay M. Emergency laboratory abnormalities in suicidal patients with acute organophosphate poisoning. *Turkish J Biochem*. 2010;35:29–34.
- Ather NA, Ara J, Khan EA, Sattar RA, Durrani R. Acute organophosphate insecticide poisoning. *J Surg Pakistan*. 2008;13:71–4.
- Çolak Ş, Erdoğan MÖ, Baydın A, Afacan MA, Kati C, Duran L. Epidemiology of organophosphate intoxication and predictors of intermediate syndrome. *Turkish J Med Sci*. 2014;44:279–82.
- Costa LG. Current issues in organophosphate toxicology. *Clin Chim Acta*. 2006;366:1–13.
- Fuentes-Delgado VH, Martínez-Saldaña MC, Rodríguez-Vázquez ML, Reyes-Romero MA, Reyes-Sánchez JL, Jaramillo-Juárez F. Renal damage induced by the pesticide methyl parathion in male Wistar rats. *J Toxicol Environ Health Part A*. 2018;81:130–41.
- Fukuto TR. Mechanism of action of organophosphorus and carbamate insecticides. *Environ Health Perspect*. 1990;87:245–54.
- Gifford RM, Chathuranga U, Lamb T, Verma V, Sattar MA, Thompson A, et al. Short-term glucose dysregulation following acute poisoning with organophosphorus insecticides but not herbicides, carbamate or pyrethroid insecticides in South Asia. *Clin Toxicol*. 2019;57:254–64.
- Hamza RZMM, Diab AE, El-Aziz E-SAA. Hyperglycemic effect of chlorpyrifos, profenofos and possible ameliorative role of propolis and ginseng. *Sci Agric PSCI Publ*. 2014;9–14.
- Hui KS. Metabolic disturbances in organophosphate insecticide poisoning. *Arch Pathol Lab Med*. 1983;107:154.

- Joshi AKR, Rajini PS. Reversible hyperglycemia in rats following acute exposure to acephate, an organophosphorus insecticide: role of gluconeogenesis. *Toxicology*. 2009;257:40–5.
- Joshi AKR, Rajini PS. Hyperglycemic and stressogenic effects of monocrotophos in rats: evidence for the involvement of acetylcholinesterase inhibition. *Exp Toxicol Pathol*. 2012;64:115–20.
- Joshi AKR, Nagaraju R, Rajini PS. Insights into the mechanisms mediating hyperglycemic and stressogenic outcomes in rats treated with monocrotophos, an organophosphorus insecticide. *Toxicology*. 2012; 294:9–16.
- Juntarawijit C, Juntarawijit Y. Association between diabetes and pesticides: a case-control study among Thai farmers. *Environ Health Prev Med*. 2018;23:3.
- Kamath V, Joshi AKR, Rajini PS. Dimethoate induced biochemical perturbations in rat pancreas and its attenuation by cashew nut skin extract. *Pestic Biochem Physiol*. 2008;90:58–65.
- Lasram MM, Annabi AB, Rezg R, Elj N, Slimen S, Kamoun A, et al. Effect of short-time malathion administration on glucose homeostasis in Wistar rat. *Pestic Biochem Physiol*. 2008;92:114–9.
- Lasram MM, Bouzid K, Douib IB, Annabi A, El Elj N, El Fazaa S, et al. Lipid metabolism disturbances contribute to insulin resistance and decrease insulin sensitivity by malathion exposure in Wistar rat. *Drug Chem Toxicol*. 2015;38:227–34.
- Liang Y, Zhan J, Liu D, Luo M, Han J, Liu X, et al. Organophosphorus pesticide chlorpyrifos intake promotes obesity and insulin resistance through impacting gut and gut microbiota. *Microbiome*. 2019;7:19.
- Montgomery MP, Kamel F, Saldana TM, Alavanja MCR, Sandler DP. Incident diabetes and pesticide exposure among licensed pesticide applicators: Agricultural Health Study, 1993-2003. *Am J Epidemiol*. 2008;167:1235–46.
- Moon JM, Chun BJ, Cho YS. Hyperglycemia at presentation is associated with in hospital mortality in non-diabetic patient with organophosphate poisoning. *Clin Toxicol*. 2016;54:252–8.
- Moore PG, James OF. Acute pancreatitis induced by acute organophosphate poisoning. *Postgrad Med J*. 1981;57:660–2.
- Mostafalou S, Eghbal MA, Nili-Ahmadabadi A, Baeeri M, Abdollahi M. Biochemical evidence on the potential role of organophosphates in hepatic glucose metabolism toward insulin resistance through inflammatory signaling and free radical pathways. *Toxicol Ind Health*. 2012;28:840–51.
- Nagaraju R, Rajini P. Adaptive response of rat pancreatic β -cells to insulin resistance induced by monocrotophos: Biochemical evidence. *Pestic Biochem Physiol*. 2016;134:39–48.
- Nagaraju R, Joshi AKR, Rajini PS. Organophosphorus insecticide, monocrotophos, possesses the propensity to induce insulin resistance in rats on chronic exposure. *J Diabetes*. 2015;7:47–59.
- Raafat N, Abass MA, Salem HM. Malathion exposure and insulin resistance among a group of farmers in Al-Sharkia governorate. *Clin Biochem*. 2012;45: 1591–5.
- Rao R, Raju G. Random blood sugar levels and pseudocholinesterase levels their relevance in organophosphorus compound poisoning. *Int J Community Med Public Health*. 2016;3:2757–61.
- Ribeiro TA, Prates KV, Pavanello A, Malta A, Tôfolo LP, Martins IP, et al. Acephate exposure during a perinatal life program to type 2 diabetes. *Toxicology*. 2016;372:12–21.
- Salek-Maghsoudi A, Hassani S, Momtaz S, Shadboorestan A, Ganjali MR, Ghahremani MH, et al. Biochemical and molecular evidence on the role of vaspin in early detection of the insulin resistance in a rat model of high-fat diet and use of diazinon. *Toxicology*. 2019;411:1–14.
- Sogorb MA, Vilanova E. Enzymes involved in the detoxification of organophosphorus, carbamate and pyrethroid insecticides through hydrolysis. *Toxicol Lett*. 2002;128:215–28.
- Sudhir U, Chandrashekar, Pai R, Sunil H, Rao M, Kempgowda P. Glycemic changes in acute anticholinesterase insecticide poisoning. *West Lond Med J*. 2013;5:27–33.
- Velmurugan G, Venkatesh Babu DD, Ramasamy S. Prolonged monocrotophos intake induces cardiac oxidative stress and myocardial damage in rats. *Toxicology*. 2013;307:103–8.
- Velmurugan G, Ramprasath T, Swaminathan K, Mithieux G, Rajendhran J, Dhivakar M, et al. Gut microbial degradation of organophosphate insecticides-induces glucose intolerance via gluconeogenesis. *Genome Biol*. 2017;18:8.

Weizman Z, Sofer S. Acute pancreatitis in children with anticholinesterase insecticide intoxication. *Pediatrics*. 1992;90:204–6.

Yousefizadeh S, Farkhondeh T, Samarghandian S. Age-related diazinon toxicity impact on blood glucose, lipid profile and selected biochemical indices in male rats. *Curr Aging Sci*. 2019, epub ahead of print.

Yurumez Y, Durukan P, Yavuz Y, Ikizceli I, Avsarogullari L, Ozkan S, et al. Acute organophosphate poisoning in university hospital emergency room patients. *Intern Med*. 2007;46:965–9.