

ADOPTED: 22 January 2019

doi: 10.2903/j.efsa.2019.5602

Safety and efficacy of L-threonine produced by fermentation with *Corynebacterium glutamicum* [REDACTED] for all animal species

EFSA Panel on Additives and Products or Substances used in Animal Feed (FEEDAP), Vasileios Bampidis, Giovanna Azimonti, Maria de Lourdes Bastos, Henrik Christensen, Birgit Dusemund, Maryline Kouba, Mojca Kos Durjava, Marta López-Alonso, Secundino López Puente, Francesca Marcon, Baltasar Mayo, Alena Pechová, Mariana Petkova, Yolanda Sanz, Roberto Edoardo Villa, Ruud Woutersen, Lucio Costa, Noël Dierick, Gerhard Flachowsky, Alberto Mantovani, Robert John Wallace, Jordi Tarrés-Call and Fernando Ramos

Abstract

Following a request from the European Commission, the Panel on Additives and Products or Substances used in Animal Feed (FEEDAP) was asked to deliver a scientific opinion on L-threonine produced by fermentation with *Corynebacterium glutamicum* [REDACTED] when used as a nutritional additive in feed and water for drinking for all animal species and categories. The product under assessment is L-threonine produced by fermentation with a [REDACTED] strain of *C. glutamicum* ([REDACTED]). L-Threonine produced by *C. glutamicum* [REDACTED] is considered safe for the target species when supplemented in appropriate amounts to the diet. The FEEDAP Panel has concerns on the safety of the simultaneous oral administration of L-threonine via water for drinking and feed. L-Threonine produced using *C. glutamicum* [REDACTED] is safe for the consumer. The additive is not a skin or eye irritant and is not a skin sensitiser. Although the workers can be exposed by inhalation, the results of an acute inhalation study showed that risk of adverse effects by inhalation is low. L-Threonine produced using *C. glutamicum* [REDACTED] is safe for the environment. The product under assessment is considered an efficacious source of the amino acid L-threonine for all animal species. For L-threonine to be as efficacious in ruminants as in non-ruminant species, it requires protection against degradation in the rumen.

© 2019 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

Keywords: nutritional additive, amino acid, L-threonine, safety, efficacy, *Corynebacterium glutamicum*

Requestor: European Commission

Question number: EFSA-Q-2018-00506

Correspondence: feedap@efsa.europa.eu

Panel members: Giovanna Azimonti, Vasileios Bampidis, Maria de Lourdes Bastos, Henrik Christensen, Birgit Dusemund, Maryline Kouba, Mojca Kos Durjava, Marta López-Alonso, Secundino López Puente, Francesca Marcon, Baltasar Mayo, Alena Pechová, Mariana Petkova, Fernando Ramos, Yolanda Sanz, Roberto Edoardo Villa and Ruud Woutersen.

Acknowledgements: The EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed) wishes to thank the following for the support provided to this scientific output: Montserrat Anguita, Jaume Galobart and Paola Manini.

Legal notice: Relevant information or parts of this scientific output have been blackened in accordance with the confidentiality requests formulated by the applicant pending a decision thereon by the European Commission. The full output has been shared with the European Commission, EU Member States and the applicant. The blackening will be subject to review once the decision on the confidentiality requests is adopted by the European Commission.

Suggested citation: EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), Bampidis V, Azimonti G, Bastos ML, Christensen H, Dusemund B, Kouba M, Kos Durjava M, López-Alonso M, López Puente S, Marcon F, Mayo B, Pechová A, Petkova M, Sanz Y, Villa RE, Woutersen R, Costa L, Dierick N, Flachowsky G, Mantovani A, Wallace RJ, Tarrés-Call J and Ramos F, 2019. Scientific Opinion on the safety and efficacy of L-threonine produced by fermentation with *Corynebacterium glutamicum* [REDACTED] for all animal species. EFSA Journal 2019;17(2):5602, 13 pp. <https://doi.org/10.2903/j.efsa.2019.5602>

ISSN: 1831-4732

© 2019 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs License](https://creativecommons.org/licenses/by-nd/4.0/), which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.



The EFSA Journal is a publication of the European Food Safety Authority, an agency of the European Union.



Table of contents

Abstract.....	1
1. Introduction.....	4
1.1. Background and Terms of Reference.....	4
1.2. Additional information.....	4
2. Data and methodologies.....	5
2.1. Data.....	5
2.2. Methodologies.....	5
3. Assessment.....	5
3.1. Characterisation.....	5
3.1.1. Characterisation of the production organism.....	5
3.1.2. Manufacturing process.....	6
3.1.3. Characterisation of the additive.....	6
3.1.3.1. Impurities.....	6
3.1.3.2. Physical characteristics.....	7
3.1.3.3. Stability and homogeneity.....	7
3.1.3.4. Physico-chemical incompatibilities.....	7
3.1.4. Conditions of use.....	8
3.2. Safety.....	8
3.2.1. Safety for the target species.....	8
3.2.2. Safety for the consumer.....	8
3.2.3. Safety for the user.....	8
3.2.3.1. Conclusions on safety for the user.....	8
3.2.4. Safety for the environment.....	8
3.3. Efficacy.....	9
3.4. Post-market monitoring.....	9
4. Conclusions.....	9
Documentation provided to EFSA.....	9
Chronology.....	9
References.....	10
Abbreviations.....	11
Annex A – Executive summary of the evaluation report of the European Union Reference Laboratory for feed additives on the methods of analysis for L-threonine produced by fermentation with <i>Corynebacterium glutamicum</i>	13

1. Introduction

1.1. Background and Terms of Reference

Regulation (EC) No 1831/2003¹ establishes the rules governing the Community authorisation of additives for use in animal nutrition. In particular, Article 4(1) of that Regulation lays down that any person seeking authorisation for a feed additive or for a new use of a feed additive shall submit an application in accordance with Article 7.

The European Commission received a request from CJ Europe GmbH² for authorisation of the product L-threonine, feed grade, produced by fermentation with *Corynebacterium glutamicum* [REDACTED] when used as a feed additive for all animal species (category: nutritional additives; functional group: amino acids, their salts and analogues).

According to Article 7(1) of Regulation (EC) No 1831/2003, the Commission forwarded the application to the European Food Safety Authority (EFSA) as an application under Article 4(1) (authorisation of a feed additive or new use of a feed additive). The particulars and documents in support of the application were considered valid by EFSA as of 20 April 2018.

According to Article 8 of Regulation (EC) No 1831/2003, EFSA, after verifying the particulars and documents submitted by the applicant, shall undertake an assessment in order to determine whether the feed additive complies with the conditions laid down in Article 5. EFSA shall deliver an opinion on the safety for the target animals, consumer, user and the environment and on the efficacy of the product L-threonine produced by fermentation with *C. glutamicum* [REDACTED] when used as a nutritional additive in feed and water for drinking for all animal species under the proposed conditions of use (see Section 3.1.4).

1.2. Additional information

L-Threonine produced by eight different strains of *Escherichia coli* (minimum content of 98% on dry matter basis) is currently authorised as a nutritional feed additive for use in all animal species.³ The product under assessment, L-threonine produced by the [REDACTED] strain of *C. glutamicum* [REDACTED] has not been previously authorised as feed additive in the European Union (EU).

L-Threonine is authorised for use in food,⁴ cosmetics⁵ and as a veterinary medicinal product.^{6,7}

L-Threonine is described in a monograph of the European Pharmacopoeia (MG 01/2008:1049) (Ph. Eur., 2016).

The Scientific Panel on Additives and Products or Substances used in Animal Feed (FEEDAP) issued ten opinions on the safety and efficacy of L-threonine produced by genetically modified strains of *E. coli* (EFSA FEEDAP Panel, 2013, 2014a,b,c,d, 2015a,b, 2016a,b, 2017a, 2018a).

The Joint FAO/WHO Expert Committee on Food Additives evaluated L-threonine as a food flavouring agent (JECFA; WHO, 2012).

¹ Regulation (EC) No 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition. OJ L 268, 18.10.2003, p. 29.

² CJ Europe GmbH, Ober der Roeth 4, 65824 Schwalbach am Taunus, Germany.

³ Commission implementing regulation (EC) 2016/1220 of 26 July 2016 concerning the authorisation of L-threonine produced by *Escherichia coli* as feed additive for all animal species. OJ L 201, 27.7.2016, p. 5.

⁴ Regulation (EU) No 609/2013 of the European Parliament and of the Council of 12 June 2013 on food intended for infants and young children, food for special medical purposes, and total diet replacement for weight control and repealing Council Directive 92/52/EEC, Commission Directives 96/8/EC, 1999/21/EC, 2006/125/EC and 2006/141/EC, Directive 2009/39/EC of the European Parliament and of the Council and Commission Regulations (EC) No 41/2009 and (EC) No 953/2009, OJ L 181, 29.6.2013, p. 35.

⁵ Commission Decision of 9 February 2006 amending Decision 96/335/EC establishing an inventory and a common nomenclature of ingredients employed in cosmetic products. OJ L 97, 5.4.2006, pp. 1–528.

⁶ Commission Regulation (EU) No 37/2010 of 22 December 2009 on pharmacologically active substances and their classification regarding maximum residue limits in foodstuffs of animal origin. OJ L 15, 20.1.2010, p. 1.

⁷ Regulation (EC) No 470/2009 of the European Parliament and of the Council of 6 May 2009 laying down Community procedures for the establishment of residue limits of pharmacologically active substances in foodstuffs of animal origin, repealing Council Regulation (EEC) No 2377/90 and amending Directive 2001/82/EC of the European Parliament and of the Council and Regulation (EC) No 726/2004 of the European Parliament and of the Council. OJ L 152, 16.6.2009, p. 11.

2. Data and methodologies

2.1. Data

The present assessment is based on data submitted by the applicant in the form of a technical dossier⁸ in support of the authorisation request for the use of L-threonine produced by fermentation with *C. glutamicum* [REDACTED] as a feed additive.

The FEEDAP Panel used the data provided by the applicant together with data from other sources, such as previous risk assessments by EFSA or other expert bodies, and experts' knowledge to deliver the present output.

EFSA has verified the European Union Reference Laboratory (EURL) report as it relates to the methods used for the control of the L-threonine produced by fermentation with *C. glutamicum* [REDACTED] in animal feed. The Executive Summary of the EURL report can be found in Annex A.⁹

2.2. Methodologies

The approach followed by the FEEDAP Panel to assess the safety and the efficacy of L-threonine produced by fermentation with *C. glutamicum* [REDACTED] is in line with the principles laid down in Regulation (EC) No 429/2008¹⁰ and the relevant guidance documents: Guidance on nutritional additives (EFSA FEEDAP Panel, 2012a), Guidance on the identity, characterisation and conditions of use of feed additives (EFSA FEEDAP Panel, 2017b), Guidance on the assessment of the safety of feed additives for the target species (EFSA FEEDAP Panel, 2017c), Guidance on the assessment of the safety of feed additives for the consumer (EFSA FEEDAP Panel, 2017d), Guidance on studies concerning the safety of use of the additive for users/workers (EFSA FEEDAP Panel, 2012b), Technical Guidance for assessing the safety of feed additives for the environment (EFSA, 2008) and Guidance on the characterisation of microorganisms used as feed additives or as production organisms (EFSA FEEDAP Panel, 2018b).

3. Assessment

The subject of the present assessment is L-threonine (minimum 98.5%) produced by fermentation with a [REDACTED] strain of *C. glutamicum* ([REDACTED]). It is intended to be used as a nutritional additive (functional group amino acids, their salts and analogues) to feed and water for drinking in all animal species and categories.

Under European Union (EU) conditions, L-threonine seems to be the second most limiting amino acid, after L-lysine, in pigs and the third most limiting, after the sulphur amino acids and L-lysine, in poultry.

3.1. Characterisation

3.1.1. Characterisation of the production organism

The additive under assessment is produced by a [REDACTED] strain of *Corynebacterium glutamicum*, [REDACTED]

C. glutamicum is considered suitable for qualified presumption of safety (QPS) approach when used for production purposes (EFSA BIOHAZ Panel, 2019) on condition that the production strain is free from possible antibiotic resistance.

The susceptibility of the strain was tested against the list of antibiotics proposed for *Corynebacterium* in the technical Guidance on the characterisation of microorganisms used as feed

⁸ FEED dossier reference: FAD-2018-0035.

⁹ The full report is available on the EURL website: https://ec.europa.eu/jrc/sites/jrcsh/files/finirep_fad-2018-0035_lthreonine.pdf

¹⁰ Commission Regulation (EC) No 429/2008 of 25 April 2008 on detailed rules for the implementation of Regulation (EC) No 1831/2003 of the European Parliament and of the Council as regards the preparation and the presentation of applications and the assessment and the authorisation of feed additives. OJ L 133, 22.5.2008, p. 1.

additives or as production organisms (EFSA FEEDAP Panel, 2018a,b).

C. glutamicum

therefore fulfils all criteria for qualification as QPS.

3.1.2. Manufacturing process

¹⁴

3.1.3. Characterisation of the additive

L-Threonine (International Union of Pure and Applied Chemistry (IUPAC) name: (2*S*,3*R*)-2-amino-3-hydroxybutanoic acid; synonyms: 2-amino-3-hydroxybutyric acid, α -amino- β -hydroxybutyric acid), a compound identified with the Chemical Abstracts Service (CAS) No 72-19-5 and the European Inventory of Existing Commercial Chemical Substances (EINECS) No 200-774-1, has a molecular weight of 119.12 Da. The molecular formula of L-threonine is C₄H₉NO₃. The structural formula is given in Figure 1.




Figure 1: Structural formula of L-threonine

The additive contains by specification $\geq 98.5\%$ L-threonine 'as is' ($\geq 99.0\%$ L-threonine on a dry matter basis), $\leq 0.5\%$ moisture and $\leq 0.1\%$ ash.¹⁶ The analysis of five batches of the additive showed an average of threonine of 99.2% 'as is' (range 99.1–99.2%).¹⁷ The loss on drying was 0.14% (range 0.13–0.16%) and the ash 0.03% (range 0.02–0.03%). Other amino acids (lysine, serine, glutamic acid, isoleucine and valine) represented 0.06% and sulfate was 0.01% (in all batches).¹⁸ The amount of unidentified material was lower than 1% on a dry matter basis.

The specific optical rotation was measured in three batches of the final product and the average was -28.4° (range -28.3 to -28.5°),¹⁹ which is within the range established for L-threonine in the European Pharmacopoeia (-29.0 to -27.6°) and demonstrates the identity of the L-enantiomer.

3.1.3.1. Impurities

Three batches were analysed for heavy metals (cadmium, mercury and lead) and arsenic and in all cases the levels were found under the limit of detection (LOD).²⁰ None of these amounts were considered of concern.

The microbiological quality of three batches of the product was tested by counting total bacterial count ($< 10^3$ colony forming units (CFU)/g) *Salmonella* spp. (negative in 25 g), *E. coli* (negative), coliform bacteria (negative) and filamentous fungi and yeasts ($< 5 \times 10^2$ CFU/g).²¹ Regarding mycotoxins, ochratoxin A; aflatoxins B₁, B₂, G₁, G₂; zearalenone; fumonisin B₁ and B₂; and deoxynivalenol were analysed in three batches of the additive. All values were below the LOD and are of no concern.²² Pesticides (358 species) were analysed in three batches and the values were below the LOD.²³

¹⁴ Technical dossier/Section II.3 and III.B.2.2.1.

¹⁶ Technical dossier/Section II.1.3 and annex II.1.1.

¹⁷ Technical dossier/Section II/Annex II.1.3., analytical method AOAC 999.13.

¹⁸ Technical dossier/Section II/Annex II.1.3.

¹⁹ Technical dossier/Supplementary information November 2018/Annex SIn 1.

²⁰ Technical dossier/Section II/Annex II.1.2. LOD (in $\mu\text{g}/\text{kg}$) was 1 for arsenic, lead and cadmium and 5 for mercury.

²¹ Technical dossier/Section II/Annex II.1.2 and Supplementary information November 2018/02 SIn CJE L-thr.

²² Technical dossier/Section II/Annex II.1.2. LOD (in $\mu\text{g}/\text{kg}$) was 0.1 for ochratoxin A and aflatoxins, 1.5 for zearalenone, 5 for fumonisin and 0.5 for deoxynivalenol.

²³ Technical dossier/Section II/Annex II.1.5. LOD in $\mu\text{g}/\text{kg}$ was < 0.8 .

Dioxins (polychlorinated dibenzofurans (PCDF), polychlorinated dibenzo(*p*)dioxins (PCDD)) and dioxin-like polychlorinated biphenyls (DL-PCBs) were measured in three batches of the final product.²⁴ PCDD/F were < 0.05 µg WHO-TEQ/kg. Dioxin-like PCBs were < 0.16 µg WHO-TEQ/kg.

The absence of viable cells in the final product was tested: [REDACTED]

3.1.3.2. Physical characteristics

The additive is an off-white free flowing powder. It has a pH of 4.5–7 in 10% water solution, a bulk density of 700–900 kg/m³. Its melting point is 256°C and its solubility in water at 20°C is 97.6 g/L.²⁶

The particle size distribution of the final product (three batches) was analysed by sieving.²⁷ The fractions of particles having a diameter < 125, and < 75 µm ranged 36–42, and 10–18% (v/w), respectively. The applicant demonstrated that the particle size distribution was very similar to the L-threonine produced by strain *C. glutamicum* [REDACTED] (three batches analysed by sieving showed fractions of particles having a diameter < 125, and < 75 µm ranged 37–47, and 12–16% (v/w), respectively).

No information on the dusting potential of the product under assessment was provided.

3.1.3.3. Stability and homogeneity

No information on the shelf life, stability (in premixtures and feedingstuffs) and capacity of the additive to distribute homogeneously in feed of the additive under assessment was provided. The applicant provided information on the shelf life, stability in premixtures and feedingstuffs and on the capacity of L-threonine to distribute homogeneously in feed performed with an L-threonine originating from a different strain (*C. glutamicum* [REDACTED]).²⁸ As the production process is the same and the product characteristics are very similar, the FEEDAP Panel considers that the results of those studies can be applicable to the product under assessment.

A previous study on the shelf life of L-threonine produced by *C. glutamicum* [REDACTED] when stored in sealed brown glass at 25°C and 40°C for 6 months showed no losses.

L-Threonine produced by *C. glutamicum* [REDACTED] (three batches) was stable in a premixture containing 4% choline chloride when supplemented at 5% and stored in aluminium vacuum bags at 25°C for 6 months.

L-Threonine produced by *C. glutamicum* [REDACTED] (three batches) was stable in a complete feed for chicken for fattening when supplemented via premixture at 0.4%. The basal diet consisted of maize, soybean meal and wheat. The samples were stored at 25°C in aluminium vacuum bags for 3 months. Losses ranged from 0% to 5%.²⁹

L-Threonine produced by *C. glutamicum* [REDACTED] (three batches) was stable in water for drinking when supplemented at 0.05%. Samples were stored in aluminium vacuum bags at 25 and 40°C for 48 h.

The capacity of the L-threonine produced by *C. glutamicum* [REDACTED] to distribute homogeneously in feed was studied in the premixture described above, in the mash feed described above and in a different pelleted feed. The pelleted feed was supplemented with 0.2% L-threonine and conditioned at 72°C, pelleted at 82°C and dried at 60–65°C. The coefficients of variation were 3%, 2% and 4%, respectively.

3.1.3.4. Physico-chemical incompatibilities

No physico-chemical incompatibilities in feed are expected with other additives, medicinal products or feed materials.

²⁴ Technical dossier/Section II/Annex II.1.2.

²⁶ Technical dossier/Section II/Annex II.3.1.

²⁷ Technical dossier/Section II/Annexes II.1.6 and II.1.7.

²⁸ Technical dossier/Section II/Annexes II.4.1 to II.4.8.

²⁹ Technical dossier/Section II/Annex II.4.3.

3.1.4. Conditions of use

It is proposed that L-threonine will be used in feeds to achieve an adequate amino acid profile and to meet the L-threonine requirements for all animal species. It can be added directly to feedingstuffs or complementary feedingstuffs, or via a premixture. It is also proposed to use the additive in water for drinking. No inclusion levels have been proposed, as the requirements, in quantitative terms, depend on the species, the physiological state of the animal, the performance level, the environmental conditions and the amino acid composition of the unsupplemented diet.³⁰

3.2. Safety

3.2.1. Safety for the target species

Concerns from the use of the additive may arise from residues of the fermentation process/production strain remaining in the final product. The additive is highly purified (> 99%), is produced by fermentation using a strain that belongs to a species and that qualifies for the QPS approach for safety assessment. Therefore, the FEEDAP Panel concludes that L-threonine produced by *C. glutamicum* [REDACTED] is safe for the target species provided that it is supplemented in appropriate amounts to the diets. Due to the risk of nutritional imbalances and hygienic reasons associated to the use of amino acids via water for drinking (EFSA FEEDAP Panel, 2010), the FEEDAP Panel has concerns on the safety of the simultaneous oral administration of threonine-containing additives via feed and water for drinking.

3.2.2. Safety for the consumer

The amino acid L-threonine, supplemented to feed, will be incorporated into proteins of tissues and/or products of animal origin and any of their potential excess will be metabolised and excreted as urea/uric acid and carbon dioxide. Therefore, the composition of tissues and products of animal origin will not be affected by the use of L-threonine in animal nutrition.

The product under assessment is produced by fermentation using a *C. glutamicum* strain which fulfils the qualifications for the QPS approach to safety assessment. Therefore, the FEEDAP Panel concludes that the use of L-threonine produced by *C. glutamicum* [REDACTED] in animal nutrition is safe for the consumer.

3.2.3. Safety for the user

No studies to support the safety of the additive for the user were performed using the additive under assessment as test item. The applicant submitted an acute inhalation toxicity study, an *in vitro* skin irritation test, and eye irritation/corrosion test and a skin sensitisation test performed using L-threonine produced by *C. glutamicum* [REDACTED] as test item (EFSA FEEDAP Panel, 2019).³¹ As the production process and the product characteristics are very similar, the FEEDAP Panel considers that the conclusions of these studies are applicable to the product under assessment.

3.2.3.1. Conclusions on safety for the user

The additive is not a skin or eye irritant and is not a skin sensitiser. Although the workers can be exposed by inhalation, the results of an acute inhalation study showed that risk of adverse effects by inhalation is low.

3.2.4. Safety for the environment

The amino acid L-threonine is a physiological and natural component of the proteins of living organisms. When consumed, it will be absorbed, and the non-absorbed fraction will be incorporated into the intestinal microbial mass and excreted as such. The absorbed L-threonine will be incorporated into body protein or excreted as urea/uric acid and as carbon dioxide.

L-Threonine produced using *C. glutamicum* [REDACTED] is safe for the environment.

³⁰ Technical dossier/Section II.5.1.

³¹ Technical dossier/Section III/Annexes III.3.2 to III.3.5.

3.3. Efficacy

Efficacy studies are not required for amino acids which naturally occur in the proteins of plants and animals. The nutritional role of L-threonine is well established in the scientific literature. Since most of the studies have been performed with supplemental L-threonine, the product L-threonine produced by *C. glutamicum* [REDACTED] is regarded as an effective source of the amino acid L-threonine.

The efficacy of L-threonine for both non-ruminant and ruminant species was described in previous opinions (EFSA FEEDAP Panel, 2013, 2014a). Supplemental L-threonine is degraded by ruminal microbiota if not given in a protected form.

3.4. Post-market monitoring

The FEEDAP Panel considers that there is no need for specific requirements for a post-market monitoring plan other than those established in the Feed Hygiene Regulation³² and Good Manufacturing Practice.

4. Conclusions

L-Threonine produced by *C. glutamicum* [REDACTED] is considered safe for the target species when supplemented in appropriate amounts to the diet. The FEEDAP Panel has concerns on the safety of the simultaneous oral administration of L-threonine via feed and water for drinking.

The use of L-threonine produced by *C. glutamicum* [REDACTED] in animal nutrition is safe for the consumer.

The additive is not a skin or eye irritant and is not a skin sensitiser. Although the workers can be exposed by inhalation, the results of an acute inhalation study showed that risk of adverse effects by inhalation is low.

L-Threonine produced by *C. glutamicum* [REDACTED] is safe for the environment.

The product L-threonine is regarded as an effective source of the amino acid L-threonine for all non-ruminant species. For the supplemental L-threonine to be as efficacious in ruminants as in non-ruminant species, it requires protection against degradation in the rumen.

Documentation provided to EFSA

- 1) Feed grade L-threonine produced by fermentation with *Corynebacterium glutamicum* [REDACTED]. June 2018. Submitted by CJ Europe GmbH.
- 2) Feed grade L-threonine produced by fermentation with *Corynebacterium glutamicum* [REDACTED]. Supplementary information. November 2018. Submitted by CJ Europe GmbH.
- 3) Evaluation report of the European Union Reference Laboratory for Feed Additives on the Methods of Analysis for L-threonine produced by fermentation with *Corynebacterium glutamicum* [REDACTED].
- 4) Comments from Member States.

Chronology

Date	Event
5/6/2018	Dossier received by EFSA
19/6/2018	Reception mandate from the European Commission
31/7/2018	Application validated by EFSA – Start of the scientific assessment
9/10/2010	Request of supplementary information to the applicant in line with Article 8(1)(2) of Regulation (EC) No 1831/2003 – Scientific assessment suspended. <i>Issues: characterisation of the additive and manufacturing process</i>
31/10/2018	Comments received from Member States
15/11/2018	Reception of supplementary information from the applicant - Scientific assessment re-started
30/11/2018	Reception of the Evaluation report of the European Union Reference Laboratory for Feed Additives
22/1/2019	Opinion adopted by the FEEDAP Panel. End of the Scientific assessment

³² Regulation (EC) No 1831/2003 of the European Parliament and of the Council of 22 September 2003 laying down requirements for feed hygiene. OJ L 35, 8.2.2005, p. 1.

References

- EFSA (European Food Safety Authority), 2008. Technical Guidance of the Scientific Panel on Additives and Products or Substances used in Animal Feed (FEEDAP) for assessing the safety of feed additives for the environment. EFSA Journal 2008;6(10):842, 28 pp. <https://doi.org/10.2903/j.efsa.2008.842>
- EFSA BIOHAZ Panel (EFSA Panel on Biological Hazards), Koutsoumanis K, Allende A, Alvarez-Ordóñez A, Bolton D, Bover-Cid S, Chemaly M, Davies R, Hilbert F, Lindqvist R, Nauta M, Peixe L, Ru G, Simmons M, Skandamis P, Suffredini E, Cocconcelli PS, Fernandez Escamez PS, Maradona MP, Querol A, Suarez JE, Sundh I, Vlák J, Barizzone F, Correia S and Herman L, 2019. Statement on the update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 9: suitability of taxonomic units notified to EFSA until September 2019. EFSA Journal 2019;17(1):5555, 46 pp. <https://doi.org/10.2903/j.efsa.2019.5555>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances Used in Animal Feed), 2010. Scientific Opinion on the use of feed additives authorised/applied for use in feed when supplied via water. EFSA Journal 2010;8(12):1956, 9 pp. <https://doi.org/10.2903/j.efsa.2010.1956>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2012a. Guidance for the preparation of dossiers for nutritional additives. EFSA Journal 2012;10(1):2535, 14 pp. <https://doi.org/10.2903/j.efsa.2012.2535>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2012b. Guidance on studies concerning the safety of use of the additive for users/workers. EFSA Journal 2012;10(1):2539, 5 pp. <https://doi.org/10.2903/j.efsa.2012.2539>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2013. Scientific opinion on the safety and efficacy of L-threonine produced by *Escherichia coli* (FERM BP-10942) for all animal species based on a dossier submitted by Ajinomoto Eurolysine SAS. EFSA Journal 2013;11(7):3319, 19 pp. <https://doi.org/10.2903/j.efsa.2013.3319>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2014a. Scientific Opinion on the safety and efficacy of L-threonine (ThreAMINO) produced by *Escherichia coli* K-12 (DSM25086) for all animal species and categories based on a dossier by Evonik Industries A.G. EFSA Journal 2014;12(2):3564, 2 pp. <https://doi.org/10.2903/j.efsa.2014.3564>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2014b. Scientific Opinion on the safety and efficacy of L-threonine produced by *Escherichia coli* for all animal species, based on a dossier submitted by HELM AG on behalf of Global Bio-Chem Technology. EFSA Journal 2014;12(5):3674, 3 pp. <https://doi.org/10.2903/j.efsa.2014.3674>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2014c. Scientific Opinion on the safety and efficacy of L-threonine produced by *Escherichia coli* for all animal species, based on a dossier submitted by HELM AG on behalf of Star Lake Bioscience Co. EFSA Journal 2014;12(6):3726, 14 pp. <https://doi.org/10.2903/j.efsa.2014.3726>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2014d. Scientific Opinion on the safety and efficacy of L-threonine produced by *Escherichia coli* (FERM BP-11383) for all animal species, based on a dossier submitted by Ajinomoto Eurolysine S.A.S. EFSA Journal 2014;12(10):3825, 14 pp. <https://doi.org/10.2903/j.efsa.2014.3825>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2015a. Scientific Opinion on the safety and efficacy of L-threonine produced by *Escherichia coli* for all animal species based on a dossier submitted by HELM AG on behalf of Meihua Holdings Group Co. Ltd. EFSA Journal 2015;13(3):4051, 17 pp. <https://doi.org/10.2903/j.efsa.2015.4051>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2015b. Scientific Opinion on the safety and efficacy of L-threonine produced by *Escherichia coli* strains NRRL B-30843, DSM26131, KCCM11133P or DSM 25085 for all animal species based on a dossier submitted by AMAC EEIG. EFSA Journal 2015;13(9):4236, 34 pp. <https://doi.org/10.2903/j.efsa.2015.4236>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2016a. Scientific opinion on the safety of L-threonine produced by fermentation using *Escherichia coli* CGMCC 3703, for all animal species based on a dossier submitted by GBT Europe GmbH. EFSA Journal 2016;14(1):4344, 13 pp. <https://doi.org/10.2903/j.efsa.2016.4344>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), 2016b. Scientific opinion on the safety of L-threonine, technically pure, produced by fermentation with *Escherichia coli* CGMCC7.58 for all animal species based on a dossier submitted by Feedway Europe NV. EFSA Journal 2016;14(5):4470, 11 pp. <https://doi.org/10.2903/j.efsa.2016.4470>

- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), Rychen G, Aquilina G, Azimonti G, Bampidis V, Bastos ML, Bories G, Chesson A, Cocconcelli PS, Flachowsky G, Gropp J, Kolar B, Kouba M, Lopez-Alonso M, Lopez Puente S, Mantovani A, Mayo B, Ramos F, Saarela M, Villa RE, Wester P, Costa L, Dierick N, Leng L, Glandorf B, Herman L, Kärenlampi S, Aguilera J, Tarrés-Call J and Wallace RJ, 2017a. Scientific Opinion on the safety and efficacy of L-threonine produced by fermentation with *Escherichia coli* CGMCC 11473 for all animal species. EFSA Journal 2017;15(7):4939, 16 pp. <https://doi.org/10.2903/j.efsa.2017.4939>
- EFSA FEEDAP Panel (EFSA Panel on additives and products or substances used in animal feed), Rychen G, Aquilina G, Azimonti G, Bampidis V, Bastos ML, Bories G, Chesson A, Cocconcelli PS, Flachowsky G, Gropp J, Kolar B, Kouba M, López-Alonso M, López Puente S, Mantovani A, Mayo B, Ramos F, Saarela M, Villa RE, Wallace RJ, Wester P, Anguita M, Galobart J and Innocenti ML, 2017b. Guidance on the identity, characterisation and conditions of use of feed additives. EFSA Journal 2017;15(10):5023, 12 pp. <https://doi.org/10.2903/j.efsa.2017.5023>
- EFSA FEEDAP Panel (EFSA Panel on additives and products or substances used in animal feed), Rychen G, Aquilina G, Azimonti G, Bampidis V, Bastos ML, Bories G, Chesson A, Cocconcelli PS, Flachowsky G, Gropp J, Kolar B, Kouba M, López-Alonso M, López Puente S, Mantovani A, Mayo B, Ramos F, Saarela M, Villa RE, Wallace RJ, Wester P, Anguita M, Galobart J, Innocenti ML and Martino L, 2017c. Guidance on the assessment of the safety of feed additives for the target species. EFSA Journal 2017;15(10):5021, 19 pp. <https://doi.org/10.2903/j.efsa.2017.5021>
- EFSA FEEDAP Panel (EFSA Panel on additives and products or Substances used in Animal Feed), Rychen G, Aquilina G, Azimonti G, Bampidis V, Bastos ML, Bories G, Chesson A, Cocconcelli PS, Flachowsky G, Gropp J, Kolar B, Kouba M, López-Alonso M, López Puente S, Mantovani A, Mayo B, Ramos F, Saarela M, Villa RE, Wallace RJ, Wester P, Anguita M, Dujardin B, Galobart J and Innocenti ML, 2017d. Guidance on the assessment of the safety of feed additives for the consumer. EFSA Journal 2017;15(10):5022, 17 pp. <https://doi.org/10.2903/j.efsa.2017.5022>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), Bampidis V, Azimonti G, Bastos ML, Christensen H, Dusemund B, Kouba M, Kos Durjava M, López-Alonso M, López Puente S, Marcon F, Mayo B, Pechová A, Petkova M, Ramos F, Sanz Y, Villa RE, Woutersen R, Cocconcelli PS, Costa L, Dierick N, Flachowsky G, Glandorf B, Herman L, Kärenlampi S, Mantovani A, Saarela M, Anguita M, Tarrés-Call J and Wallace RJ, 2018a. Scientific Opinion on the safety and efficacy of L-threonine produced by fermentation using *Escherichia coli* CGMCC 7.232 for all animal species. EFSA Journal 2018;16(10):5458, 17 pp. <https://doi.org/10.2903/j.efsa.2018.5458>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), Rychen G, Aquilina G, Azimonti G, Bampidis V, Bastos ML, Bories G, Chesson A, Cocconcelli PS, Flachowsky G, Gropp J, Kolar B, Kouba M, López-Alonso M, López Puente S, Mantovani A, Mayo B, Ramos F, Saarela M, Villa RE, Wallace RJ, Wester P, Glandorf B, Herman L, Kärenlampi S, Aguilera J, Anguita M, Brozzi R and Galobart J, 2018b. Guidance on the characterisation of microorganisms used as feed additives or as production organisms. EFSA Journal 2018;16(3):5206, 24 pp. <https://doi.org/10.2903/j.efsa.2018.5206>
- EFSA FEEDAP Panel (EFSA Panel on Additives and Products or Substances used in Animal Feed), Bampidis V, Azimonti G, Bastos ML, Christensen H, Dusemund B, Kouba M, Kos Durjava M, López-Alonso M, López Puente S, Marcon F, Mayo B, Pechová A, Petkova M, Ramos F, Sanz Y, Villa RE, Woutersen R, Costa L, Dierick N, Glandorf B, Flachowsky G, Herman L, Mantovani A, Saarela M, Anguita M, Tarrés-Call J, and Wallace RJ, 2019. Safety and efficacy of L-Threonine produced by fermentation with *Corynebacterium glutamicum* [REDACTED] for all animal species. EFSA Journal 2019;17(2):5603, 16 pp. <https://doi.org/10.2903/j.efsa.2019.5603>
- Ph. Eur. (European Pharmacopoeia), 2016. European Directorate for the Quality of Medicines and Health Care. 9th Edition.
- WHO (World Health Organization), 2012. Evaluation of certain food additives: seventy-sixth report of the Joint FAO/WHO Expert Committee on Food Additives (JECFA). WHO technical report series No 974.

Abbreviations

BIOHAZ	EFSA Panel on Biological Hazards
CAS	Chemical Abstracts Service
CFU	colony forming unit
DL-PCB	dioxin-like polychlorinated biphenyls
EINECS	European Inventory of Existing Commercial chemical Substances
EURL	European Union Reference Laboratory
FCC	Food Chemical Codex
FEEDAP	EFSA Panel on Additives and Products or Substances used in Animal Feed
IEC	ion exchange chromatography
IUPAC	International Union of Pure and Applied Chemistry

JECFA	Joint FAO/WHO Expert Committee on Food Additives
█	█
LOD	limit of detection
PCDD	polychlorinated dibenzodioxin
PCDF	polychlorinated dibenzofuran
PCR	polymerase chain reaction
QPS	qualified presumption of safety
TEQ	Toxic equivalent
VIS/FLD	visible or fluorescence detection
WHO	World Health Organization

Annex A – Executive summary of the evaluation report of the European Union Reference Laboratory for feed additives on the methods of analysis for L-threonine produced by fermentation with *Corynebacterium glutamicum* [REDACTED]

In the current application, authorisation is sought under Article 4(1) for L-threonine produced by fermentation with *Corynebacterium glutamicum* [REDACTED], under the category/functional group 3(c) 'nutritional additives'/amino acids, their salts and analogues', according to Annex I of Regulation (EC) No 1831/2003. Authorisation is sought for all animal species.

According to the Applicant, the product is an off-white powder with a minimum purity of 98.5%.

L-Threonine is intended to be mixed either in premixtures or added directly to feedingstuffs or in addition to water. However, the Applicant did not propose a minimum or maximum L-threonine content in feedingstuffs.

For the quantification of L-threonine in the feed additive, premixtures and feedingstuffs, the Applicant submitted the ring-trial validated Community method (Commission Regulation (EC) No 152/2009) based on ion exchange chromatography coupled with photometric detection (IEC-VIS). This method, designed only for the analysis of premixtures and feedingstuffs, does not distinguish between the salts and the amino acid enantiomers. The following performance characteristics were reported for the quantification of total threonine: a relative standard deviation for repeatability (RSDr) ranging from 1.9% to 2.7%, and a relative standard deviation for reproducibility (RSDR) ranging from 3.8% to 5.2%.

For the quantification of L-threonine in the feed additive, the EURL identified the ring-trial validated method EN ISO 17180:2013 based on IEC coupled with post-column derivatisation and visible or fluorescence detection (IEC-VIS/FLD). The following performance characteristics are reported: a RSDr ranging from 0.7% to 1.4%; and a RSDR ranging from 1.9% to 2.3%. In addition, the EURL identified the 'L-threonine monograph' of the Food Chemical Codex (FCC) for the identification of L-threonine in the feed additive.

Within the dossier, the Applicant presented experimental data obtained analysing threonine in water with the AOAC official method 999.13 based on IEC-VIS/FLD. The results presented are considered sufficient to demonstrate the suitability of the procedure for the analysis of the amino acid in water. Hence, the EURL recommends for official control this method to quantify threonine in water.

In the frame of this authorisation the EURL recommends for official control (i) the 'L-threonine monograph' of the FCC based on infrared absorption for the identification of L-threonine in the feed additive; (ii) the ring-trial validated method EN ISO 17180:2013 based on IEC-VIS/FLD to quantify free threonine in the feed additive and premixtures (containing more than 10% threonine); (iii) the Community method based on IEC-VIS for the quantification of threonine in premixtures and feedingstuffs; and (iv) the analytical method described by AOAC (999.13) based on IEC-VIS/FLD to quantify threonine in water.

Further testing or validation of the methods to be performed through the consortium of National Reference Laboratories as specified by Article 10 (Commission Regulation (EC) No 378/2005), as last amended by Regulation (EU) 2015/1761) is not considered necessary.