












Rapid Communication  
Virology



# Molecular characterization of a pigeon paramyxovirus type 1 virus isolated from Eurasian collared doves in Iran, 2017

Reza Esmaelzadeh-Dizaji <sup>1</sup>, Aidin Molouki <sup>2</sup>, Hossein Hosseini <sup>3</sup>,  
Mohammad Hossein Fallah-Mehrabadi <sup>2</sup>, Zahra Ziafati-Kafi <sup>4</sup>, Azin Takalou <sup>5</sup>,  
Nava Eram <sup>1</sup>, Niloufar Kumar <sup>5</sup>, Alireza Ashuri <sup>1</sup>, Naser Sadri <sup>4</sup>,  
Arash Ghalyanchi-Langeroudi <sup>4,\*</sup>

<sup>1</sup>Department of Poultry Diseases, Faculty of Veterinary Medicine, University of Tehran, Tehran 1419963111, Iran

<sup>2</sup>Department of Poultry Diseases, RAZI Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Karaj 3197619751, Iran

<sup>3</sup>Department of Clinical Sciences, Faculty of Veterinary Medicine, Islamic Azad University, Karaj Branch, Karaj 3149968111, Iran

<sup>4</sup>Department of Microbiology and Immunology, Faculty of Veterinary Medicine, University of Tehran, Tehran 1419963111, Iran

<sup>5</sup>Department of Microbiology, Faculty of Veterinary Sciences, Science and Research Branch, Islamic Azad University, Tehran 1477893855, Iran

 OPEN ACCESS

Received: Sep 23, 2021

Revised: Dec 19, 2021

Accepted: Dec 25, 2021

Published online: Jan 27, 2022

\*Corresponding author:

Arash Ghalyanchi-Langeroudi

Department of Microbiology and Immunology,  
Faculty of Veterinary Medicine, University of  
Tehran, Tehran 1419963111, Iran.

Email: arashghalyanchi@gmail.com

ghalyana@ut.ac.ir

https://orcid.org/0000-0001-7914-0863

## ABSTRACT

In September 2017, an outbreak with high mortality, which showed the typical signs of ND, occurred among a flock of more than 2000 Eurasian collared doves in Konarak, southeast of Iran. A confirmed pigeon paramyxovirus type 1 strain was isolated from the brain tissues of the dead doves. The isolate, which was called Pigeon/Iran/Konarak/Barin/2017, was classified as a highly velogenic NDV. Complete genome sequencing and phylogenetic analysis showed that the isolate belonged to subgenotype XXI.2, which has never been reported from Iran before. The isolate had the highest homology (96.15%) with early 2010s Italian isolates. Further studies will be required to understand the diversity better.

**Keywords:** Pigeon paramyxovirus serotype-1 (PPMV-1); whole genome; fusion gene; phylogenetic study; subgenotype XXI.2; Iran

## INTRODUCTION

NDV has a single-stranded negative-sense and non-segmented RNA genome [1]. The genome contains six genes of nucleocapsid (NP), phosphoprotein (P), fusion protein (F), matrix protein (M), hemagglutinin-neuraminidase (HN), and RNA large polymerase protein (L) that encode structural proteins with the same names. Although the HN and F glycoproteins are involved in the attachment and fusion of the virus to the host cell, the internal proteins are involved in several functions, such as transcription and replication of the virus [2,3].

© 2022 The Korean Society of Veterinary Science  
This is an Open Access article distributed under the  
terms of the Creative Commons Attribution Non-  
Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-  
commercial use, distribution, and reproduction in any  
medium, provided the original work is properly cited.

**ORCID iDs**

Reza Esmaeelzadeh-Dizaji  
<https://orcid.org/0000-0001-9044-1659>  
 Aidin Molouki  
<https://orcid.org/0000-0002-3430-0955>  
 Hossein Hosseini  
<https://orcid.org/0000-0003-2696-621X>  
 Mohammad Hossein Fallah-Mehrabadi  
<https://orcid.org/0000-0002-1221-7771>  
 Zahra Ziafati-Kafi  
<https://orcid.org/0000-0002-8824-1276>  
 Azin Takalou  
<https://orcid.org/0000-0003-4328-6785>  
 Nava Eram  
<https://orcid.org/0000-0002-3276-1707>  
 Niloufar Kumar  
<https://orcid.org/0000-0002-6810-0054>  
 Alireza Ashuri  
<https://orcid.org/0000-0002-8982-3067>  
 Naser Sadri  
<https://orcid.org/0000-0002-5200-2820>  
 Arash Ghalyanchi-Langeroudi  
<https://orcid.org/0000-0001-7914-0863>

**Author Contributions**

Conceptualization: Ghalyanchi-Langeroudi A, Hosseini H; Data curation: Ziafati-Kafi Z, Esmaeelzadeh-Dizaji R, Ashuri A; Formal analysis: Ziafati-Kafi Z, Molouki A, Fallah-Mehrabadi MH; Funding acquisition: Takalou A, Eram N, Kumar N; Investigation: Takalou A, Esmaeelzadeh-Dizaji R, Molouki A; Methodology: Ghalyanchi-Langeroudi A, Molouki A; Project administration: Ghalyanchi-Langeroudi A; Resources: Kumar N, Eram N; Software: Ziafati-Kafi Z, Molouki A; Supervision: Ghalyanchi-Langeroudi A, Hosseini H, Molouki A; Validation: Ghalyanchi-Langeroudi A; Visualization: Hosseini H; Writing - original draft: Esmaeelzadeh-Dizaji R, Ziafati-Kafi Z; Writing - review & editing: Molouki A, Ghalyanchi-Langeroudi A.

**Conflict of Interest**

The authors declare no conflict of interest.

NDV is divided into two classes (Class I and II) with one and 21 genotypes, respectively. According to the latest classification system, two new genotypes of XX, XXI, and some isolates of genotype VI, have been assigned to the NDV variant frequently reported from Columbiformes [4].

Although infections of pigeons by pigeon paramyxovirus type 1 (PPMV-1) have been reported since the 1930s, major events, such as the 3<sup>rd</sup> ND panzootic, are believed to have started during the late 1970s and have continued until now [5-8]. Indeed, potential transmission from Columbiformes to poultry has always been a concern in the poultry industry.

Doves are domesticated birds that come in close contact with backyard and commercial poultry [9]. They can also serve as reservoirs for different pathogens, such as ND [10]. In this study, the causative agent of an outbreak, i.e., PPMV-1, was characterized and analyzed phylogenetically in a flock of Eurasian collared doves. The PPMV-1 isolate belonged to subgenotype XXI.2, which has never been reported in the authors' geographical region.

## MATERIALS AND METHODS

### Case history

In September 2017, an outbreak with the typical symptoms of ND in a flock of more than 2000 wild Eurasian collared doves was reported in Konarak, Sistan, and Baluchestan southeast Iran (coordinates: 25°21'37"N 60°23'58"E). Brain tissue samples of dead doves were aseptically collected and stored at -70°C, which were used for virus isolation and identification according to the OIE protocols (OIE, 2018).

### Pathogenicity study

The MDT test was conducted according to the OIE guidelines [11,12]. In addition, multiple candlings were used to determine the embryo time of death.

### RNA extraction and RT-PCR

An RNX™-Plus Kit (CinnaGen, Tehran, Iran) was used for RNA extraction from the HI-confirmed allantoic fluids according to the manufacturer's instructions and stored at -70°C until needed. The cDNA was synthesized by applying of Revert Aid First Strand cDNA Synthesis Kit according to the manufacturer's instructions (Fermentas-Thermo Fisher Scientific, Canada). A partial fusion gene was amplified to confirm the presence of the NDV using primers and protocols described previously [13].

### Genome sequencing

The primary results were confirmed by the generation of cDNA libraries for next-generation sequencing (NGS) on a HiSeq instrument (Illumina, USA) by BGI (China). The raw sequence data were analyzed and assembled within a customized workflow on the Galaxy platform, as described previously [14]. The sequences were assembled, edited, and analyzed using CLC sequence viewer 8 (CLC bio Anhus, Denmark).

### Phylogenetic analyses

A phylogenetic tree for the F gene sequence was made using the standard pilot dataset prepared by the NDV consortium [4] (as of October 30, 2020). For HN and whole-genome comparisons, the sequences were selected based on the highest BLAST homologies and the originally

classified isolates in the NDV consortium datasets. The Maximum Likelihood method and a GTR+G model with a bootstrap of either 500 replicates were carried out using MEGA X [23].

## RESULTS

An MDT of 52 h was obtained when the pathogenicity of the HI-positive isolate was examined. Preliminary PCR also showed that the isolate was the NDV. Amino acid (aa) sequence of the F protein cleavage site further showed that the virus contained a <sup>112</sup>RRQKRF<sup>117</sup> motif, an indicator of a virulent strain. The isolate was called Avian\_avulavirus\_1\_isolate\_pigeon/Iran/Konarak/Barin/2017 (hereafter called Barin/2017), and its complete genome was deposited to GenBank under accession number MH044693.1. The size of the genome of the isolate was 15,192 bp.

According to BLAST analysis using the F gene sequence, Barin/2017 shared at most 96.15% identity with the PPMV-1 strains isolated from Israel and Italy in the early 2010s (namely MH377298.1 and N638234.1, respectively). Furthermore, Barin/2017 clustered to subgenotype XXI.2 (formerly subgenotype VIi under Diel et al. 2012 classification based on phylogenetic analysis of the F gene using the standard dataset of the NDV consortium [15]) (**Supplementary Data 1** and **Fig. 1**). Moreover, the genetic distance of the isolate was ~4% (below the 5% threshold) with other XXI.2 isolates, as shown in **Supplementary Data 2**.


BLAST and phylogenetic analyses based on the HN and the whole genome (15,192 b) were also conducted; similar results showing the same XXI.2 isolates were obtained.

## DISCUSSION

NDV genotypes, such as VI, XX, and XXI, are sometimes referred to as PPMV-1 (shown in **Fig. 1** as a subtree derived from **Supplementary Data 1**), which are considered the most genetically diverse groups among the NDV strains [2]. Over the years, they have adapted to Columbiformes and became generally affiliated with them. Moreover, sporadic outbreaks have been reported in poultry farms [16], making them an important issue that must be monitored and studied extensively.

In this study, the Barin/2017 isolate was categorized into subgenotype XXI.2, and is the first report of a subgenotype XXI.2 strain of an Iranian PPMV-1. Only two other complete genomes sequenced strains, belonging to subgenotype XXI.2, are currently available in GenBank, with the accession numbers of MH377298.1 and MG456676.1; the latter is an Iranian PPMV-1 and has some diversity in nucleotide sequences compared to the present isolate. Overall, Barin/2017 may belong to the subgenotype XXI.2 that has mutated since the early 2010s and was found in Iran or other parts of the world.

Only a few published reports on subgenotype XXI.2 are available. For example, in 2012, Bonfante et al. [17] reported three PPMV-1 strains classified into the subgenotype XXI.2 cluster (formerly VIi sub-genotype and sublineage 4a). Similarly, such viruses were isolated from collared doves during the surveillance carried out on wild birds in Italy in 2010–2011. In a different study, Snoeck et al. [18] reported two pigeon-derived subgenotype XXI.2 strains isolated from live markets in Nigeria. The analyses showed that Barin/2017 shared the same cleavage site motif of <sup>112</sup>RRQKRF<sup>117</sup> with these subgenotype XXI.2 PPMV-1 isolates.



**Fig. 1.** Phylogenetic tree showing that Barin/2017 belonged to NDV subgenotype XXI.2 (marked with ●). The pigeon paramyxovirus type 1 subtree was derived from **Supplementary Data 1**, which was made using the F gene sequence of Barin/2017 and the standard pilot dataset of the NDV consortium [14].

Interestingly, the PPMV-1 subgenotype XXI.1.1 strains (previously called VIg) reported by other Iranian groups [19,20] did not show up more than 90% identity (the threshold set for the genotype and subgenotype proposed by the NDV consortium [4]) to Barin/2017, showing that other PPMV-1 genotypes/subgenotypes may also exist in Iran. Further study will be needed to understand the diversity of Iranian NDV originated from Columbiformes.

The subgenotype XXI.2 may be present and still circulating in other parts of the world, despite Barin/2017 being the last subgenotype XXI.2 reported.

This study provided the full genome sequence and characteristics of a subgenotype XXI.2 PPMV-1 strain, aiming to improve global NDV data. ND caused by PPMV-1 remains an ongoing panzootic since the 1980s, and it has become endemic in domestic and wild free-flying Columbiformes in most parts of the world. Therefore, more studies on this group of viruses will be needed because the circulation of PPMV-1 is a persistent threat to poultry farming in Iran and other countries.

## ACKNOWLEDGEMENTS

The authors wish to thank many colleagues for stimulating discussions and suggestions. The University of Tehran supported this study.

## SUPPLEMENTARY MATERIALS

### Supplementary Data 1

Evolutionary analysis inferred using the Maximum Likelihood method and General Time Reversible model [21] using the standard pilot dataset of the NDV consortium [4] and the F gene sequence of Barin/2017. The isolate clustered to the subgenotype XII.2 strains. The bootstrap consensus tree inferred from 500 replicates [22] was taken to represent the evolutionary history of the taxa analyzed. A discrete Gamma distribution was used to model the evolutionary rate differences among the sites (four categories [+G, parameter = 0.4095]). All positions containing gaps and missing data were eliminated (complete deletion option). A total of 1662 positions were in the final dataset. Evolutionary analyses were conducted in MEGA X [23]. Barin/2017 was marked with ●.

[Click here to view](#)

### Supplementary Data 2

The evolutionary distance based on the sequence of the F gene. Barin/2017 showed ~4% distance (below the 5% threshold) to the subgenotype XXI.2 strains in the standard dataset of the NDV consortium, suggesting it belonged to the same cluster. The numbers of base substitutions per site from between the sequences are shown. Standard error estimate(s) are shown above the diagonal. The analyses were conducted using the Maximum Composite Likelihood model [24]. All positions containing gaps and missing data were eliminated (complete deletion option). There were 1662 positions in the final dataset. Evolutionary analyses were conducted in MEGA X [23].

[Click here to view](#)

## REFERENCES

1. Yusoff K, Tan WS. Newcastle disease virus: macromolecules and opportunities. *Avian Pathol.* 2001;30(5):439-455.  
[PUBMED](#) | [CROSSREF](#)
2. Dimitrov KM, Ramey AM, Qiu X, Bahl J, Afonso CL. Temporal, geographic, and host distribution of avian paramyxovirus 1 (Newcastle disease virus). *Infect Genet Evol.* 2016;39:22-34.  
[PUBMED](#) | [CROSSREF](#)
3. Gogoi P, Ganar K, Kumar S. Avian paramyxovirus: a brief review. *Transbound Emerg Dis.* 2017;64(1):53-67.  
[PUBMED](#) | [CROSSREF](#)
4. Dimitrov KM, Abolnik C, Afonso CL, Albina E, Bahl J, Berg M, et al. Updated unified phylogenetic classification system and revised nomenclature for Newcastle disease virus. *Infect Genet Evol.* 2019;74:103917.  
[PUBMED](#) | [CROSSREF](#)
5. Alexander DJ. Newcastle disease and other avian paramyxoviruses. *Rev Sci Tech.* 2000;19(2):443-462.  
[PUBMED](#) | [CROSSREF](#)

6. Doyle T. The virus diseases of animals with special reference to those of poultry. *J Comp Pathol Ther.* 1933;46:90-107.  
[CROSSREF](#)
7. Suarez DL, Miller PJ, Koch G, Mundt E, Rautenschlein S. Newcastle disease, other avian paramyxoviruses, and avian metapneumovirus infections. In: Swayne DE, editor. *Diseases of Poultry.* 2020, 109-166.
8. Yang CY, Shieh HK, Lin YL, Chang PC. Newcastle disease virus isolated from recent outbreaks in Taiwan phylogenetically related to viruses (genotype VII) from recent outbreaks in western Europe. *Avian Dis.* 1999;43(1):125-130.  
[PUBMED](#) | [CROSSREF](#)
9. Ali S, Ripley SD. *Handbook of the Birds of India and Pakistan Together With Those of Bangladesh, Nepal, Bhutan and Sri Lanka: Stone Curlews to Owls, vol. 3.* Oxford: Oxford University Press; 1981.
10. Terregino C, Cattoli G, Grossele B, Bertoli E, Tisato E, Capua I. Characterization of Newcastle disease virus isolates obtained from Eurasian collared doves (*Streptopelia decaocto*) in Italy. *Avian Pathol.* 2003;32(1):63-68.  
[PUBMED](#) | [CROSSREF](#)
11. Wei T, Deng Q, Li H, Pan C, Zhai G, Yuan Y, et al. Molecular characterization of two novel sub-sublineages of pigeon paramyxovirus type 1 in China. *Arch Virol.* 2018;163(11):2971-2984.  
[PUBMED](#) | [CROSSREF](#)
12. OIE. *Newcastle Disease, Biological Standards Commission, Manual of Diagnostic Tests and Vaccines for Terrestrial Animals: Mammals, Birds and Bees.* 7th ed. Paris: World Organisation for Animal Health; 2012, 555-574.
13. Kant A, Koch G, Van Roozelaar DJ, Balk F, Huurne AT. Differentiation of virulent and non-virulent strains of Newcastle disease virus within 24 hours by polymerase chain reaction. *Avian Pathol.* 1997;26(4):837-849.  
[PUBMED](#) | [CROSSREF](#)
14. Dimitrov KM, Sharma P, Volkening JD, Goraichuk IV, Wajid A, Rehmani SF, et al. A robust and cost-effective approach to sequence and analyze complete genomes of small RNA viruses. *Virology.* 2017;14(1):72.  
[PUBMED](#) | [CROSSREF](#)
15. Diel DG, da Silva LH, Liu H, Wang Z, Miller PJ, Afonso CL. Genetic diversity of avian paramyxovirus type 1: proposal for a unified nomenclature and classification system of Newcastle disease virus genotypes. *Infect Genet Evol.* 2012;12(8):1770-1779.  
[PUBMED](#) | [CROSSREF](#)
16. Alexander DJ, Banks J, Collins MS, Manvell RJ, Frost KM, Speidel EC, et al. Antigenic and genetic characterisation of Newcastle disease viruses isolated from outbreaks in domestic fowl and turkeys in Great Britain during 1997. *Vet Rec.* 1999;145(15):417-421.  
[PUBMED](#) | [CROSSREF](#)
17. Bonfante F, Terregino C, Heidari A, Monne I, Salviato A, Taddei R, et al. Identification of APMV-1 associated with high mortality of collared doves (*Streptopelia decaocto*) in Italy. *Vet Rec.* 2012;171(13):327.  
[PUBMED](#) | [CROSSREF](#)
18. Snoeck CJ, Adeyanju AT, Owoade AA, Couacy-Hymann E, Alkali BR, Ottosson U, et al. Genetic diversity of Newcastle disease virus in wild birds and pigeons in West Africa. *Appl Environ Microbiol.* 2013;79(24):7867-7874.  
[PUBMED](#) | [CROSSREF](#)
19. Mayahi M, Shapouri M, Jafari R, Farsani M. Characterization of isolated pigeon paramyxovirus-1 (PMV-1) and its pathogenicity in broiler chickens. *Vet Res Forum.* 2017;8(1):15.
20. Rezaei Far A, Peighambari SM, Pourbakhsh SA, Ashtari A, Soltani M. Co-circulation of genetically distinct groups of avian paramyxovirus type 1 in pigeon Newcastle disease in Iran. *Avian Pathol.* 2017;46(1):36-43.  
[PUBMED](#) | [CROSSREF](#)
21. Nei M, Kumar S. *Molecular Evolution and Phylogenetics.* Oxford: Oxford University Press; 2000.
22. Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution.* 1985;39(4):783-791.  
[PUBMED](#) | [CROSSREF](#)
23. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol.* 2018;35(6):1547-1549.  
[PUBMED](#) | [CROSSREF](#)
24. Tamura K, Nei M, Kumar S. Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc Natl Acad Sci U S A.* 2004;101(30):11030-11035.  
[PUBMED](#) | [CROSSREF](#)