

Original Article



Comparison between adjuvant chemotherapy and adjuvant radiotherapy/chemoradiotherapy after radical surgery in patients with cervical cancer: a meta-analysis

Kwang-Beom Lee ^{1,*}, Seung-Hyuk Shim ^{2,*}, Jong-Min Lee ³

¹Department of Obstetrics and Gynecology, Gachon University Gil Medical Center, Gachon University College of Medicine, Incheon, Korea

²Department of Obstetrics and Gynecology, Konkuk University School of Medicine, Seoul, Korea

³Department of Obstetrics and Gynecology, Kyung Hee University Hospital at Gangdong, Kyung Hee University School of Medicine, Seoul, Korea



Received: Jan 31, 2018

Revised: Mar 12, 2018

Accepted: Apr 3, 2018

Correspondence to

Jong-Min Lee

Department of Obstetrics and Gynecology,
Kyung Hee University Hospital at Gangdong,
Kyung Hee University School of Medicine,
892 Dongnam-ro, Gangdong-gu, Seoul 05278,
Korea.

E-mail: kgo02@hanmail.com

*Kwang-Beom Lee and Seung-Hyuk Shim contributed equally to this work and should be considered as co-first authors.

Copyright © 2018. Asian Society of Gynecologic Oncology, Korean Society of Gynecologic Oncology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Kwang-Beom Lee

<https://orcid.org/0000-0003-4634-176X>

Seung-Hyuk Shim

<https://orcid.org/0000-0001-8043-2257>

Jong-Min Lee

<https://orcid.org/0000-0002-0562-5443>

ABSTRACT

Objective: To estimate the effect of adjuvant chemotherapy (AC) on the prognosis in cervical cancer patients with intermediate- or high-risk factors after radical hysterectomy (RH) compared to that for adjuvant radiotherapy (AR).

Methods: The Embase and MEDLINE databases and the Cochrane Library were searched for published studies comparing cervical cancer patients who received AC with those who received AR after RH. The endpoints were patient oncologic outcome. Random-effects meta-analytical models were used to calculate the pooled estimates of the effect of AC on mortality/recurrence.

Results: Two randomized trials and eleven observational studies (AC, 942 patients; AR, 1,721 patients) met our search criteria. There were no significant differences in mortality and any recurrence between two groups. The results for distant recurrence favored the AC group (pooled odds ratio: 0.69; 95% confidence interval: 0.54–0.88; $p=0.03$). In subgroup analyses (for study design, histology, indication for adjuvant treatment, AR type, AC type, and lymph node metastasis), there was no significant increase in mortality and recurrence for AC compared with that for AR.

Conclusion: Compared to AR, AC showed similar survival outcomes in cervical cancer patients undergoing RH and also appeared to reduce the risk of distant recurrence.

Keywords: Uterine Cervical Neoplasms; Hysterectomy, Drug Therapy; Radiotherapy; Chemoradiotherapy; Meta-Analysis

INTRODUCTION

The International Agency of Research on Cancer estimated that cervical cancer was the fourth most common cancer in women and the seventh overall worldwide in 2012, with an estimated 528,000 new cases. There were 266,000 cervical cancer-related deaths estimated worldwide in 2012, accounting for 7.5% of all female cancer-related deaths [1]. In Korea, there were an

Funding

This work was supported by Konkuk University Medical Center Research Grant 2017.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Author Contributions

Conceptualization: L.K.B., S.S.H., L.J.M.;
Data curation: L.K.B., S.S.H.; Formal analysis:
L.K.B., S.S.H., L.J.M.; Funding acquisition:
L.K.B., S.S.H., L.J.M.; Investigation: L.K.B.,
S.S.H., L.J.M.; Methodology: L.K.B., S.S.H.;
Project administration: L.K.B., S.S.H., L.J.M.;
Resources: L.K.B., S.S.H., L.J.M.; Software:
S.S.H.; Supervision: L.K.B., S.S.H., L.J.M.;
Validation: L.K.B., S.S.H., L.J.M.; Visualization:
L.K.B., S.S.H., L.J.M.; Writing - original draft:
L.K.B., S.S.H., L.J.M.; Writing - review &
editing: L.K.B., S.S.H., L.J.M.

estimated approximately 3,500 new cases and the age-standardized incidence rate was 9.5 per 100,000 persons in 2013 [2]. Most gynecologic oncology surgeons prefer radical surgery in patients with early-stage disease [3]. The Gynecologic Oncology Group (GOG) randomized trial of postoperative radiation therapy (RT) included 277 patients with negative nodes and local high-risk features meeting the Sedlis criteria. Overall, there was a 47% reduction in the risk recurrence. In addition, the SWOG demonstrated that postoperative treatment with chemoradiation resulted in significantly improved pelvic disease control and survival when compared to RT alone in patients who met the Peters criteria [4,5]. So, adjuvant RT has been administered to improve oncologic outcomes in various settings such as RT alone, concurrent chemoradiation therapy, sequential chemoradiation therapy, RT followed by additional chemotherapy, sandwich therapy with chemotherapy and RT. However, some clinicians have performed adjuvant chemotherapy (AC) in patients with certain risk factor(s) for disease recurrence and suggested that AC could be a reasonable option. There are considerable advantages to surgery without adjuvant radiotherapy (AR), including the preservation of vaginal pliability and ovarian function and the opportunity to avoid potential late radiation-related complications.

The present meta-analysis aimed to quantify and compare the effects of AC with any treatment strategies including AR on oncologic outcome in cervical cancer patients with intermediate- or high-risk factors after surgery.

MATERIALS AND METHODS

1. Literature search

A systematic review and meta-analysis were performed using previously described reporting guidelines [6-8]. The Embase and MEDLINE databases and the Cochrane Central Register for Controlled Trials database were searched up to October 2017, irrespective of language. Pre-publication papers were also included. The search strategy is described in Appendix 1. Titles and abstracts were checked to identify potentially eligible studies. The full texts were then reviewed in detail. References were manually screened to find additional studies. Two authors (SHS and KBL) independently performed all searches.

2. Eligibility criteria

The inclusion criteria for this meta-analysis were as follows: 1) a randomized controlled trial (RCT), prospective/retrospective cohort study, nested case-control study; 2) or population-based case-control study of participants receiving primary radical hysterectomy (RH) with lymphadenectomy for cervical cancer (stage I-II); 3) AC following RH, as intervention; 4) AR including radiotherapy or concurrent chemoradiotherapy (CCRT) following RH as a comparison; and 5) outcomes of mortality or recurrence rates measured via relative risks (RRs), odds ratios (ORs), or hazard ratios (HRs) with 95% confidence intervals (CIs) or sufficient data for calculation. For studies with duplicated data, the most recent or instructive study was selected. Single-arm cohort studies and case reports were excluded.

3. Data extraction

The following data were extracted from each study: first author; publication year; study design, location, and period; patient ages; sample size; tumor stage; histology; lymph node metastasis (LNM), parametrial involvement (PMI), resection margin involvement (RMI),

tumor size, deep stromal invasion (DSI), lymphovascular space invasion (LSVI), AC details (regimen, dose, cycle); AR details (dose, use of CCRT, concomitant chemotherapy regimen), follow-up duration; recurrence; death from disease; and indications for adjuvant treatment. Pathological risk factors including LNM, PMI, and RMI were considered high-risk factors [4]; those including bulky tumor (>4 cm diameter), DSI, and LSVI were considered intermediate risk factors [5]. Each study was systematically reviewed for features that could introduce bias, similarity of risk factors for prognosis, and similarity of follow-up durations between the AC and AR groups. Three authors (KBL, SHS, and JML) independently extracted data and discrepancies were jointly reviewed until consensus was reached.

4. Quality assessment

For non-randomized studies (NRSs), the quality of each study was evaluated using the nine-star Newcastle-Ottawa scale (NOS) in three categories: selection, comparability, and exposure (case-control studies) or outcomes (cohort studies) [9]. Based on quality assessment standards from previous meta-analyses [10], the present meta-analysis defined a study with five or more stars as high-quality. To evaluate the study quality for RCTs, the following features were assessed: randomization procedure, estimation of sample size, blinding and allocation concealment, loss to follow-up, dropout, and intention-to-treat analysis [11]. Study quality was quantified using the Jadad/Oxford quality scoring system [12]. Three authors (KBL, SHS, and JML) independently evaluated study quality and discrepancies were jointly reviewed until consensus was reached.

5. Data generation and analysis

The primary endpoint was the mortality rate. The secondary endpoints were the total, local, and distant recurrence rates. The ORs and 95% CIs for the mortality or recurrence rates for the AC and AR groups were calculated from the original data of each study. Cross-study heterogeneity was examined using the Cochran Q test and the I^2 statistic [13,14]. Q test $p < 0.1$ [13] or I^2 statistic $> 50\%$ [14] indicated substantial heterogeneity between studies. A random-effects model was used (DerSimonian-Laird method) to estimate the combined OR [15]. Subgroup analyses according to study design (RCT or NRS), histology (squamous cell carcinoma [SCC] or adenocarcinoma [ADC] or SCC plus ADC), type of AR (RT only or CCRT only or RT plus CCRT), indications for adjuvant treatment (intermediate-risk only or high-risk only), status of LNM (yes or no), and AC regimen (platinum combination with taxane or platinum combination with non-taxane) were performed.

Sensitivity analysis was conducted by withdrawing one study at a time from the meta-analysis to evaluate its effect on the pooled OR [8]. Publication bias was determined using the Begg-Mazumdar rank correlation [16,17] and fail-safe N [18] tests. All analyses were performed using R version 3.0.2 (R Foundation, Vienna, Austria; <http://cran.r-project.org/mirrors.html>). A two-sided p value < 0.05 was considered significant.

RESULTS

1. Literature search

From 2,057 records, 49 papers were identified for detailed full-text review. Finally, 13 studies were included in the meta-analysis [19-31]. **Fig. 1** depicts the literature search process. Supplementary **Table 1** shows the excluded studies and the reasons for exclusion.

Comparison of adjuvant treatment after surgery




Fig. 1. Flow diagram of the literature search process.

Table 1. Characteristics of studies included in the meta-analysis

Study/design	Stage (No.)	Histology, No. (%)	AR	AC regimen
Seki et al. [19]/NRS	IB–IIB (22)	ADC/ADSCC 22 (100%)	CCRT 86 (76%, weekly CDDP), RT only 27 (24%)	TP/NTP
	IB–IIB (113)	SCC 90, ADC/ADSCC 23 (20%)		
Matsuo et al. [20]/NRS	IB1–IIB (319)	SCC 156 (49%), ADC/ADSCC 149 (47%)	CCRT 502 (66%, weekly CDDP), RT only 253 (36%)	TP/NTP
	IB1–IIB (753)	SCC 597 (79%), ADC/ADSCC 154 (20%)		
Takekuma et al. [21]/NRS	IB1–IIB (37)	SCC 24 (65%), ADC/ADSCC 13 (35%)	CCRT 74 (100%, triweekly FP)	TP/NTP
	IB1–IIB (74)	SCC 48 (65%), ADC/ADSCC 26 (35%)		
Li et al. [22]/NRS	IB1–IIA1 (65)	SCC 59 (91%), ADC 6 (9%)	CCRT 68 (100%, weekly CDDP)	TP
	IB1–IIA1 (68)	SCC 63 (93%), ADC 23 (7%)		
Jung et al. [23]/NRS	IB1–IIA (85)	SCC 61 (72%), ADC 21 (25%), ADSCC 3 (4%)	CCRT 177 (100%, weekly CDDP)	TP/NTP
	IB1–IIA (177)	SCC 138 (78%), ADC 26 (15%), ADSCC 13 (7%)		
Shimada et al. [24]/NRS	IB–IIB (64)	ADC/ADSCC 64 (100%)	CCRT (weekly CDDP) or RT only	TP/NTP
	IB–IIB (69)	ADC/ADSCC 69 (100%)		
Hosaka et al. [25]/NRS	IB1–IIB (32)	SCC 24 (75%), ADC/ADSCC 8 (25%)	RT only 49 (100%)	TP
	IB1–IIB (49)	SCC 47 (96%), ADC/ADSCC 2 (4%)		
Mossa et al. [26]/NRS	IB1–IIA (127)	SCC 127 (100%)	RT only 136 (100%)	NTP
	IB1–IIA (136)	SCC 136 (100%)		
Lee et al. [27]/NRS	IB1–IIA (38)	SCC 31 (82%), ADC 2 (5%), ADSCC 5 (13%)	RT only 42 (100%)	TP/NTP
	IB1–IIA (42)	SCC 33 (79%), ADC 3 (7%), ADSCC 5 (12%)		
Hosaka et al. [28]/NRS	IB1–IIB (28)	SCC 28 (100%)	RT only 42 (100%)	NTP
	IB1–IIB (42)	SCC 42 (100%)		
Lahousen et al. [29]/RCT	IB–IIB (52)	SCC 28 (100%)	RT only 24 (100%)	NTP
		SCC 24 (100%)		
Iwasaka et al. [30]/NRS	IB–IIB (53)	SCC 43 (81%), ADSCC 3 (6%), others 7 (13%)	RT only 127 (100%)	NTP
	IB–IIB (127)	SCC 107 (84%), ADSCC 7 (6%), others 13 (10%)		
Curtin et al. [31]/RCT	IB–IIA (88)	SCC 52 (58%), ADC 30 (39%)	RT only 45 (100%)	NTP
RCTs combined	IB–IIB (134)	SCC 104 (74%), ADC 30 (21%)	CCRT 0	
NRSs combined	IB–IIB (805)	SCC 553 (64%), ADC/ADSCC 296 (34%)	CCRT 907 of 1,583 (57%)	
	IB–IIB (1,581)	SCC 1211 (73%), ADC/ADSCC 351 (21%)		
All combined	IB–IIB (805)	SCC 553 (64%), ADC/ADSCC 296 (34%)	CCRT 907 of 1,652 (55%)	
	IB–IIB (1,581)	SCC 1211 (73%), ADC/ADSCC 351 (21%)		

AC, adjuvant chemotherapy; ADC, adenocarcinoma; ADSCC, adenosquamous carcinoma; AR, adjuvant radiotherapy; CCRT, concurrent chemoradiotherapy; CDDP, cisplatin; FP, fluorouracil and cisplatin; NRS, non-randomized study; NTP, nontaxane and platinum; RCT, randomized controlled trial; RT, radiation therapy; SCC, squamous cell carcinoma; TP, taxane and platinum.

2. Study characteristics

Table 1 lists the study characteristics. Thirteen papers, published between 1996 and 2017, included 942 patients who received AC after RH and 1,721 who received AR after RH. Two studies were RCTs [29,31] and 11 were NRSs [19-28,30]. The studies were conducted in Japan [19-21,24,25,28,30], Korea [23,27], China [22], Italy [26], Austria [29], and the US [31]. For the RCTs, the quality score was 3 (Jadad/Oxford quality scoring system, **Supplementary Table 2**). For the NRSs, the quality scores were 6 or 7 (**Supplementary Table 3**). All included NRSs received three stars for selection. Three studies received one star for comparability for 1 controlled confounder (e.g., disease status after CCRT); eight studies received two stars because they satisfied additional comparability criteria. Nine studies lost stars for exposure for failure to adequately present follow-up losses.

All of the patients had International Federation of Gynecology and Obstetrics (FIGO) stage IB–IIB disease. The median ages in the AC and AR groups were 48 and 49 years, respectively, with median follow-up periods of 66 months. The SCC and ADC histology types were included in 9 studies [19-23,25,27,30,31]; three involved only the SCC type [26,28,29] and one involved only the ADC type [24]. In all studies combined, the AC and AR groups showed similar distributions of pathological risk factors such as bulky tumor (44% vs. 41%), LSVI (69% vs. 70%), DSI (63% vs. 66%), and RMI (1% vs. 2%); however, the AC group showed slightly lower rates of LNM (64% vs. 75%) and PMI (25% vs. 38%) compared with those in the AR group.

Regarding the indication for adjuvant treatment, all enrolled patients had high-risk factors for recurrence in three studies [20,21,26] and only intermediate-risk factors in two studies [22,27], whereas the other studies featured a mixture of intermediate- and high-risk factors [19,23-25,28-31]. Two studies involved only patients with LNM [20,26]. In 4 studies, >1/2 stromal invasion was considered DSI [20,23,24,27]; in 3 studies, >2/3 stromal invasion was considered DSI [22,25,28]; and in 2 studies, >3/4 stromal invasion was considered as DSI [30,31].

Regarding the AC regimen, all included studies involved platinum-based combination chemotherapy; 2 with only platinum + taxane [22,25], 5 with only platinum + non-taxane [26,28-31], and the remaining seven with a mixture of platinum + taxane and platinum + non-taxane [19-21,23,24,27]. Regarding the type of AR, 6 involved RT only without concomitant chemotherapy [25-30] whereas 3 involved CCRT only [21-23] and three featured a mixture of RT and CCRT [19,20,24]. All patients who underwent CCRT received concomitant cisplatin-based chemotherapy.

3. Meta-analysis of the impact of AC on survival compared to that of AR

Twelve studies compared AC to AR in terms of survival, with a combined total of 486 deaths (164/920 patients receiving AC vs. 322/1,608 patients receiving AR) [20-31]. No significant difference in mortality rates was observed between the AC and AR groups (OR=0.73; 95% CI=0.51-1.04; $p=0.08$), with low cross-study heterogeneity ($p=0.10$ and $I^2=34$) (**Fig. 2A**). In the sensitivity analysis, no significant increase was found in mortality in the AC group compared to that in the AR group after any single study was omitted (**Supplementary Table 4**). There was no evidence of publication bias ($p=0.55$, Begg-Mazumdar rank correlation test). The fail-safe N test indicated that 18 more studies with significant findings would be needed to render the two-sided p value non-significant. The funnel plot for publication bias was asymmetric (**Fig. 3A**). However, other quantified tests for publication bias indicated that the observed overall effect is robust ($p=0.55$, Begg-Mazumdar rank correlation test); The fail-safe N test 184 indicated that 18 more studies with significant findings would be needed to render the 2-sided p -value non-significant.

Comparison of adjuvant treatment after surgery




Fig. 2. (A) ORs for the risk of mortality in each study and all studies combined; AC was compared with AR following radical hysterectomy in a meta-analysis based on the random-effects model. (B) ORs for the risk of any recurrence in each study and all studies combined based on the random-effects model. (C) ORs for the risk of local recurrence in each study and all studies combined based on the random-effects model. (D) ORs for the risk of distant recurrence in each study and all studies combined based on the random-effects model. AC, adjuvant chemotherapy; AR, adjuvant radiotherapy; CI, confidence interval; OR, odds ratio.




Fig. 3. (A) Funnel plots for identifying publication bias in the meta-analysis of mortality (n=12). The Begg-Mazumdar rank correlation test indicates no evidence of publication bias (p=0.55). (B) Funnel plots for identifying publication bias in the meta-analysis of any recurrence (n=13). The Begg-Mazumdar rank correlation test indicates no evidence of publication bias (p=0.68). (C) Funnel plots for identifying publication bias in the meta-analysis of local recurrence (n=12). The Begg-Mazumdar rank correlation test indicates no evidence of publication bias (p=0.84). (D) Funnel plots for identifying publication bias in the meta-analysis of distant recurrence (n=12). The Begg-Mazumdar rank correlation test indicates no evidence of publication bias (p=0.55). OR, odds ratio.

4. AC and mortality risk in subgroup meta-analyses

Table 2 shows the results of subgroup analyses comparing mortality rates for AC and AR. The Forest plots are shown in **Supplementary Fig. 1**. **Supplementary Fig. 1A** illustrates the ORs for AC and mortality for each study and the pooled ORs for study design (NRS or RCT). There were 2 RCTs [29,31]; the pooled OR was 0.64 (95% CI=0.26–1.56; p=0.33; p=0.67 and I²=0). There were 10 NRSs [20–28,30]; the pooled OR was 0.73 (95% CI=0.49–1.08; p=0.12; p=0.06 and I²=40), indicating no significant difference in mortality rates between groups.

Comparison of adjuvant treatment after surgery
Table 2. Subgroup analysis comparing AC versus AR following radical hysterectomy for mortality

Characteristics	No. of studies	AC patients (No.)	AR patients (No.)	OR (95% CI) for mortality	p value	Study heterogeneity	
						I ²	p value, Cochran Q
Study design							
RCT	2	72	69	0.64 (0.26–1.56)	0.33	0%	0.67
NRS	10	848	1,539	0.73 (0.49–1.08)	0.12	40%	0.06
Histology type							
SCC	3	183	202	0.63 (0.39–1.01)	0.054	0%	0.27
ADC	2	86	92	0.54 (0.26–1.12)	0.10	0%	0.42
All	8	673	1,337	0.78 (0.48–1.26)	0.31	36%	0.17
Indications for adjuvant treatment							
High-risk	6	554	1,100	0.92 (0.65–1.29)	0.63	24%	0.25
Intermediate-risk	2	103	110	0.66 (0.08–5.48)	0.70	0%	0.62
High- or intermediate-risk	4	157	263	0.41 (0.14–1.17)	0.10	51%	0.12
AR details							
CCRT	4	506	821	0.96 (0.55–1.68)	0.89	29%	0.34
RT	7	350	465	0.63 (0.43–0.92)	0.02	0%	0.40
CCRT or RT	1	64	69	0.63 (0.28–1.39)		NA	NA
AC details							
Platinum combination with taxane only	2	97	117	0.28 (0.03–2.44)	0.25	37%	0.21
Platinum combination with non-taxane only	5	280	374	0.67 (0.45–0.99)	0.04	0%	0.50
Mixed	5	543	1,117	0.84 (0.50–1.42)	0.51	41%	0.19
Status of LNM							
LNM	5	505	1,012	0.85 (0.53–1.38)	0.52	46%	0.15
No LNM	2	103	110	0.66 (0.08–5.48)	0.70	0%	0.62
Mixed	5	194	337	0.62 (0.36–1.06)	0.08	0%	0.20

AC, adjuvant chemotherapy; ADC, adenocarcinoma; AR, adjuvant radiotherapy; CCRT, concurrent chemoradiotherapy; CI, confidence interval; LNM, lymph node metastasis; NA, not applicable; NRS, non-randomized study; OR, odds ratio; SCC, squamous cell carcinoma; RCT, Randomized controlled trial; RT, radiotherapy.

Supplementary Fig. 1B shows the pooled ORs for mortality according to histology type (SCC or ADC [including adenosquamous carcinoma] or SCC plus ADC [including adenosquamous carcinoma]). In three studies including only patients with SCC type [26,28,29], there was no significant difference in mortality between groups (OR=0.63; 95% CI=0.39–1.01; p=0.054; p=0.27 and I²=0). Two studies included only patients with ADC [24] and provided mortality data according to ADC [19]. There was no significant difference in the mortality between the groups (OR=0.54; 95% CI=0.26–1.12; p=0.10; p=0.42 and I²=0). Similarly, no significant differences in mortality were noted between groups (OR=0.78; 95% CI=0.48–1.26; p=0.31; p=0.17 and I²=36) in eight studies including patients with the ADC plus SCC histological type [20-23,25,27,30,31].

Supplementary Fig. 1C shows the pooled ORs for mortality according to the indications for adjuvant treatment. Six studies included only patients with high risk [20,21,26] or provided mortality data according to high risk [23,24,29]. Pooled data from these studies revealed no significant difference between the AC and AR groups (OR=0.92; 95% CI=0.65–1.29; p=0.63), with low cross-study heterogeneity (p=0.25 and I²=24). In the two studies that included only intermediate risk [22,27], no significant difference in recurrence was found between the two groups (OR=0.66; 95% CI=0.08–5.48; p=0.70), with low cross-study heterogeneity (p=0.62 and I²=0).

Supplementary Fig. 1D shows the pooled ORs for mortality according to AR type. In seven studies featuring RT only [25-31], there was no significant increase in mortality in the AC group relative to that in the control group (OR=0.63; 95% CI=0.43–0.92; p=0.02; p=0.40 and I²=0). Four studies included only patients with CCRT as comparison [21-23] and provided mortality data according to CCRT [20]. Pooled data from these studies revealed no

significant difference between the AC and AR groups (OR=0.96; 95% CI=0.55–1.68; $p=0.89$), with low cross-study heterogeneity ($p=0.34$ and $I^2=29$).

Supplementary Fig. 1E shows the pooled ORs for mortality according to the AC regimen. In 2 studies featuring platinum combination with taxane only [22,24], there was no significant difference in mortality between groups (OR=0.28; 95% CI=0.03–2.44; $p=0.25$; $p=0.21$ and $I^2=37$). In 5 studies featuring platinum combination with non-taxane only [26,28–31], there was no significant increase in mortality in AC group relative to the control group (OR=0.67; 95% CI=0.45–0.99; $p=0.04$; $p=0.50$ and $I^2=0$).

Supplementary Fig. 1F shows the pooled ORs for mortality according to the LNM status. Five studies included only patients with LNM [20,26] and provided mortality data according to LNM [23,24,29]. Pooled data from these studies revealed no significant difference between the AC and AR groups (OR=0.85; 95% CI=0.53–1.38; $p=0.52$; $p=0.15$ and $I^2=46$).

5. Meta-analysis of the impact of AC on recurrence compared with that for AR

Thirteen studies comprised a total of 2,663 patients with a combined total of 754 recurrences (253/942 patients receiving AC vs. 501/1,721 patients receiving AR) [19–31]. No significant difference in recurrence rates was observed between the AC and AR groups (OR=0.86; 95% CI=0.67–1.11; $p=0.26$), with low cross-study heterogeneity ($p=0.08$ and $I^2=20$) (**Fig. 2B**). In the sensitivity analysis, no study significantly influenced the pooled OR for AC and recurrence (**Supplementary Table 5**). No publication bias was found ($p=0.68$, Begg-Mazumdar rank correlation test) and the funnel plot was symmetrical (**Fig. 3B**).

Twelve studies assessing 2,530 patients had reported on local and distant recurrence [19–23,25–31]. No significant difference in the local recurrence rates was observed between the two groups (OR=1.20; 95% CI=0.74–1.94; $p=0.46$), with significant cross-study heterogeneity ($p=0.02$ and $I^2=50$) (**Fig. 2C**). No publication bias was found ($p=0.84$) and the funnel plot was symmetrical (**Fig. 3C**). The results for distant recurrence favored the AC group (OR=0.69; 95% CI=0.54–0.88; $p=0.03$), with low cross-study heterogeneity ($p=0.76$ and $I^2=0$) (**Fig. 2D**). No publication bias was found ($p=0.55$) and the funnel plot was symmetrical (**Fig. 3D**). The fail-safe N test indicated that 14 more studies with significant findings would be needed to render the two-sided p value non-significant.

6. Meta-analysis of the impact of AC on recurrence compared with that for AR

Table 3 shows the results of subgroup analyses comparing recurrence in AC versus that for AR. The Forest plots are illustrated in **Supplementary Fig. 2**. **Supplementary Fig. 2A** shows the ORs for AC, the recurrence for each study, and the pooled ORs for study design (NRS or RCT). There were 2 RCTs [29,31]; the pooled OR was 1.20 (95% CI=0.53–2.70, $p=0.66$; $p=0.35$ and $I^2=0$). There were 11 NRSs [19–28,30]; the pooled OR was 0.83 (95% CI=0.63–1.09; $p=0.18$; $p=0.06$ and $I^2=27$), indicating no significant difference in recurrence rates between groups.

Supplementary Fig. 2B shows the pooled ORs for recurrence according to histology type (SCC or ADC [including adenosquamous carcinoma] or SCC plus ADC [including adenosquamous carcinoma]). In 3 studies including only patients with SCC type [26,28,29], there was no significant difference in recurrence between groups (OR=0.51; 95% CI=0.09–3.01; $p=0.46$; $p=0.02$ and $I^2=83$). In eight studies including patients with the ADC plus SCC histological type [20–23,25,27,30,31], there was also no significant difference in recurrence between groups (OR=1.06; 95% CI=0.85–1.32; $p=0.60$; $p=0.62$ and $I^2=0$). Two studies

Comparison of adjuvant treatment after surgery
Table 3. Subgroup analysis comparing AC versus AR following radical hysterectomy for recurrence

Characteristics	Number of studies	AC patients (No.)	AR patients (No.)	OR (95% CI) for recurrence	p value	Study heterogeneity	
						I ²	p value, Cochran Q
Study design							
RCT	2	72	69	1.20 (0.53–2.70)	0.66	0%	0.35
NRS	11	870	1,652	0.83 (0.63–1.09)	0.18	27%	0.06
Histology type							
SCC	3	183	202	0.51 (0.09–3.01)	0.46	83%	0.02
ADC	2	86	92	0.42 (0.17–1.06)	0.07	34%	0.22
All	8	673	1,337	1.06 (0.85–1.32)	0.60	0%	0.62
Indications for adjuvant treatment							
High-risk	4	510	1,045	0.93 (0.67–1.30)	0.69	31%	0.32
Intermediate-risk	2	103	110	2.13 (0.45–10.01)	0.34	25%	0.25
High- or intermediate-risk	5	179	376	0.63 (0.30–1.30)	0.21	52%	0.06
AR details							
CCRT	4	506	821	1.04 (0.81–1.35)	0.74	0%	0.37
RT	7	350	465	0.82 (0.55–1.23)	0.33	19%	0.10
CCRT or RT	2	86	182	0.54 (0.29–1.01)	0.054	0%	0.72
AC details							
Platinum combination with taxane only	2	97	117	1.45 (0.16–12.93)	0.74	70%	0.07
Platinum combination with non-taxane only	5	280	374	0.84 (0.46–1.53)	0.57	49%	0.04
Mixed	6	565	1,230	0.87 (0.62–1.21)	0.40	24%	0.41
Status of LNM							
LNM	3	473	971	0.94 (0.64–1.40)	0.77	45%	0.19
No LNM	2	103	110	2.13 (0.45–10.01)	0.34	25%	0.25
Mixed	6	216	450	0.72 (0.44–1.20)	0.21	27%	0.10

AC, adjuvant chemotherapy; ADC, adenocarcinoma; AR, adjuvant radiotherapy; CCRT, concurrent chemoradiotherapy; CI, confidence interval; LNM, lymph node metastasis; NRS, non-randomized study; OR, odds ratio; RCT, Randomized controlled trial; RT, radiotherapy; SCC, squamous cell carcinoma.

included only patients with ADC [24] and provided recurrence data according to ADC [19]. There was also no significant difference in recurrence between the group (OR=0.42; 95% CI=0.17–1.06; p=0.07; p=0.22 and I²=34).

Supplementary Fig. 2C shows the pooled ORs for recurrence according to the indications for adjuvant treatment. Four studies included only patients with high risk [20,21,26] or provided recurrence data according to high risk [23]. Pooled data from these studies revealed no significant difference between the AC and AR groups (OR=0.93; 95% CI=0.67–1.30; p=0.69), with low cross-study heterogeneity (p=0.32 and I²=31). In 2 studies that only assessed intermediate risk [22,27], no significant difference in recurrence was found between the 2 groups (OR=2.13; 95% CI=0.45–10.01; p=0.34), with low cross-study heterogeneity (p=0.25 and I²=25).

Supplementary Fig. 2D shows the pooled ORs for recurrence according to AR type. In 7 studies featuring RT only [25–31], there was no significant difference in recurrence between the 2 groups (OR=0.82; 95% CI=0.55–1.23; p=0.33; p=0.10 and I²=19). Four studies included only patients receiving CCRT as comparison [21–23] and provided recurrence data according to CCRT [20]. Pooled data from these studies revealed no significant difference between the AC and AR groups (OR=1.04; 95% CI=0.81–1.35; p=0.74), with low cross-study heterogeneity (p=0.37 and I²=0).

Supplementary Fig. 2E shows the pooled ORs for recurrence according to the AC regimen. In 2 studies featuring platinum combination with taxane only [22,25], there was no significant difference in recurrence between groups (OR=1.45; 95% CI=0.16–12.93; p=0.74; p=0.07 and I²=70). In 5 studies featuring platinum combination with non-taxane only [26,28–31], there

was no significant difference in recurrence between groups (OR=0.84; 95% CI=0.46–1.53; $p=0.57$; $p=0.04$ and $I^2=49$).

Supplementary Fig. 2F shows the pooled ORs for recurrence according to the LNM status. Three studies included only patients with LNM [20,26] and provided recurrence data according to LNM [23]. Pooled data from these studies revealed no significant difference between the AC and AR groups (OR=0.94; 95% CI=0.64–1.40; $p=0.77$; $p=0.19$ and $I^2=45$).

DISCUSSION

To the best of our knowledge, this is the first meta-analysis to comprehensively evaluate the effects of AC on oncologic outcome in cervical cancer patients with intermediate- or high-risk factors after surgery compared to those for AR. The results of our meta-analysis indicate that patients receiving AC after surgery have comparable survival outcomes and recurrence rates to those who undergo AR or adjuvant CCRT. This pattern was consistently observed in the subgroup analyses of study design, histology, AR type, indications for adjuvant treatment, LNM status, and AC regimen.

It is controversial as to whether histologic type influences survival and the pattern of recurrence in patients with early-stage cervical cancer [32–35]. Although studies are limited, ADC may be more resistant when RT is added after surgery, and role of adjuvant RT for ADC remains uncertain [36]. The results of subgroup analysis including ADC plus SCC histologic type showed no significant difference in recurrence and mortality rates between groups.

Although few data specifically address the result of AC after radical surgery in patients meeting the Sedlis criteria, the literature addressing this issue reported a 3-year disease-free survival rate of 94.6% and a 5-year cervical cancer-specific survival of 96.2% [37]. Due to discretion and experience of the gynecologic oncologists, the risk factors that indicate adjuvant treatment can differ between clinicians. In each of the analyzed studies, each group was compared with the same indications. The pooled data from these studies revealed no significant difference in mortality and recurrence rates between the AC and AR groups.

The results of retrospective studies suggested that adjuvant systemic chemotherapy treatment decreased distant recurrence when compared to CCRT for patients with high-risk factors [19,20]. However, other studies reported that recurrence patterns and rates did not differ significantly between groups with high-risk factors [21,23,29,31]. In a study of patients with intermediate risk factors, distant control was not significantly improved but local failure did not occur in the chemotherapy group [27]. In our analysis, no significant difference in local recurrence rates was observed between groups. However, distant recurrence favored the AC group (OR=0.69; 95% CI=0.54–0.88; $p=0.03$), with low cross-study heterogeneity ($p=0.76$ and $I^2=0$). If lymph node metastasis is a surrogate of distant metastasis, this result may be evidence of a direct effect of systemic chemotherapy on distant micrometastatic lesions.

Platinum is the most common active agent in the treatment of cervical cancer and platinum-containing combination regimens, topotecan and paclitaxel combination regimens, platinum single agent, and paclitaxel single agent have been considered appropriate treatments for advanced or recurrent cervical cancer [38]. Although the optimal chemotherapeutic agents

have not been determined, AC with platinum-based combination regimen were used in the included studies. The GOG 240 trial demonstrated the effects of the addition of bevacizumab to paclitaxel and cisplatin [39]. Similarly, in the Japanese Gynecologic Oncology Group (JGOG) 1,065 trial, neoadjuvant chemotherapy with nedaplatin and irinotecan were reported to have favorable response rates [40]. These data show the potential of the addition of such combination regimens in the adjuvant setting. There were no significant differences in mortality among 2 studies including platinum and taxane combination therapy (OR=0.28; 95% CI=0.03–2.44; p=0.25). In addition, the mortality in the AC group did not increase significantly in five studies that using platinum and non-taxane combinations (OR=0.67; 95% CI=0.45–0.99; p=0.04). The recurrence rates between groups did not differ significantly, irrespective of the combination regimen.

RCTs overcome numerous weaknesses associated with NRSs and provide the best available data regarding the effect of AC in surgically treated cervical cancer patients. Our meta-analysis included 2 RCTs with 3 stars in the Jadad scoring system. However, the study by Curtin et al. [31] was closed prematurely owing to insufficient accrual; the initial accrual goal was 160 patients but it enrolled only 89 patients. Thus, the data were unable to show that the AC and AR were equivalent. Similarly, the planned accrual goal after statistical sample size calculation was 160 patients for the study by Lahousen et al. [29]. Their final enrolled number was only 52, yielding suboptimal power. Recently, the JGOG launched a confirmatory clinical trial of AC in high-risk, stage IB–IIB cervical cancer after primary radical hysterectomy (JGOG 1082). If the non-inferiority of AC is confirmed in terms of survival compared with AR in this prospective trial, AC could be accepted as the new standard treatment.

Our study had several limitations; thus, the results should be interpreted with caution. First, the majority of the included studies were NRS, although 2 RCTs were also included. While we rigorously performed subgroup-analyses to control for potential confounders, unknown confounders such as surgical quality may not have been excluded. Second, the studies were conducted in different institutions with presumably varying surgical expertise. Third, it was not possible to perform a subgroup analysis of the number of AC cycles or dose of AC because of a lack of data. Therefore, the effect of AC according to these features could not be analyzed. Finally, our meta-analysis did not provide adverse events profiles related to AC or AR. No RCT directly compared adverse events between groups, but several studies suggested lower complication rates with AC [17,20,24].

The results of the present meta-analysis suggest that AC showed similar survival outcomes in surgically treated cervical cancer patients and reduced the risk of distant recurrence when compared with AR. The JGOG trial may provide more evidence on the role of AC in these patients, establishing this strategy be used as an alternative standard treatment option.

SUPPLEMENTARY MATERIALS

Supplementary Table 1

Excluded studies with reasons

[Click here to view](#)

Supplementary Table 2

Jadad score for the risk of bias and quality assessment of RCTs

[Click here to view](#)

Supplementary Table 3

Newcastle-Ottawa scale for the risk of bias and quality assessment of non-randomized studies

[Click here to view](#)

Supplementary Table 4

Sensitivity analysis comparing adjuvant chemotherapy (AC) versus adjuvant radiotherapy (AR) for mortality in which one study at a time was omitted and the pooled odds ratio for all remaining studies was calculated

[Click here to view](#)

Supplementary Table 5

Sensitivity analysis comparing AC versus AR for recurrence in which one study at a time was omitted and the pooled OR for all remaining studies was calculated

[Click here to view](#)

Supplementary Fig. 1

The association between AC and mortality in subgroup meta-analyses is shown according to (A) study design, (B) histology type, (C) indication for adjuvant treatment, (D) AR type, (E) AC type, and (E) status of LNM. The size of each square is proportional to the sample size in each study, and the horizontal line through the square indicates the 95% CI for that study. For the pooled analysis, the diamond indicates the pooled value, and the right and left ends of the diamond indicate the 95% CI for the analysis.

[Click here to view](#)

Supplementary Fig. 2

The association between AC and recurrence in subgroup meta-analyses is shown according to (A) study design, (B) histology type, (C) indication for adjuvant treatment, (D) AR type, (E) AC type, and (E) status of LNM. The size of each square is proportional to the sample size in each study, and the horizontal line through the square indicates the 95% CI for that study. For the pooled analysis, the diamond indicates the pooled value, and the right and left ends of the diamond indicate the 95% CI for the analysis.

[Click here to view](#)

REFERENCES

1. International Agency for Research on Cancer; World Health Organization. GLOBOCAN 2012: estimated cancer incidence, mortality, and prevalence worldwide in 2012 [Internet]. Lyon: IARC; 2014 [cited year month day]. Available from: http://globocan.iarc.fr/Pages/fact_sheets_cancer.aspx.

2. Shim SH, Kim H, Sohn IS, Hwang HS, Kwon HS, Lee SJ, et al. Nationwide cervical cancer screening in Korea: data from the National Health Insurance Service Cancer Screening Program and National Cancer Screening Program, 2009–2014. *J Gynecol Oncol* 2017;28:e63.
[PUBMED](#) | [CROSSREF](#)
3. Ryu HS, Kang SB, Kim KT, Chang KH, Kim JW, Kim JH. Efficacy of different types of treatment in FIGO stage IB2 cervical cancer in Korea: results of a multicenter retrospective Korean study (KGOG-1005). *Int J Gynecol Cancer* 2007;17:132-6.
[PUBMED](#) | [CROSSREF](#)
4. Peters WA 3rd, Liu PY, Barrett RJ 2nd, Stock RJ, Monk BJ, Berek JS, et al. Concurrent chemotherapy and pelvic radiation therapy compared with pelvic radiation therapy alone as adjuvant therapy after radical surgery in high-risk early-stage cancer of the cervix. *J Clin Oncol* 2000;18:1606-13.
[PUBMED](#) | [CROSSREF](#)
5. Sedlis A, Bundy BN, Rotman MZ, Lentz SS, Muderspach LI, Zaino RJ. A randomized trial of pelvic radiation therapy versus no further therapy in selected patients with stage IB carcinoma of the cervix after radical hysterectomy and pelvic lymphadenectomy: a Gynecologic Oncology Group Study. *Gynecol Oncol* 1999;73:177-83.
[PUBMED](#) | [CROSSREF](#)
6. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000;283:2008-12.
[PUBMED](#) | [CROSSREF](#)
7. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
[PUBMED](#) | [CROSSREF](#)
8. Shim SH, Kim SN, Jung PS, Dong M, Kim JE, Lee SJ. Impact of surgical staging on prognosis in patients with borderline ovarian tumours: a meta-analysis. *Eur J Cancer* 2016;54:84-95.
[PUBMED](#) | [CROSSREF](#)
9. Wells GA, Shea B, O'connell D, Peterson J, Welch V, Losos M, et al., editors. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. The 3rd Symposium on Systematic Reviews: Beyond the Basics; 2000 Jul 3–5; Oxford, United Kingdom. Ottawa: Ottawa Hospital Research Institute; 2000.
10. Aziz O, Constantinides V, Tekkis PP, Athanasiou T, Purkayastha S, Paraskeva P, et al. Laparoscopic versus open surgery for rectal cancer: a meta-analysis. *Ann Surg Oncol* 2006;13:413-24.
[PUBMED](#) | [CROSSREF](#)
11. OuYang PYXie C, Mao YP, Zhang Y, Liang XX, Su Z, et al. Significant efficacies of neoadjuvant and adjuvant chemotherapy for nasopharyngeal carcinoma by meta-analysis of published literature-based randomized, controlled trials. *Ann Oncol* 2013;24:2136-46.
[PUBMED](#) | [CROSSREF](#)
12. Jadad AR, Moore RA, Carroll D, Jenkinson C, Reynolds DJ, Gavaghan DJ, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996;17:1-12.
[PUBMED](#) | [CROSSREF](#)
13. Dickersin K, Berlin JA. Meta-analysis: state-of-the-science. *Epidemiol Rev* 1992;14:154-76.
[PUBMED](#) | [CROSSREF](#)
14. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557-60.
[PUBMED](#) | [CROSSREF](#)
15. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177-88.
[PUBMED](#) | [CROSSREF](#)
16. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50:1088-101.
[PUBMED](#) | [CROSSREF](#)
17. Sterne JA, Egger M. Funnel plots for detecting bias in meta-analysis: guidelines on choice of axis. *J Clin Epidemiol* 2001;54:1046-55.
[PUBMED](#) | [CROSSREF](#)
18. Orwin RG. A fail-safe N for effect size in meta-analysis. *J Educ Stat* 1983;8:157-9.
19. Seki T, Tanabe H, Nagata C, Suzuki J, Suzuki K, Takano H, et al. Adjuvant therapy after radical surgery for stage IB-IIB cervical adenocarcinoma with risk factors. *Jpn J Clin Oncol* 2017;47:32-8.
[PUBMED](#) | [CROSSREF](#)

20. Matsuo K, Shimada M, Aoki Y, Sakamoto M, Takeshima N, Fujiwara H, et al. Comparison of adjuvant therapy for node-positive clinical stage IB-IIB cervical cancer: systemic chemotherapy versus pelvic irradiation. *Int J Cancer* 2017;141:1042-51.
[PUBMED](#) | [CROSSREF](#)
21. Takekuma M, Kasamatsu Y, Kado N, Kuji S, Tanaka A, Takahashi N, et al. Adjuvant chemotherapy versus concurrent chemoradiotherapy for high-risk cervical cancer after radical hysterectomy and systematic lymphadenectomy. *Int J Clin Oncol* 2016;21:741-7.
[PUBMED](#) | [CROSSREF](#)
22. Li L, Song X, Liu R, Li N, Zhang Y, Cheng Y, et al. Chemotherapy versus radiotherapy for FIGO stages IB1 and IIA1 cervical carcinoma patients with postoperative isolated deep stromal invasion: a retrospective study. *BMC Cancer* 2016;16:403.
[PUBMED](#) | [CROSSREF](#)
23. Jung PS, Kim DY, Lee SW, Park JY, Suh DS, Kim JH, et al. Clinical role of adjuvant chemotherapy after radical hysterectomy for FIGO stage IB-IIA cervical cancer: comparison with adjuvant RT/CCRT using inverse-probability-of-treatment weighting. *PLoS One* 2015;10:e0132298.
[PUBMED](#) | [CROSSREF](#)
24. Shimada M, Nishimura R, Hatae M, Hiura M, Takehara K, Tase T, et al. Comparison of adjuvant chemotherapy and radiotherapy in patients with cervical adenocarcinoma of the uterus after radical hysterectomy: SGSG/TGCU Intergroup surveillance. *Eur J Gynaecol Oncol* 2013;34:425-8.
[PUBMED](#)
25. Hosaka M, Watari H, Takeda M, Moriwaki M, Hara Y, Todo Y, et al. Treatment of cervical cancer with adjuvant chemotherapy versus adjuvant radiotherapy after radical hysterectomy and systematic lymphadenectomy. *J Obstet Gynaecol Res* 2008;34:552-6.
[PUBMED](#) | [CROSSREF](#)
26. Mossa B, Mossa S, Marziani R. Adjuvant chemotherapy versus radiation therapy after radical surgery in high-risk positive node stage IB/IIA cervical cancer. *Eur J Gynaecol Oncol* 2010;31:545-50.
[PUBMED](#)
27. Lee KB, Lee JM, Ki KD, Lee SK, Park CY, Ha SY. Comparison of adjuvant chemotherapy and radiation in patients with intermediate risk factors after radical surgery in FIGO stage IB-IIA cervical cancer. *Int J Gynecol Cancer* 2008;18:1027-31.
[PUBMED](#) | [CROSSREF](#)
28. Hosaka M, Watari H, Kato T, Odagiri T, Konno Y, Endo D, et al. Clinical efficacy of paclitaxel/cisplatin as an adjuvant chemotherapy for patients with cervical cancer who underwent radical hysterectomy and systematic lymphadenectomy. *J Surg Oncol* 2012;105:612-6.
[PUBMED](#) | [CROSSREF](#)
29. Lahousen M, Haas J, Pickel H, Hackl A, Kurz C, Ogris H, et al. Chemotherapy versus radiotherapy versus observation for high-risk cervical carcinoma after radical hysterectomy: a randomized, prospective, multicenter trial. *Gynecol Oncol* 1999;73:196-201.
[PUBMED](#) | [CROSSREF](#)
30. Iwasaka T, Kamura T, Yokoyama M, Matsuo N, Nakano H, Sugimori H. Adjuvant chemotherapy after radical hysterectomy for cervical carcinoma: a comparison with effects of adjuvant radiotherapy. *Obstet Gynecol* 1998;91:977-81.
[PUBMED](#)
31. Curtin JP, Hoskins WJ, Venkatraman ES, Almadrones L, Podratz KC, Long H, et al. Adjuvant chemotherapy versus chemotherapy plus pelvic irradiation for high-risk cervical cancer patients after radical hysterectomy and pelvic lymphadenectomy (RH-PLND): a randomized phase III trial. *Gynecol Oncol* 1996;61:3-10.
[PUBMED](#) | [CROSSREF](#)
32. Look KY, Brunetto VL, Clarke-Pearson DL, Averette HE, Major FJ, Alvarez RD, et al. An analysis of cell type in patients with surgically staged stage IB carcinoma of the cervix: a Gynecologic Oncology Group study. *Gynecol Oncol* 1996;63:304-11.
[PUBMED](#) | [CROSSREF](#)
33. Balbergen A, Ewing-Graham PC, Hop WC, Struijk P, Helmerhorst TJ. Prognostic factors in adenocarcinoma of the uterine cervix. *Gynecol Oncol* 2004;92:262-7.
[PUBMED](#) | [CROSSREF](#)
34. Park JY, Kim DY, Kim JH, Kim YM, Kim YT, Nam JH. Outcomes after radical hysterectomy in patients with early-stage adenocarcinoma of uterine cervix. *Br J Cancer* 2010;102:1692-8.
[PUBMED](#) | [CROSSREF](#)
35. Galic V, Herzog TJ, Lewin SN, Neugut AI, Burke WM, Lu YS, et al. Prognostic significance of adenocarcinoma histology in women with cervical cancer. *Gynecol Oncol* 2012;125:287-91.
[PUBMED](#) | [CROSSREF](#)

36. Hopkins MP, Peters WA 3rd, Andersen W, Morley GW. Invasive cervical cancer treated initially by standard hysterectomy. *Gynecol Oncol* 1990;36:7-12.
[PUBMED](#) | [CROSSREF](#)
37. Lee KB, Kim YS, Lee JM. Oncologic outcomes of adjuvant chemotherapy alone after radical surgery for stage IB-IIA cervical cancer patients. *J Gynecol Oncol* 2018;29:e5.
[PUBMED](#) | [CROSSREF](#)
38. National Comprehensive Cancer Network (US). NCCN Clinical Practice Guidelines in Oncology. Cervical cancer, version 1.2018 [Internet]. Fort Washington, PA: National Comprehensive Cancer Network; 2018 [cited 2018 Jan 6]. Available from: https://www.nccn.org/professionals/physician_gls/pdf/cervical.pdf.
39. Tewari KS, Sill MW, Long HJ 3rd, Penson RT, Huang H, Ramondetta LM, et al. Improved survival with bevacizumab in advanced cervical cancer. *N Engl J Med* 2014;370:734-43.
[PUBMED](#) | [CROSSREF](#)
40. Yamaguchi S, Nishimura R, Yaegashi N, Kiguchi K, Sugiyama T, Kita T, et al. Phase II study of neoadjuvant chemotherapy with irinotecan hydrochloride and nedaplatin followed by radical hysterectomy for bulky stage Ib2 to IIb, cervical squamous cell carcinoma: Japanese Gynecologic Oncology Group study (JGOG 1065). *Oncol Rep* 2012;28:487-93.
[PUBMED](#) | [CROSSREF](#)

Appendix 1. The Search Strategy

PUBMED		
#1	"Uterine Cervical Neoplasms"[Mesh]	67,380
#2	(cervical[TIAB] OR cervix[TIAB]) AND (Neoplasm*[TIAB] OR cancer*[TIAB] OR tumor[TIAB] OR tumors[TIAB] OR tumour*[TIAB] OR carcinoma*[TIAB] OR Malignanc*[TIAB] OR Neoplasia*[TIAB] OR Adenocarcinoma*[TIAB])	96,969
#3	#1 OR #2	112,551
#4	"Postoperative Period"[Mesh] OR "Postoperative Complications"[Mesh] OR "Postoperative Care"[Mesh]	559,967
#5	Postoperativ*[TIAB] OR Post-operativ*[TIAB]	497,845
#6	surger*[TIAB] OR Surgical[TIAB]	1,577,480
#7	"Hysterectomy"[Mesh] OR Hysterectom*[TIAB]	43,368
#8	#4 OR #5 OR #6 OR #7	2,033,382
#9	"Chemotherapy, Adjuvant"[Mesh]	35,097
#10	Adjuvant*[TIAB] OR addition*[TIAB]	2,294,559
#11	Chemotherap*[TIAB] OR Pharmacotherap*[TIAB] OR "Drug therapy"[TIAB] OR "Drug therapies"[TIAB]	417,878
#12	#10 AND #11	85,746
#13	#9 OR #12	108,009
#14	#3 AND #8 AND #13	1,532
#15	#14 AND English[Lang]	1,283
EMBASE		
#1	'uterine cervix cancer'/exp	80,905
#2	((cervical OR cervix) NEAR/5 (Neoplasm* OR cancer* OR tumor* OR tumour* OR carcinoma* OR Adenocarcinoma* OR Malignanc* OR Neoplasia*)):ab,ti	95,438
#3	#1 OR #2	117,217
#4	'postoperative period'/exp OR 'postoperative care'/exp OR 'postoperative complication'/exp	942,737
#5	Postoperativ*:ab,ti OR Post-operativ*:ab,ti	660,468
#6	surger*:ab,ti OR Surgical:ab,ti	2,022,870
#7	'hysterectomy'/exp OR Hysterectom*:ab,ti	71,538
#8	#4 OR #5 OR #6 OR #7	2,741,955
#9	'adjuvant chemotherapy'/exp	39,846
#10	((Adjuvant* OR addition*) NEAR/5 (Chemotherap* OR Pharmacotherap* OR 'Drug therapy' OR 'Drug therapies')):ab,ti	52,410
#11	#9 OR #10	75,340
#12	#3 AND #8 AND #11	1,376
#13	#12 AND [english]/lim	1,218
Cochrane		
#1	MeSH descriptor: [Uterine Cervical Neoplasms] explode all trees	2,000
#2	(cervical:ab,ti,kw or cervix:ab,ti,kw) near/5 (Neoplasm*:ab,ti,kw or cancer*:ab,ti,kw or tumor*:ab,ti,kw or tumour*:ab,ti,kw or carcinoma*:ab,ti,kw or Adenocarcinoma*:ab,ti,kw or Malignanc*:ab,ti,kw or Neoplasia*:ab,ti,kw)	4,004
#3	#1 or #2	4,004
#4	MeSH descriptor: [Postoperative Period] explode all trees	5,354
#5	MeSH descriptor: [Postoperative Complications] explode all trees	34,368
#6	MeSH descriptor: [Postoperative Care] explode all trees	4,406
#7	MeSH descriptor: [Hysterectomy] explode all trees	1,843
#8	Postoperativ*:ab,ti,kw or Post-operativ*:ab,ti,kw or surger*:ab,ti,kw or Surgical:ab,ti,kw or Hysterectom*:ab,ti,kw	160,575
#9	#4 or #5 or #6 or #7 or #8	162,152
#10	MeSH descriptor: [Chemotherapy, Adjuvant] explode all trees	3,863
#11	(Adjuvant*:ab,ti,kw or addition*:ab,ti,kw) near/5 (Chemotherap*:ab,ti,kw or Pharmacotherap*:ab,ti,kw or "Drug therapy":ab,ti,kw or "Drug therapies":ab,ti,kw)	9,721
#12	#10 or #11	9,721
#13	#3 and #9 and #12	88
#14	#13 in Cochrane Reviews (Reviews only), Other Reviews and Trials	87

Comparison of adjuvant treatment after surgery
Appendix 2. Detailed characteristics of studies included in the meta-analysis

Study /design	Location/ Treatment study period	Recurrence (No.)	Death of disease (No.)	Median age (yr)	Median follow-up period (Mo)	Stage, No. (%)	Histology, No. (%)	LNM, No. (%)	PMI, No. (%)	RMI, No. (%)	Tumor size, No. (%)	DSI, No. (%)	LWSI, No. (%)	Indication of adjuvant treatment	Detailed reasons for adjuvant treatment	Median EBRT (Gy)	AR details		AC details
																	Radiation type	Concomitant chemotherapy regimen	
Seki et al. /NRS	Japan/ 2001-2010	3	2	47	48	IB11 (50%) II11 (50%)	ADC/ADSCC 22 (100%)	8 (36%)	7 (32%)	0	>4 cm, 17 (77%)	NR	NR	Intermediate- or high-risk	Bulky tumor (>4 cm), PMI, LNM	NR	NR	Platinum + Taxane/ Platinum + non-Taxane	
Matsuo et al. /NRS	Japan/ 2004-2008	114	82	47	64.5	IB1141 (44%) IB2161 (19%) IIA134 (11%) IIB83 (26%) IB1310 (41%) IB2134 (18%) IIA90 (12%) IIB219 (29%)	ADC/ADSCC 23 (20%) SCC 156 (49%) ADC/ADSC 149 (47%)	319 (100%)	112 (35%)	NR	>4 cm, 111 (36%)	241 (85%)	270 (87%)	High-risk	LNM	NR	NR	Platinum + Taxane/ Platinum + non-Taxane	
Takekuma et al. /NRS	Japan/ 2002-2013	10	6	45	52	IB119 (24%) IB2114 (38%) II114 (38%) IB125 (34%) IB222 (30%) II27 (37%)	SCC 24 (65%) ADC/ADSCC 13 (35%) SCC 48 (65%) ADC/ADSCC 26 (35%)	16 (43%)	16 (43%)	NR	>4 cm, 23 (64%)	NR	34 (92%)	High-risk	PMI, LNM	NR	NR	Platinum + Taxane/ Platinum + non-Taxane	
Li et al. /NRS	China/ 2010-2013	5	1	49	34	IB122 (34%) IIA143 (66%) IB128 (41%) IIA140 (59%)	SCC 59 (91%) ADC 6 (9%) SCC 63 (98%) ADC 23 (7%)	0	0	NR	<4 cm, 1/3-2/3 thickness 26 (40%) 65 (100%) 2/3-full thickness 16 (25%) <4 cm, 1/3-2/3 thickness 20 (44%) 68 (100%) 2/3-full thickness 17 (25%)	NR	NR	Intermediate-risk	isolated DSI	NR	NR	Platinum + Taxane	
Jung et al. /NRS	Korea/ 1992-2012	11	3	44	47	IB167 (79%) IB211 (13%) IIA7 (8%) IB121 (68%) IB231 (18%) IIA25 (14%)	SCC 61 (79%) ADC 21 (25%) ADSCC 3 (4%) SCC 138 (78%) ADC 26 (15%) ADSCC 13 (7%)	27 (32%)	8 (10%)	1 (1%)	30 mm (1%)	63 (76%)	46 (54%)	Intermediate- or high-risk	intermediate risk factors (tumor size >4 cm, DSI, LWSI) or ≥ high-risk factor (LNM, RMI, PMI)	45-55	CCRT 177 (100%)	Platinum + Taxane/ Platinum + non-Taxane	
Shimada et al. /NRS	Japan/ 1997-2003	18	13	NR	NR	IB-IB164 (100%) IB-IB164 (100%)	ADC/ADSCC 64 (100%)	13 (20%)	12 (19%)	1 (2%)	>4 cm, 26 (41%)	19 (30%)	26 (41%)	Intermediate- or high-risk	DSI, LWSI, PMI, LNM, RMI	NR	NR	Platinum + Taxane/ Platinum + non-Taxane	
		28	20	NR	NR	IB-IB164 (100%)	ADC/ADSCC 69 (100%)	27 (39%)	14 (20%)	7 (10%)	>4 cm, 28 (41%)	17 (25%)	35 (51%)	Intermediate- or high-risk	CCRT or RT only	45-50	CCRT or RT only	Platinum + Taxane/ Platinum + non-Taxane	

(continued to the next page)

Comparison of adjuvant treatment after surgery

Appendix 2. (Continued) Detailed characteristics of studies included in the meta-analysis

Study /design	Location/Treatment study period	Recurrence (No.)	Death of disease (No.)	Median age (yr)	Median follow-up period (Mo)	Stage, No. (%)	Histology, No. (%)	LNM, No. (%)	PMI, No. (%)	RMI, No. (%)	Tumor size, No. (%)	DSI, No. (%)	LWSI, No. (%)	Indication of adjuvant treatment	Detailed reasons for adjuvant treatment	AR details		AC details Regimen	
																Median dose of EBRT (Gy)	Radiation type		Concomitant chemotherapy regimen
Hosaka et al. /NRS	Japan /2003-2007	7	1	48	>36	IB1 9 (28%) IB2 8 (28%) IIA 0	SCC 24 (75%) ADC/ADSCC 8 (25%)	>2 LNM 11 (34%)	7 (22%)	0	>4 cm, 25 (78%)	19 (59%) >2/3 thickness	24 (75%) >2/3 thickness	Intermediate- or high-risk	DSI (>2/3 thickness), LWSI, PMI, LNM, tumor diameter >4 cm	50	RT only 49 (100%)	No	Platinum + Taxane
Mossa et al. /NRS	Italy /1971-2001	51	39	47*	120	IB1 9 (18%) IB2 24 (19%) IIA 3 (6%) IIB 25 (51%)	SCC 47 (96%) ADC/ADSCC 2 (4%)	>2 LNM 22 (45%)	16 (33%)	0	>4 cm, 16 (33%)	27 (55%) >2/3 thickness	40 (82%)	High-risk	LNM	50	RT only 49 (100%)	No	Platinum + non-Taxane
Lee et al. /NRS	Korea /1997-2005	3	0	54.5	49	IB1 22 (58%) IB2 10 (26%) IIA 6 (16%) IIB 31 (74%)	SCC 31 (82%) ADC 2 (5%) ADSCC 5 (13%)	0	0	0	>3 cm, 15 (39%)	28 (74%) >2/3 thickness	17 (45%)	Intermediate-risk	DSI, LWSI, large tumor size (3-6 cm), close vaginal resection margin (<1 cm)	45	RT only 42 (100%)	No	Platinum + Taxane/ Platinum + non-Taxane
Hosaka et al. /NRS	Japan /2000-2002	1	0	52*	>36	IB1 11 (39%) IB2 9 (32%) IIA 2 (7%) IIB 6 (21%)	SCC 28 (100%)	3 (11%)	5 (18%)	0	>4 cm, 10 (36%)	11 (39%) >2/3 thickness	27 (96%)	Intermediate- or high-risk	DSI (>2/3 thickness), LWSI, PMI, LNM, tumor diameter >4 cm	50	RT only 42 (100%)	No	Platinum + non-Taxane
Lahousen et al. /RCT	Austria /1989-1995	8	4	51	49	IB-IB	SCC 28 (100%)	19 (68%)	NR	0	NR	NR	28 (100%)	Intermediate- or high-risk	LNM, LWSI	50	RT only 24 (100%)	No	Platinum + non-Taxane
Iwasaka et al. /NRS	Japan /1983-1988	13	9	54*	75	IB 30 (57%) IIA 8 (15%) IIB 15 (28%) IIB 7 (13%)	SCC 43 (81%) ADSCC 3 (6%) others 7 (13%)	22 (42%)	16 (30%)	0	NR	35 (66%) >3/4 thickness	NR	Intermediate- or high-risk	LNM, DSI (< 3/4 thickness), PMI	41.4-50.4	RT only 127 (100%)	No	Platinum + non-Taxane
Curtin et al. /RCT	US /1987-1994	9	6	45*	36	IB 74 (83%) IIA 14 (16%) IIB 11 (12%)	SCC 52 (58%) ADC 30 (39%)	33 (37%)	9 (10%)	NR	>4 cm, 33 (37%)	45 (51%) >3/4 thickness	NR	Intermediate- or high-risk	LNM, tumor diameter >4 cm, DSI (<3/4 thickness), nonsquamous histology, PMI, RMI	45	RT only 45 (100%)	No	Platinum + non-Taxane
RCTs combined		17	10	47†	41†	IB-IB	SCC 104 (74%) ADC 30 (21%)	19 of 28 (68%) 14 of 24 (58%)	9 of 89 (10%) 0 of 24	0 of 28 (0%) 0 of 24	>4 cm, 45 of 89 (51%) >3/4 thickness (33 of 89) (37%)	45 of 89 (51%) >3/4 thickness (24 of 24) (100%)	28 of 89 (31%) 24 of 24 (100%)						CCRT 0

(continued to the next page)

Comparison of adjuvant treatment after surgery

Appendix 2. (Continued) Detailed characteristics of studies included in the meta-analysis

Study /design	Location/Treatment study period	Recurrence (No.)	Death of disease (No.)	Median age (yr)	Median follow-up period (Mo)	Stage, No. (%)	Histology, No. (%)	LNMM, No. (%)	PMI, No. (%)	RMI, No. (%)	Tumor size	DSI, No. (%)	LWSI, No. (%)	Indication of adjuvant treatment	Detailed reasons for adjuvant treatment	AR details		AC details
																EBRT (Gy)	Medication chemotherapy regimen	
NRSs combined	AC (n=870)	236	154 of 848	48†	66†	IB 535 (66%) IIA 151 (19%)	SCC 553 (64%) ADC/ADSCC 296 (34%)	535 of 838 (64%)	183 of 743 (25%)	2 of 449 (1%)	>4 cm, 292 of 658 (44%)	432 of 684 (63%)	444 of 656 (68%)					
	AR (n=1,652)	487	308 of 1,539	49†		IIB 119 (15%) IB 1002 (63%) IIA 280 (18%)	SCC 1211 (73%) ADC/ADSCC 351 (21%)	1,207 of 1,603 (75%)	573 of 1,516 (38%)	16 of 755 (2%)	>4 cm, 543 of 1,339 (41%)	881 of 1,329 (66%)	931 of 1,335 (70%)		CCRT907 of 1,583 (57%)			
	All combined (n=942)	253	164 of 920	48†	64†	IB 535 (66%) IIA 151 (19%)	SCC 553 (64%) ADC/ADSCC 296 (34%)	554 of 866 (64%)	183 of 743 (25%)	2 of 477 (1%)	>4 cm, 292 of 658 (44%)	432 of 684 (63%)	472 of 684 (69%)					
AR (n=1,721)	AR	501	322 of 1,608	49†		IIB 119 (15%) IB 1002 (63%) IIA 280 (18%)	SCC 1211 (73%) ADC/ADSCC 351 (21%)	1,221 of 1,627 (75%)	573 of 1,516 (38%)	16 of 779 (2%)	>4 cm, 543 of 1,339 (41%)	881 of 1,329 (66%)	955 of 1,359 (70%)		CCRT907 of 1,652 (55%)			
	All combined (n=1,721)	501	322 of 1,608	49†		IIB 299 (19%)	SCC 1211 (73%) ADC/ADSCC 351 (21%)	1,221 of 1,627 (75%)	573 of 1,516 (38%)	16 of 779 (2%)	>4 cm, 543 of 1,339 (41%)	881 of 1,329 (66%)	955 of 1,359 (70%)		CCRT907 of 1,652 (55%)			

AC, adjuvant chemotherapy; ADC, adenocarcinoma; ADSCC, adenosquamous carcinoma; AR, adjuvant radiotherapy; CCRT, concurrent chemoradiotherapy; DSI, deep stromal invasion; EBRT, external beam radiotherapy; LNMM, lymph node metastasis; LWSI, lymphovascular space invasion; NR, not reported; NRS, non-randomized study; PMI, parametrial invasion; RCT, randomized controlled trial; RH, radical hysterectomy; RMI, resection margin involved; RT, radiation therapy; SCC, squamous cell carcinoma.
 †Mean; †weighted average.