

## Letter to the Editor

## Smoking and breast cancer

P Terry<sup>\*,1</sup> and KC Johnson<sup>2</sup><sup>1</sup>Department of Epidemiology and Social Medicine, Albert Einstein College of Medicine, Bronx, NY 10461, USA; <sup>2</sup>Centre for Chronic Disease Prevention and Control, Population and Public Health Branch, Health Canada, Tunney's Pasture, P.L. # 0601C1, Ottawa, Ontario, Canada, K1A 0L2British Journal of Cancer (2003) 88, 1500. doi:10.1038/sj.bjc.6600933 www.bjcancer.com  
© 2003 Cancer Research UK

## Sir

The conclusion of the Collaborative Group on Hormonal Factors in Breast Cancer (Collaborative Group on Hormonal Factors and Breast Cancer, 2002) that cigarette smoking has little or no independent effect on the risk of developing breast cancer may not be warranted. Several recent studies of breast cancer risk suggest that a positive association with smoking may emerge 30–40 years after commencement (Terry and Rohan, 2002), particularly among chronic heavy smokers. There is also evidence to suggest that smoking prior to a first full-term pregnancy may be an important exposure (Russo *et al*, 1982; Band *et al*, 2002; Terry and Rohan,

2002). In addition, an increasing number of studies suggest that a positive association between smoking and breast cancer risk may be stronger among (or limited to) women with certain genotypes (Terry and Rohan, 2002). To what extent the Collaborative Group's examination of smoking as 'ever vs never' or 'current vs never' could account for such possible time-dependent or genetic associations is unclear. The possibility of an association between cigarette smoking and breast cancer risk among certain groups of women should not be ruled out based solely on the Collaborative Group's analyses.

## REFERENCES

- Band PR, Le ND, Fang R, Deschamps M (2002) Carcinogenic and endocrine disrupting effects of cigarette smoke and risk of breast cancer. *Lancet* 360(9339): 1044–1049
- Collaborative Group on Hormonal Factors Breast Cancer (2002) Alcohol, tobacco and breast cancer – collaborative reanalysis of individual data from 53 epidemiological studies, including 58 515 women with breast cancer and 95 067 women without the disease. *Br J Cancer* 87: 1234–1245
- Russo J, Tay LK, Russo IH (1982) Differentiation of the mammary gland and susceptibility to carcinogenesis. *Breast Cancer Res Treat* 2(1): 5–73
- Terry PD, Rohan TE (2002) Cigarette smoking and the risk of breast cancer in women: a review of the literature. *Cancer Epidemiol, Biomarkers Prev* 11: 953–971

\*Correspondence: Dr P Terry; E-mail: Terry2@niehs.nih.gov