



Journal of Epidemiology and Global Health

ISSN (Online): 2210-6014

ISSN (Print): 2210-6006

Journal Home Page: <https://www.atlantis-press.com/journals/jegh>

Fall-related injuries in a low-income setting: Results from a pilot injury surveillance system in Rawalpindi, Pakistan

Junaid A. Bhatti, Umar Farooq, Mudassir Majeed, Jahangir S. Khan, Junaid A. Razzak, Muhammad M. Khan

To cite this article: Junaid A. Bhatti, Umar Farooq, Mudassir Majeed, Jahangir S. Khan, Junaid A. Razzak, Muhammad M. Khan (2015) Fall-related injuries in a low-income setting: Results from a pilot injury surveillance system in Rawalpindi, Pakistan, Journal of Epidemiology and Global Health 5:3, 283–290, DOI:

<https://doi.org/10.1016/j.jegh.2015.01.003>

To link to this article: <https://doi.org/10.1016/j.jegh.2015.01.003>

Published online: 23 April 2019



Fall-related injuries in a low-income setting: Results from a pilot injury surveillance system in Rawalpindi, Pakistan

Junaid A. Bhatti ^{a,b,c,1,*}, Umar Farooq ^{d,1}, Mudassir Majeed ^d,
Jahangir S. Khan ^d, Junaid A. Razzak ^e, Muhammad M. Khan ^d

^a Sunnybrook Health Sciences Centre Research Institute, Toronto, Canada

^b Department of Surgery, University of Toronto, Toronto, Canada

^c Institute for Clinical Evaluative Sciences, Toronto, Canada

^d Surgical Unit I, Department of Surgery, Holy Family Hospital, Rawalpindi, Pakistan

^e Department of Emergency Medicine, The Aga Khan University, Karachi, Pakistan

Received 17 July 2014; received in revised form 7 January 2015; accepted 28 January 2015

Available online 26 February 2015

KEYWORDS

Falls;
Pakistan;
Surveillance

Abstract This study assessed the characteristics and emergency care outcomes of fall-related injuries in Pakistan. This study included all fall-related injury cases presenting to emergency departments (EDs) of the three teaching hospitals in Rawalpindi city from July 2007 to June 2008. Out of 62,530 injury cases, 43.4% ($N = 27,109$) were due to falls. Children (0–15 years) accounted for about two out of five of all fall-related injuries. Compared with women aged 16–45 years, more men of the same age group presented with fall-related injuries (50% vs. 42%); however, compared with men aged 45 years or more, about twice as many women of the same age group presented with fall-related injuries (16% vs. 9%, $P < 0.001$). For each reported death due to falls ($n = 57$), 43 more were admitted ($n = 2443$, 9%), and another 423 were discharged from the EDs ($n = 24,142$, 91%). Factors associated with death or inpatient admission were: aged 0–15 years (adjusted odds ratio [aOR] = 1.35), aged 45 years or more (aOR = 1.94), male gender (aOR = 1.15), falls occurring at home (aOR = 3.38), in markets (aOR = 1.43), on work sites (aOR = 4.80), and during playing activities (aOR = 1.68). This ED-based surveillance

Abbreviations: ED; emergency department; LMICs; low- and middle-income countries; SD; standard deviation; WHO; World Health Organization

* Corresponding author at: Sunnybrook Health Sciences Centre Research Institute, 2075 Bayview Avenue, G1-06, Toronto M4N 3M5, Canada. Tel.: +1 416 480 4055; fax: +1 416 480 6049.

E-mail addresses: junaid.bhatti@yahoo.com, junaid.bhatti@ices.on.ca (J.A. Bhatti).

¹ Equally contributed.

<http://dx.doi.org/10.1016/j.jegh.2015.01.003>

2210-6006/© 2015 Ministry of Health, Saudi Arabia. Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

study indicated that fall prevention interventions in Pakistan should target children, older adult women, homes, and work sites.

© 2015 Ministry of Health, Saudi Arabia. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Injuries account for about 12% of the disease burden worldwide and have a major impact on the health system [1]. Out of 5 million deaths that were attributed to injuries in 2000, falls accounted for about 424,000 deaths, making it the second leading cause of unintentional injury deaths worldwide after road traffic injuries [1–3]. The Global Burden of Disease report has estimated that fall-related deaths have increased by 55% from 348,000 deaths annually in 1990 to about 540,000 deaths in 2010 [4]. More worrying is the fact that this burden was disproportionately distributed as more than 82% of fall-related mortality and 92% of disability adjusted life year (DALY) loss occurred in low- and middle-income countries (LMICs) [5].

Research conducted in developed countries showed that fall-related injuries were frequent in children, older adults, and occupational settings [6]. The economic costs of such injuries are overwhelming because of the high number of childhood injury visits, hospital care of the older adult population, and productivity losses [2,7–10]. The most immediate consequence of falls, however, is engaging available healthcare resources in dealing with injury problems that are amenable to prevention [11].

The situation in LMICs could be much worse where the burden of fall-related injuries might affect performance of the healthcare system more rapidly than high-income settings because of limited resources [12]. To date, few surveillance studies have been carried out to assess the impact of fall-related injuries in LMIC settings, in particular on emergency care systems that could be overwhelmed by the care of such injuries [13,14]. This lack of data on local injury problems could hamper appropriate resource allocation to prevent and control fall-related injuries in LMICs [15].

Pakistan is a low-income country with a population of over 160 million inhabitants [16]. One previous health survey conducted at the national level indicated that falls accounted for almost half of the injury incidence in the country and were more frequent in urban settings than rural settings [17]. Another national survey indicated that fall-related injuries were the most common mechanism of

injuries after injuries that occur on roads [18,19]. Both surveys, however, lacked clarity about circumstances of fall-injuries because both classified all injuries occurring on roads as road traffic injuries. More worryingly, none of the previous studies had evaluated the impact of this burden on the healthcare system, in particular, the emergency care outcomes that could be used for monitoring healthcare burden of these injuries, as well as informing about preventive interventions [7,12,20]. The aim of this study is to assess the characteristics and emergency care outcomes of fall-related injuries in an urban setting in Pakistan.

2. Methods

2.1. Study setting and population

This prospective observational study was conducted in Rawalpindi city which had a population of 1.6 million in 2006 and an urbanization rate of around 4% [21,22]. An injury surveillance system was established at the emergency departments (EDs) of three teaching hospitals, namely: Holy Family Hospital, Benazir Bhutto Shaheed Hospital, and District Headquarter Hospital. A fall in the injury cases was defined as “a sudden and unintentional change in position resulting in an individual landing at a lower level such as on an object, the floor, or the ground, with or without injury” [23]. This definition was concordant with the Prevention of Falls Network Europe Consensus [24] and the International Classification of Disease W00–W19 [25]. All cases presenting to EDs over a period from July 1, 2007 to June 30, 2008 with a fall-related injury were included in the study. The questionnaire used in this study distinguished fall-related injuries from road traffic injuries, as the variable used for defining both mechanisms ensured that both events were recorded as mutually exclusive [26,27]. Ethical approval of the study protocol was obtained from Rawalpindi Medical College research ethics council before beginning the study.

2.2. Data collection

Data were recorded on minimal dataset questionnaires for injury surveillance by the World Health

Organization (WHO) [26]. This one-page questionnaire was translated into Urdu and was back-translated into English to ensure consistency [28]. The questionnaire was filled through face-to-face interviews after the initial management of these patients and after obtaining informed consent. In each hospital, data collection was coordinated by the resident in surgery. Four data collectors from the attending hospital staff were nominated in each ED which ensured data collection 24 h a day. Questionnaires were then transferred to a central location every third day where they were verified by the principal investigators. These were then coded on an Excel® spreadsheet by two data entry operators; 10% of the entries were checked by principal investigators.

2.3. Measurements

The variables included were age, gender, place, activity, nature, severity and outcome of the fall-related injury cases. Severity of injury was defined as mild, moderate, or severe according to the surveillance guidelines [26]. Outcome was defined as the discharge from the ED, admission to an inpatient surgical ward, or death at presentation or during the stay in the ED. The maximum period of stay in the ED was 24 h; further stay required inpatient admission. Intent of injury was defined as unintentional or intentional; the intent was further described as self-harm or assaults (interpersonal violence) [29].

2.4. Analysis

The age and gender distributions of fall-related injury cases were assessed according to three age categories: 0–15 years, 16–45 years and 45 years or more [26]. Age distribution of fall-related injuries in males and females was compared using chi-square test. A significant *P* value meant that for a given injury characteristic, e.g., site, activity, intent, severity, nature, or outcome, the age distribution was different between male and female cases. For assessing factors associated with adverse ED outcomes, the category of deaths was combined with inpatient admission because of fewer numbers. For this purpose, a logistic regression analysis with backward selection strategy was computed where all variables associated with ED outcomes (death or admission) with *P* value ≤ 0.20 were included in the analysis [30]. Statistical Package for Social Sciences Version 20.0 was used for these analyses.

3. Results

During the 12-month period, 62,530 people presented with injuries to EDs of the three teaching hospitals. Of these, 43.4% ($N = 27,109$) were fall-related injuries. The majority of these injuries were among males (75.0%; $N = 20,280$). The mean age of males was 21.2 years ($SD = 16.8$) and females was 24.7 years ($SD = 19.9$).

In males, fall-related injuries were sustained mostly at homes (45.8%) and on road sites (30.9%), whereas in females four out of five (80.1%) fall-related injuries occurred at home (Table 1). About half of injuries in females (49.1%) occurred during work activity compared with about a third in males (33.6%). A third of fall-related injuries in males (35.4%) and females (35.6%) occurred during playing. Almost all injuries were unintentional (99.8%) in males and females. About a third of injuries were of moderate severity in males (35.4%) and females (36.2%). About a third of injuries resulted in open wounds in males (37.8%) and females (31.5%); one in five injuries resulted in either a concussion or a fracture in males (12.1% and 10.0%, respectively) and females (11.9% and 13.4%, respectively).

Two out of five of the fall-injury cases were children (0–15 years); 42.1% were males and 41.9% were females (Fig. 1). Row proportions may not add to 100% because of missing values in $< 0.5\%$. Compared with females aged 16–45 years, more males of the same age presented to the ED (49.5% vs. 42.2%); however, females aged 45 years or more were twice as likely to present with fall injuries than males of the same age (15.9% vs. 8.9% $P < 0.001$). Compared with female children, nearly two-thirds (64.3%) of fall-related injuries in male children (0–15 years) occurred at home. Male and female children accounted for almost all playing-related falls (90.4% in males and 93.8% in females). About three quarters of work site- and activity-related falls occurred in males and females aged 16–45 years. The severity of injuries, such as moderate or severe injury, as well as nature of injury, such as fractures, sprains, bruises and organ system injury, was significantly higher ($P < 0.001$) in women aged 45 years or more as compared with men of the same ages.

Fall-related injury cases accounted for 57 deaths during the study period. Out of all fall-related injury cases, approximately one tenth required inpatient admission: 8.9% in males and 9.4% in

Table 1 Characteristics of fall-related injuries presenting to the three teaching hospitals in Rawalpindi (Jul 2007–Jun 2008).

| Age (in years) <i>N</i> | Males | | | | | Females | | | | | <i>P</i> * |
|-------------------------|----------|--------|-----------|------------|----------|----------|--------|-----------|------------|----------|------------|
| | <i>N</i> | (%) | 0–15 % | 16–45 % | >45 % | <i>N</i> | (%) | 0–15 % | 16–45 % | >45 % | |
| Site | | | | | | | | | | | |
| – Home | 9279 | (45.8) | 64.3 | 27.4 | 8.3 | 5471 | (80.1) | 43.5 | 40.6 | 15.9 | <0.001 |
| – School | 724 | (3.6) | 82.4 | 15.3 | 1.2 | 243 | (3.6) | 88.3 | 10.6 | 0.5 | 0.09 |
| – Roads | 6257 | (30.9) | 22.1 | 69.9 | 8.0 | 734 | (10.8) | 26.9 | 54.4 | 18.9 | <0.001 |
| – Market | 2026 | (10.0) | 19.8 | 69.5 | 10.8 | 203 | (3.0) | 15.5 | 58.0 | 26.4 | <0.001 |
| – Work site | 1616 | (8.0) | 6.9 | 83.3 | 9.8 | 57 | (0.8) | 5.0 | 75.6 | 15.4 | 0.26 |
| – Other | 378 | (1.9) | 18.0 | 69.1 | 13.5 | 121 | (1.8) | 28.4 | 54.6 | 17.2 | 0.01 |
| Activity | | | | | | | | | | | |
| – Work | 6819 | (33.6) | 12.5 | 73.8 | 13.7 | 3352 | (49.1) | 10.0 | 66.4 | 23.7 | <0.001 |
| – Playing | 7188 | (35.4) | 90.4 | 8.1 | 1.6 | 2429 | (35.6) | 93.8 | 4.7 | 1.4 | <0.001 |
| – Traveling | 5083 | (25.1) | 15.8 | 75.5 | 8.9 | 609 | (8.9) | 17.4 | 60.4 | 22.1 | <0.001 |
| – Other | 1190 | (5.9) | 32.2 | 50.7 | 17.0 | 439 | (6.4) | 32.6 | 37.3 | 30.2 | <0.001 |
| Intent | | | | | | | | | | | |
| – Unintentional | 20,161 | 99.8 | 42.2 | 49.7 | 8.4 | 6794 | 99.8 | 42.0 | 42.2 | 16.1 | <0.001 |
| – Deliberate self-harm | 12 | 0.1 | 71.0 | 83.8 | 14.2 | 4 | 0.1 | 50.0 | 50.0 | 0 | 0.88 |
| – Assault | 19 | 0.1 | 44.9 | 105.8 | 9.0 | 5 | 0.1 | 40.0 | 40.0 | 20.0 | 0.58 |
| Injury severity | | | | | | | | | | | |
| – None | 923 | (4.5) | 44.3 | 45.7 | 10.0 | 399 | (5.8) | 30.1 | 49.7 | 20.0 | <0.001 |
| – Mild | 12,016 | (59.3) | 38.2 | 53.8 | 8.1 | 3907 | (57.2) | 41.9 | 44.1 | 14.0 | <0.001 |
| – Moderate | 7171 | (35.4) | 47.9 | 43.5 | 8.6 | 2473 | (36.2) | 43.7 | 37.9 | 18.4 | <0.001 |
| – Severe | 170 | (0.8) | 55.2 | 29.6 | 15.0 | 50 | (0.7) | 40.0 | 28.7 | 30.7 | 0.04 |
| Nature of injury | | | | | | | | | | | |
| – Fracture | 2017 | (10.0) | 48.2 | 39.4 | 12.6 | 915 | (13.4) | 30.3 | 42.7 | 27.2 | <0.001 |
| – Sprain/strain | 908 | (4.5) | 29.1 | 59.8 | 10.9 | 604 | (8.8) | 18.0 | 61.4 | 20.3 | <0.001 |
| – Cut/bite/open wound | 7670 | (37.8) | 39.2 | 53.1 | 7.8 | 2153 | (31.5) | 48.5 | 39.9 | 11.5 | <0.001 |
| – Bruise | 1142 | (5.6) | 36.6 | 54.6 | 9.3 | 416 | (6.1) | 35.1 | 48.4 | 16.8 | <0.001 |
| – Concussion | 2460 | (12.1) | 63.8 | 29.4 | 8.5 | 815 | (11.9) | 64.2 | 25.0 | 10.7 | 0.02 |
| – Organ system injury | 3602 | (17.8) | 37.4 | 55.0 | 4.7 | 1196 | (17.5) | 39.7 | 43.3 | 16.9 | <0.001 |
| – Other | 2481 | (12.2) | 38.8 | 53.1 | 11.1 | 730 | (10.7) | 39.2 | 44.9 | 15.9 | <0.001 |
| Outcome | | | | | | | | | | | |
| – Discharged | 18,081 | (89.2) | 40.4 | 51.5 | 8.2 | 6043 | (88.5) | 40.8 | 43.8 | 15.3 | <0.001 |
| – Admitted | 1799 | (8.9) | 59.7 | 30.2 | 10.5 | 644 | (9.4) | 52.8 | 27.2 | 20.9 | <0.001 |
| – Died | 40 | (0.2) | 42.6 | 50.3 | 17.0 | 17 | (0.3) | 50.5 | 33.8 | 12.9 | 0.60 |
| – Other | 360 | (1.8) | 37.9 | 53.0 | 9.0 | 125 | (1.8) | 41.2 | 34.5 | 25.4 | <0.001 |

* Difference in age distribution between male and female for a given factor.

females. The likelihood of inpatient admission or death was significantly higher for those aged 0–15 years or 45 years or more than those aged 16–45 years (Table 2). Similarly, the likelihood of inpatient admission or death due to falls was higher in males than females, for falls occurring at homes, in markets, on work sites, or other sites than those on roads. Similarly, the likelihood of inpatient admission or death was higher for falls occurring during playing and other activities than those during traveling-related activities.

4. Discussion

This study presented for the first time the full age and gender spectrum of fall-related injuries in a large sample of patients presenting to EDs in Pakistan. The study showed that falls resulted in a significant number of ED visits, even more than road traffic injuries [27], in the three teaching hospitals over the study period. Most of those affected from fall-related injuries were relatively young, and children accounted for a significant proportion




Fig. 1 Age and gender-wise distributions of fall-related injuries in Rawalpindi city.

of such injuries. Children and females were mostly injured at home, whereas males aged 16–45 years were injured on road sites. Most of the children were injured while playing, whereas young and

older adults were injured during work activities. Fractures and concussions were commonly observed in children and older adults after open wound presentations. For each reported death due to falls, 43 more were admitted, and another 423 were discharged from EDs. Factors associated with inpatient admission or death were being children or men aged 45 years or more, injuries occurring at homes, roads, markets or work sites, and during playing.

This study had several limitations. First, this study was conducted in tertiary care hospitals, thus a significant number of mild to moderate injury patients consulting private health care services could not be accounted here [16]. Another study where an alternate source for reporting specific injuries was possible using the same data, e.g., assaults, showed that EDs reporting in the three hospitals accounted for about 33–43% of all assault-related injuries in Rawalpindi [31]. Secondly, outcome was recorded in the ED and

Table 2 Factors associated with emergency department (ED) outcomes of fall-related injuries in Rawalpindi city (Jul 07–Jun 08).

| | Discharged from ED | | Admitted | | Died | | <i>P</i> * | Death or admission vs. discharge from ED | |
|-----------------------------------|--------------------|-------|----------|------|----------|------|------------|--|-----------|
| | <i>N</i> | (%) | <i>N</i> | (%) | <i>N</i> | (%) | | aOR [†] | 95% CI |
| Age (in years) | | | | | | | <0.001 | | |
| – 0–15 | 9769 | 40.5 | 1404 | 57.5 | 26 | 45.6 | | 1.35 | 1.18–1.56 |
| – 16–45 | 11,953 | 49.6 | 715 | 29.3 | 22 | 38.6 | | 1 | |
| – >45 | 2402 | 9.9 | 324 | 13.3 | 9 | 15.8 | | 1.94 | 1.68–2.30 |
| Gender | | | | | | | 0.20 | | |
| – Female | 6043 | 25.1 | 644 | 26.4 | 17 | 29.8 | | 1 | |
| – Male | 18,081 | 74.9 | 1799 | 73.6 | 40 | 70.2 | | 1.15 | 1.04–1.27 |
| Place of injury | | | | | | | <0.001 | | |
| – Roads | 6614 | 52.31 | 212 | 8.7 | 8 | 14.0 | | 1 | |
| – Home | 12,619 | 52.3 | 1846 | 75.6 | 35 | 61.4 | | 3.38 | 2.78–4.10 |
| – School | 932 | 3.8 | 32 | 1.3 | 0 | 0.0 | | 0.64 | 0.43–0.96 |
| – Market | 2088 | 8.7 | 99 | 4.0 | 5 | 8.8 | | 1.43 | 1.11–1.84 |
| – Work site | 1471 | 6.1 | 186 | 7.6 | 5 | 8.8 | | 4.80 | 3.75–6.14 |
| – Other | 409 | 1.7 | 68 | 2.8 | 4 | 7.0 | | 4.82 | 3.57–6.52 |
| Activity at time of injury | | | | | | | <0.001 | | |
| – Traveling | 5381 | 22.3 | 177 | 7.2 | 7 | 12.3 | | 1 | |
| – Work | 9237 | 38.3 | 771 | 31.6 | 23 | 40.4 | | 0.93 | 0.74–1.15 |
| – Playing | 8154 | 33.8 | 1297 | 53.1 | 17 | 29.8 | | 1.68 | 1.34–2.12 |
| – Other | 1352 | 5.6 | 198 | 8.1 | 10 | 17.5 | | 1.91 | 1.49–2.44 |
| Intent | | | | | | | <0.001 | | |
| – Unintentional | 23,997 | 99.5 | 2430 | 99.5 | 55 | 96.5 | | 1 | |
| – Intentional | 30 | 0.1 | 7 | 0.3 | 2 | 3.5 | | 0.25 | 0.11–0.55 |
| – Unknown | 97 | 0.4 | 6 | 0.2 | 0 | 0 | | 0.15 | 0.05–0.47 |

aOR – adjusted odds ratio.

* Association of factors with deaths of admissions vs. discharged from ED.

† Hosmer and Lemeshow test *P* = 0.15.

patients were not followed to their discharge. Thus, these figures might underestimate the deaths due to fall-related injuries. Nonetheless, this study showed that ED surveillance could be a useful tool to identify burden and research priorities for specific types of injuries [26]. A more comprehensive trauma registry in low-income settings appeared to be essential for formalizing policies to reduce injury burden [13,15].

The study results were consistent with the previous work in Pakistan and other LMICs that falls were one of the leading mechanisms of injuries [17,32–34]. Since no recent census had been carried out in this city, incidence and age standardized rates could not be calculated in this study. Nevertheless, several demographic factors associated with adverse ED outcomes were identified in this study. Notably, findings showed a higher involvement of younger age groups in fall-related injuries than older adults which could be attributed to the demographic trends as the median age in Pakistan is about 20 years [16]. Similarly, this study showed that males accounted for about four out of five fall-related injuries, a trend also noted in other regional studies [14]. There could be several possible explanations for these findings. For instance, the cultural role of males as breadwinners could increase their likelihood of being involved in risk-taking activities, e.g., working at heights without proper gear or involved in repair work at homes without precautionary measures [19]. Previous studies in Pakistan noted that male vendors were at a significant risk for road traffic injuries in Pakistan [19,35]. In addition, males might enjoy a higher access to emergency services than females, i.e., in case of injury they would likely be transported to an ED than their female counterparts [19,36]. Gender differences have indeed been reported by a study on cultural determinants of ambulance use in Karachi, Pakistan [37].

A worrisome finding from the analysis of age and gender analyses was that a higher proportion, more than two times, of fall injuries occurred in older women, i.e., age >45 years than men of the same age group. The more severe consequences of fall injuries – notably fractures, concussions, bruises and organ system injuries – indicated that older Pakistani women were more vulnerable to such outcomes than men of the same age groups. Previous work had shown that older women were twice as likely to sustain fractures as males of the same age [10]. Poor nutritional status and a decrease in bone mass in women, especially in the five years following menopause, might explain

this severe injury outcome compared with men [38]. Screening of women for predisposing factors and awareness campaigns focusing on such problems might decrease the incidence of fall-related injuries in older Pakistani women [39].

Another important study finding was that about two thirds of fall-related fatalities occurred at homes. The proportion of other adverse outcomes, such as hospital admission, was also two times higher than the one reported in New Zealand where about one third of fall-related injuries occurred at homes [40]. These findings suggested that hazards at homes might not be perceived adequately by the residents. Other studies on home health hazards in Pakistani children showed that most hazards for injuries including falls go unappreciated by the parents and could potentially increase chances for falls [41,42]. This study showed that the older population might also be at a higher risk for home hazards for falls, and could require special interventions, including awareness to prevent such injuries [10,43].

This study showed that most of fall-related injuries in children occurred while playing either at homes or schools. In similar settings, this could be explained by lack of safety measures, as in unattended children and children flying kite at roof top [20,44]. Factors predisposing to child injuries had rarely been investigated in Pakistan and, currently, there are no ongoing injury prevention programs for children [17,45–47]. The high incidence of fall injuries while playing indicated the need for more supervision during playing and identification of specific risk factors for these injuries in Pakistani children [46].

A significant proportion of falls in the adult age group resulted during work-related activities in this study. Undoubtedly, unsafe work practices are adopted due to financial constraints and related factors in Pakistan [35]. Workers' safety remains an important problem and occupational injuries, particularly the incidence of fall-related injuries, had not been investigated in Pakistan [22]. The poor outcome of such injuries shown in this study indicated that a surveillance study of occupational injuries might be required in Pakistan to guide policy-making [9].

5. Conclusion

Fall-related injuries resulted in a relatively high burden on the healthcare system in this urban setting. Further analyses revealed that future research and prevention efforts should focus on fall-related injuries in childhood, at the home,

and during work activities [2,8,9]. Moreover, epidemiological methods could illustrate underlying mechanisms of such injuries and help in the development of appropriate preventive measures [15]. Finally, ED surveillance of fall-related injuries in childhood and due to work could provide useful information to prevent and control the increasing burden of fall-related injuries in LMICs like Pakistan [26].

Contribution of Authors

UF, MM, and JSK collected the data. JAB and JAR performed analysis and interpretation of results. JAB took the lead in manuscript writing. All authors participated in the critical revision of this manuscript. The final manuscript was seen and approved by all the authors.

Funding

This study was funded by the Government of Punjab, Pakistan. Funding organization had no input in the study design, conception, and the conduct of the study.

Competing interest

No competing interests were identified for any of the authors.

Acknowledgements

We are grateful to the emergency staff, director emergency, and medical superintendent of the teaching hospitals for their cooperation during the study. We highly appreciate the Government of Punjab for taking the initiative to collect data on injuries.

References

- [1] Peden M, McGee K, Sharma G. The injury chart book: a graphical overview of the global burden of injuries. Geneva: World Health Organization; 2002.
- [2] Peden M, Oyegbite K, Ozanne-Smith J, Hyder AA, Branche C, Rahman AKMF, et al., editors. World report on child injury prevention. Geneva: World Health Organization; 2008.
- [3] Murray CJL, Lopez AD, Mathers CD, Stein C. The Global Burden of Disease 2000 project: aims, methods and data sources. Geneva: World Health Organization; 2001.
- [4] Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380(9859):2095–128.
- [5] Mathers C, Boerma T, Ma Fat D. The Global Burden of Disease 2004 update. Geneva: World Health Organization; 2008.
- [6] Centers for Disease Control and Prevention. Injury research agenda: preventing injuries at home and in the community. Atlanta (GA): National Center for Injury Prevention and Control; 2002.
- [7] Lallier M, Bouchard S, St-Vil D, Dupont J, Tucci M. Falls from heights among children: a retrospective review. *J Pediatr Surg* 1999;34(7):1060–3.
- [8] Norton R, Hyder AA, Bishai D, Peden M. Unintentional Injuries. In: Jamison DT, Feachem RG, Makgoba MW, Bos ER, Baingana FK, Hofman KJ, et al., editors. Disease and mortality in sub-saharan Africa. Washington (DC): World Bank; 2006.
- [9] Rosenstock L, Cullen M, Fingerhut M. Occupational health. In: Jamison DT, Feachem RG, Makgoba MW, Bos ER, Baingana FK, Hofman KJ, et al., editors. Disease and mortality in sub-saharan Africa. Washington (DC): World Bank; 2006.
- [10] Stevens JA, Corso PS, Finkelstein EA, Miller TR. The costs of fatal and non-fatal falls among older adults. *Inj Prev* 2006;12(5):290–5.
- [11] Sartini M, Cristina ML, Spagnolo AM, Cremonesi P, Costaguta C, Monacelli F, et al. The epidemiology of domestic injurious falls in a community dwelling elderly population: an outgrowing economic burden. *Eur J Public Health* 2010;20(5):604–6.
- [12] Quansah R. Essential trauma care in Ghana: adaptation and implementation on the political tough road. *World J Surg* 2006;30(6):934–9.
- [13] Juillard C, Etoundi Mballa GA, Bilounga Ndongo C, Stevens KA, Hyder AA. Patterns of injury and violence in Yaounde Cameroon: an analysis of hospital data. *World J Surg* 2011;35(1):1–8.
- [14] Bengalor Injury Surveillance Programme. Factsheet on falls. Bangalore: National Institute of Mental Health and NeuroSciences; 2008.
- [15] Jamison DT, Mosley WH, Measham AB, Bobadilla JL, editors. Disease control priorities in developing countries. Washington (DC): Oxford University Press; 1993.
- [16] Government of Pakistan. Pakistan Economic Survey 2008–09. Islamabad: Economic Advisor's Wing, Finance Division, Government of Pakistan; 2009.
- [17] Fatmi Z, Hadden WC, Razzak JA, Qureshi HI, Hyder AA, Pappas G. Incidence, patterns and severity of reported unintentional injuries in Pakistan for persons five years and older: results of the National Health Survey of Pakistan 1990–94. *BMC Public Health* 2007;7:152.
- [18] Bachani AM, Ghaffar A, Hyder AA. Burden of fall injuries in Pakistan – analysis of the National Injury Survey of Pakistan. *East Mediterr Health J* 2011;17(5):375–81.
- [19] Ghaffar A, Hyder AA, Masud TI. The burden of road traffic injuries in developing countries: the 1st National Injury Survey of Pakistan. *Public Health* 2004;118(3):211–7.
- [20] Raja IA, Vohra AH, Ahmed M. Neurotrauma in Pakistan. *World J Surg* 2001;25(9):1230–7.
- [21] Government of Pakistan. Census report of Pakistan 1998. Islamabad: Population Census Organization, Statistics Division, Government of Pakistan; 2001.
- [22] Nishtar S, Mohamud KB, Razzak J, Ghaffar A, Ahmed A, et al. Injury prevention and control: National Action Plan for NCD prevention, control and health promotion in Pakistan. *JPMA* 2004;54(12 Suppl. 3):S57–68.
- [23] Public Health Agency of Canada. Report on seniors' falls in Canada. Ottawa: Minister of Public Works and Government Services Canada; 2005, 8.

- [24] Lamb SE, Jorstad-Stein EC, Hauer K, Becker C. Prevention of Falls Network Europe and Outcomes Consensus Group. Development of a common outcome data set for fall injury prevention trials: the prevention of falls network Europe consensus. *J Am Geriatr Soc* 2005;53(9):1618–22.
- [25] World Health Organization. International statistical classification of diseases and related health problems 10th revision: version for 2005. Geneva: World Health Organization; 2005.
- [26] Holder Y, Peden M, Krug E. Injury surveillance guidelines. Geneva: World Health Organization; 1999.
- [27] Farooq U, Bhatti JA, Siddiq M, Majeed M, Malik N, et al. Road traffic injuries in Rawalpindi city, Pakistan. *East Mediterr Health J* 2011;17(9):647–53.
- [28] Farooq U, Nasrullah M, Bhatti JA, Majeed M, Hanif M. Incidence of burns and factors associated with their hospitalisation in Rawalpindi, Pakistan. *Burns* 2011;37(3): 535–40.
- [29] Farooq U, Majeed M, Bhatti JA, Khan JS, Razzak JA, Khan MM. Differences in reporting of violence and deliberate self harm related injuries to health and police authorities, Rawalpindi, Pakistan. *PLoS One* 2010;5(2):e9373.
- [30] Lemeshow S, Hosmer DW, Klar J, Lwanga SK. Adequacy of sample size in health studies. West Sussex, England: John Wiley & Sons Ltd; 1990.
- [31] Farooq U, Majeed M, Bhatti JA, Khan JS, Razzak JA, Khan MM. Differences in reporting of violence and deliberate self harm related injuries to health and police authorities, Rawalpindi, Pakistan. *PLoS One* 2010;5(2):e9373.
- [32] Mock CN, Abantanga F, Cummings P, Koepsell TD. Incidence and outcome of injury in Ghana: a community-based survey. *Bull World Health Organiz* 1999;77(12):955–64.
- [33] Otieno T, Woodfield JC, Bird P, Hill AG. Trauma in rural Kenya. *Injury* 2004;35(12):1228–33.
- [34] Bangdiwala SI, Anzola-Perez E, Romer CC, Schmidt B, Valdez-Lazo F, et al. The incidence of injuries in young people: I. Methodology and results of a collaborative study in Brazil, Chile, Cuba and Venezuela. *Int J Epidemiol* 1990;19(1):115–24.
- [35] Ghaffar A, Hyder AA, Mastoor MI, Shaikh I. Injuries in Pakistan: directions for future health policy. *Health Policy Plann* 1999;14(1):11–7.
- [36] Dandona R, Kumar GA, Ivers R, Joshi R, Neal B, Dandona L. Characteristics of non-fatal fall injuries in rural India. *Inj Prev* 2010;16(3). <http://dx.doi.org/10.1136/ip.2009.023663>.
- [37] Razzak JA, Cone DC, Rehmani R. Emergency medical services and cultural determinants of an emergency in Karachi, Pakistan. *Prehospital Emergency Care* 2001;5(3): 312–6.
- [38] Prince RL. Calcium and vitamin D – for whom and when. *Menopause Int* 2007;13(1):35–7.
- [39] Morales-Torres J. Strategies for the prevention and control of osteoporosis in developing countries. *Clin Rheumatol* 2007;26(2):139–43.
- [40] Kool B, Ameratunga S, Robinson E, Jackson R. Hospitalisations and deaths due to unintentional falls at home among working-aged New Zealanders. *Injury* 2007;38(5):570–5.
- [41] Hyder AA, Chandran A, Khan UR, Zia N, Huang CM, et al. Childhood unintentional injuries: need for a community-based home injury risk assessments in Pakistan. *Int J Pediatr* 2012;2012:203204.
- [42] Khan UR, Chandran A, Zia N, Huang CM, Stewart De Ramirez S, Feroze A, et al. Home injury risks to young children in Karachi, Pakistan: a pilot study. *Arch Dis Child* 2013;98(11): 881–6.
- [43] Gillespie L. Preventing falls in elderly people. *BMJ* 2004; 328(7441):653–4.
- [44] Abbas I, Ali K, Mirza YB. Spectrum of mandibular fractures at a tertiary care dental hospital in Lahore. *J Ayub Med Coll Abbottabad* 2003;15(2):12–4.
- [45] Fatmi Z, Kazi A, Hadden WC, Bhutta ZA, Razzak JA, Pappas G. Incidence and pattern of unintentional injuries and resulting disability among children under 5 years of age: results of the National Health Survey of Pakistan. *Paediatr Perinat Epidemiol* 2009;23(3):229–38.
- [46] Hyder AA, Sugerman D, Ameratunga S, Callaghan JA. Falls among children in the developing world: a gap in child health burden estimations? *Acta Paediatr* 2007;96(10): 1394–8.
- [47] Hyder AA, Sugerman DE, Puvanachandra P, Razzak J, El-Sayed H, Isaza A, et al. Global childhood unintentional injury surveillance in four cities in developing countries: a pilot study. *Bull World Health Organiz* 2009;87(5):345–52.

Available online at www.sciencedirect.com

ScienceDirect