

RESEARCH

Open Access



# Enhanced bioethanol production from wheat straw hemicellulose by mutant strains of pentose fermenting organisms *Pichia stipitis* and *Candida shehatae*

Sravanthi Koti, Sai Prashanthi Govumoni, Jahnavi Gentela and L. Venkateswar Rao\*

## Abstract

The main aim of the present study was to mutate yeast strains, *Pichia stipitis* NCIM 3498 and *Candida shehatae* NCIM 3501 and assess the mutant's ability to utilize, ferment wheat straw hemicellulose with enhanced ethanol yield. The organisms were subjected to random mutagenesis using physical (ultraviolet radiation) and chemical (ethidium bromide) mutagens. The mutant and wild strains were used to ferment the hemicellulosic hydrolysates of wheat straw obtained by 2 % dilute sulphuric acid and enzymatic hydrolysis by crude xylanase separately. Among all the mutant strains, PSUV9 and CSEB7 showed enhanced ethanol production ( $12.15 \pm 0.57$ ,  $9.55 \pm 0.47$  g/L and yield  $0.450 \pm 0.009$ ,  $0.440 \pm 0.001$  g/g) as compared to the wild strains ( $8.28 \pm 0.54$ ,  $7.92 \pm 0.89$  g/L and yield  $0.380 \pm 0.006$  and  $0.370 \pm 0.002$  g/g) in both the hydrolysates. The mutant strains were also checked for their consistency in ethanol production and found stable for 19 cycles in hemicellulosic hydrolysates of wheat straw. A novel element in the present study was introduction of chemical mutagenesis in wild type as well as UV induced mutants. This combination of treatments i.e., UV followed by chemical mutagenesis was practically successful.

**Keywords:** Wheat straw, Hemicellulose, Mutant, Pentose, Yeast, Fermentation

## Background

Fossil fuel reserves are limited, and their current oxidation rate is a major global environmental concern with complex and severe climatic impacts (Stephenson et al. 2011). The increase in agro-industrial activity has led to the widespread accumulation of large quantities of lignocellulosic residues from wood, forestry, herbaceous, agricultural, solid, and various industrial wastes (José et al. 2010). These residues are collectively termed “biomass” and can be converted into ethanol fuel. Although bioethanol production has improved greatly by new technologies, further investigation is required to overcome the many remaining challenges (Yan and Shuzo 2006).

The production of ethanol from lignocellulosic biomass involves three major processes: pretreatment, hydrolysis,

and fermentation. Pretreatment is required to alter the size, structure, and chemical composition of the biomass to facilitate rapid and efficient hydrolysis (Chang and Holtzapfle 2000). Recent advances in pretreatment technology have the potential to improve the efficiency and reduce the cost of ethanol production (Mosier et al. 2003). Dilute acid pretreatment has become a state-of-the-art technology for the conversion of hemicellulose from any lignocellulosic biomass source into sugars. The fermentable sugars obtained via hydrolysis could subsequently be fermented into ethanol by ethanol-producing microorganisms (Lee et al. 1999).

Hemicellulose, a branched polymer composed of pentose and hexose sugars, can be hydrolyzed by hemicellulases or acids to release its monomeric sugars. Xylose and arabinose generally constitute a significant fraction of lignocellulosic biomass; therefore, their utilization is essential for a feasible bioethanol production process (Aristidou and Penttila 2000; Bothast et al. 2002; Koti

\*Correspondence: vrlinga@gmail.com  
Department of Microbiology, Osmania University, Hyderabad, Telangana state 500 007, India

et al. 2012; Sues et al. 2005). The microbial strain selected for the fermentation of pentose sugars has a large effect on ethanol yield (Anuj et al. 2011). Therefore, naturally xylose-fermenting yeasts such as *Candida shehatae* and *Pichia stipitis* have been widely studied because of their ability to ferment xylose into ethanol (Borbala et al. 2012). *Pichia stipitis* is considered a promising strain because it can ferment a wide range of sugars, including cellobiose (Nigam 2001). Furthermore, *Candida* species have been shown to ferment d-xylose to ethanol as the major product (Gong et al. 1983). The enhancement of microbial strains through mutation or gene cloning has gained attention in the commercial fermentation industry as a means to increase ethanol yields. Mutational enhancement of microorganisms is an old technique; however, use of this approach has resulted in improved ethanol yields at the laboratory level in several studies (Anuj et al. 2011).

In the present study, efforts were made to improve the pentose fermenting yeast strains by mutations using physical (UV irradiation) and chemical (ethidium bromide treatment) mutagens and selected mutant strains were assessed for their ability to produce enhanced yields of ethanol from wheat straw.

## Methods

### Substrate and yeast strains

The wheat straw used in the present study was obtained from Medak, Telangana state, India and the type used was *Triticum dicoccum*. The straw was dried at 60 °C in a hot air oven until the constant weight was obtained and processed in a laboratory pulverizer, sieved to attain a particle size between 1–3 mm. In order to avoid analytical interferences, the substrate was washed before the hydrolysis with tap water to make it free from dust and dried at 65 ± 0.5 °C for overnight. The cellulose, lignin and hemicellulosic fractions of pulverized wheat straw were determined according to ASTM (2007) method.

Yeast strains of *Pichia stipitis* NCIM 3498, *Candida shehatae* NCIM 3501 were obtained from National Chemical Laboratory, Pune, India and maintained on MGYPX agar (g/L: peptone, 10; yeast extract, 10; D-glucose, 20; xylose, 5; agar, 20).

### Mutagenesis of yeast strains

#### UV mutagenesis

UV mutagenesis was carried out according to the method of Winston and Ausubel (Winston and Ausubel 1990). Overnight grown cultures of *Pichia stipitis* and *Candida shehatae* (5 mL) were washed and re-suspended in 0.1 M phosphate buffer (pH 5.4) in order to achieve 10<sup>8</sup> cells per ml. The above cell suspension (2 mL) was placed in a sterile Petri dish and exposed to UV rays at a distance of

20 cm. At regular intervals (15, 30, 45, 60 min), the samples were collected and serially diluted to have 200–300 viable cells in each plate. Then the samples were plated on MGYPX agar medium and incubated at 28 °C for 48 h.

#### Chemical mutagenesis

The wild strains and UV induced mutants were grown for overnight in MGYPX medium and the cells after incubation were washed and suspended in 0.1 M phosphate buffer (pH 5.4). A stock of 0.1 mg/mL ethidium bromide was prepared and from this 1 mL of ethidium bromide was added to 9 mL of phosphate buffer containing yeast cells. After specific time intervals of 30, 60, 90, 120, 150 and 180 min of incubation, the cell suspensions were centrifuged at 3000 rpm for 5 min to remove the traces of mutagen. Cells were plated on MGYPX agar plates and incubated at 28 °C (Joanna and Ewelina 2003).

#### Enzyme assay

Xylanase assay was performed using 1 % (w/v) oat spelt xylan as substrate (Bailey et al. 1992). One unit (IU) of enzyme activity is defined as the amount of enzyme that produces 1 μmol of xylose in the reaction mixture per minute under the assay conditions used.

### Preparation of the wheat straw hemicellulosic hydrolysates

#### Pretreatment of wheat straw with NaOH

Wheat straw (250 g) was pretreated using 1.5 % (w/v) NaOH for 2 h at 100 °C with a liquid to solid ratio of 10:1 (Sai Prashanthi et al. 2013). The substrate was squeezed, washed and neutralized with tap water. Delignified filtrate obtained was analyzed for sugars and phenolic inhibitors (Miller 1959; Singleton et al. 1965).

#### Acid hydrolysis

Alkali pretreated wheat straw (50 g) was hydrolyzed at 121 °C with 2 % (v/v) sulfuric acid for 60 min, with an initial liquid to solid ratio of 10:1. The suspension was then squeezed to remove the unhydrolysed residue. The hydrolysate obtained was neutralized, detoxified and analyzed for sugars (Nigam 2001).

#### Enzyme production and enzymatic hydrolysis

The production media contained 50 g of corn cobs moistened with 50 mL mineral solution containing (g/L): KH<sub>2</sub>PO<sub>4</sub>, 28; (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 19.6; Urea, 4.2; MgSO<sub>4</sub>·7H<sub>2</sub>O, 4.2; CoCl<sub>2</sub>, 4.2; FeSO<sub>4</sub>·7H<sub>2</sub>O, 0.07; MnSO<sub>4</sub>·7H<sub>2</sub>O, 0.021; ZnSO<sub>4</sub>·7H<sub>2</sub>O, 0.019; CaCl<sub>2</sub>, 0.028; yeast extract, 7 and glucose, 15; pH 5.0 ± 0.2. The media were inoculated with 10 mL of inoculum having 10<sup>6</sup> spores/mL collected from 72 h grown culture of *Trichoderma asperellum* (Genbank accession number-KP965729). Inoculated production media were incubated under static conditions

at  $28 \pm 2$  °C and enzyme production was checked after every 24 h for 5 days. Enzyme was extracted with 500 mL of 0.05 M sodium acetate buffer on a rotary shaker at 150 rpm for 30 min. The content was filtered through muslin cloth and the filtrate was used as the enzyme source and utilized for enzymatic saccharification at a dosage of 0.25 mL (250 IU/mL, pH 4.8) per gram of alkali pretreated wheat straw (50 g) and incubated at  $50 \pm 0.5$  °C, 150 rpm for 48 h. After incubation, the hydrolysate was separated by filtration, supplemented with nutrients, sterilized and fermented to ethanol by wild and mutant strains.

#### Inoculum and fermentation media

The inoculum was developed by inoculating the wild type and mutant strains using media containing (g/L): xylose, 25; glucose, 5; yeast extract 5; malt extract, 5; peptone, 5; pH, 5.5) and incubated at 28 °C on a rotatory shaker at 200 rpm. The optical density (OD) of inoculum cultures was determined at the wavelength of 600 nm using a Systronics 117 UV-Vis spectrophotometer. Each fermentation media was inoculated with  $1.2 \times 10^8$  cells based on the conversion factor of 0.50 OD being equal to  $1 \times 10^7$  cells. The fermentation media used were 2 % acid hydrolysate and enzymatic hydrolysate consisted of 39.3 and 35.73 g/L of sugars respectively. Both the hydrolysates were supplemented with (g/L); yeast extract, 2;  $(\text{NH}_4)_2\text{SO}_4$ , 1;  $\text{K}_2\text{HPO}_4$ , 0.5; peptone, 1;  $\text{MgSO}_4$ , 0.5;  $\text{MnSO}_4$ , 0.5.

#### Ethanol fermentation

Ethanol fermentation was performed at 28 °C using wheat straw acid and enzymatic hydrolysates. As the fermentation was carried out in triplicates, average and standard deviations were calculated. Samples were collected at regular time intervals (12 h), centrifuged and supernatants obtained were examined for concentration of reducing sugars and ethanol.

#### Analytical methods

The amount of reducing sugars liberated was determined using the Dinitrosalicylic acid method with xylose/glucose as standard (Miller 1959). Ethanol concentration was analyzed by Gas chromatography (GC) (Shimadzu 2010, Japan) using ZB Wax column (30 mm  $\times$  0.25 mm) with a flame ionization detector (FID). The analysis was performed according to NREL (National Renewable Energy Laboratory) procedure LAP #001. The conditions used were: 150 °C (isothermal), program run time: 5.5 min, ethanol retention time: 2.3 min and the carrier gas: nitrogen (16 kPa), injector temperature: 175 °C, detector temperature: 250 °C, flow rate: 40 ml/min, split ratio: 1/50, velocity of  $\text{H}_2$  flow: 60 ml/min and sample quantity: 1  $\mu$ l.

The supernatant was filtered by 0.22  $\mu$ m cellulose acetate filters prior to GC analysis (Srilekha Yadav et al. 2011).

#### Statistical analysis

To assess whether there was any significant difference in ethanol production between the wild type and mutant strains of *Pichia stipitis* and *Candida shehatae* in acid and enzymatic hydrolysates of wheat straw, a paired t test was performed using Graph pad software (Graph pad software, inc., La Jolla, CA 92037 USA).

## Results and discussion

#### Compositional analysis of wheat straw

The cellulose, lignin and hemicellulosic fractions of pulverized wheat straw were determined according to ASTM (2007) method and reported in our previous study (Sai Prashanthi et al. 2013). The average percentages of acid soluble lignin, acid insoluble lignin, cellulose and hemicellulose were found to be  $16 \pm 1.15$ ,  $4.6 \pm 0.26$ ,  $32.60 \pm 0.37$  and  $24.7 \pm 0.2$  % respectively.

#### UV and EtBr mutagenesis

After separate UV and chemical mutagenesis, combination of UV and chemical mutagenesis was performed and selection of large colonies (42 colonies; from each method 7 colonies) was done on ethanol-containing medium. All these mutants were screened for maximum ethanol production in synthetic fermentation medium. The highest ethanol producing isolates were picked up from the UV followed by EtBr treatment (PSUV9, CSUV4) and EtBr mutagenesis (PSEB5, CSEB7) exhibiting 0.1 to 1.0 % survival. The mutants varied in cell size from parent and other mutants. Determination of ethanol produced by all the mutants revealed that only four mutants resulted in significant ethanol productivity in synthetic medium compared to their wild strains. In addition to the above four mutants, wild strains of *Pichia stipitis* NCIM 3498 and *Candida shehatae* NCIM 3501 were also investigated in this study as reference strains. The improvement of mutants in ethanol production (%) in fermentation medium II compared to the parent strains is summarized in Table 1.

#### Pretreatment of wheat straw with NaOH

Alkali pretreatment showed effective lignin solubilization of 70 % with minor cellulose and hemicellulose solubilization. Sugars and phenolics released during pretreatment were 0.83 and 17.28 g/L respectively.

#### Dilute acid and enzymatic hydrolysis of wheat straw

The acid hydrolysate contained 6.8 g/L of phenolics,  $0.334 \pm 0.014$  g/g reducing sugars ( $39.33 \pm 1.33$  g/L) with a practical conversion of  $41.79 \pm 1.62$  % of the total

**Table 1 Improvement of the production of ethanol in fermentation medium II by treatment with two mutagenic agents**

S. no.	Mutagenic treatment	Selected mutant strain	Ethanol production improved (%)
1	UV mutagenesis	PSUV9	46.73
		CSUV4	13.92
2	EtBr mutagenesis	PSEB5	11.39
		CSEB7	22.63

carbohydrates present in the straw which indicates that some part of the cellulose was also hydrolyzed. After detoxification, the phenolic concentration in the hydrolysate was reduced to 0.2 g/L. The sugar yield in the present study is comparable to the results reported by Chandel et al. (2007) and Canilha et al. (2008). Chandel et al. reported 30.29 g/L of total reducing sugars using 2.5 % v/v HCl at 140 °C for 30 min with S:L ratio of 1:10 where as Canilha et al. reported 37 g/L of xylose by dilute acid hydrolysis of wheat straw.

During the course of enzymatic hydrolysis, a regular increase in sugar release was observed till 48 h and remained constant thereafter. The hydrolysate contained  $0.285 \pm 0.005$  g/g sugars ( $35.73 \pm 1.25$  g/L) with a hydrolysis efficiency of  $35.73 \pm 1.25$  % after 48 h of treatment. These results are in accordance with other published results (Junhua et al. 2011). Results of the acid and enzymatic digestibility are shown in Table 2.

### Fermentation of wheat straw hydrolysates

#### Fermentation of acid hydrolysate

The results of sugar consumption and ethanol production in fermentation studies are summarized in Figs. 1, 2, 3 and 4 respectively where as the kinetic parameters are summarized in Table 3. Among the four mutants of *Pichia stipitis* (PSUV9 and PSEB5) and *Candida shehatae* (CSUV4 and CSEB7), PSUV9 showed maximum ethanol production and fermentation efficiency (FE) ( $11.93 \pm 0.38$  g/L, yield  $0.390 \pm 0.008$  g/g, FE  $75.95 \pm 0.26$  %) followed by CSEB7 ( $9.98 \pm 0.81$  g/L, yield  $0.350 \pm 0.005$  g/g, FE  $69.24 \pm 0.18$  %) and PSEB5 ( $9.65 \pm 0.74$  g/L, yield  $0.340 \pm 0.006$  g/g, FE  $66.81 \pm 0.12$  %) in 2 % dilute acid

hydrolysate (Fig. 2). Similar results were obtained by Eken-Sarakoglu and Arslan (2000) with mutant strain of *Pichia stipitis* using corncob hydrolysate. Shi et al. mutated *P. stipitis* CBS 6054 by disrupting the cytochrome c gene which has given 21 % higher ethanol yield (0.46 g/g sugar) than the parental strain (0.38 g/g sugar) from 8 % (w/v) xylose (Shi et al. 1999). Li generated an efficient mutant from *C. shehatae* ATCC 22984 by UV irradiation suggesting that the introduction of a mutation is effective for the improvement of ethanol from 0.39 to 0.42 g/g (Li et al. 2012). Among all the mutants, PSUV9 has given good yields of ethanol and this may be because of its ability of high sugar uptake, ethanol tolerance and inhibitor tolerance in the acid hydrolysate which was studied in the initial characterization studies of mutants. PSUV9 was able to grow after lag phase of 36 h at 1.5 g/L of vanillin, and 6 % of ethanol.

In case of the wild strains, the maximum concentration of ethanol by *Pichia stipitis* (PSP) in dilute acid hydrolysate was found to be  $9.61 \pm 0.39$  g/L equivalent to the yield  $0.330 \pm 0.008$  g/g and fermentation efficiency of  $64.530 \pm 0.248$  %. The yield obtained in this study for ethanol production by parent strain *P. stipitis* NCIM 3498 is comparable to the results reported earlier. Delgenes reported 0.25 g/g of ethanol yield from wheat straw dilute acid hydrolysate by *P. stipitis* (Delgenes et al. 1990). Roberto et al. and Nigam reported the same yield of ethanol i.e., 0.35 g/g from the sugarcane bagasse hydrolysate and *Eichhornia crassipes* by *P. stipitis* NRRL Y-7124 and *P. stipitis* CBS 5773 respectively (Roberto et al. 1991; Nigam 2002). In the present study, the parent strain of *Candida shehatae* (CSP) has produced  $8.35 \pm 0.36$  g/L of ethanol which was equivalent to the yield of  $0.310 \pm 0.007$  g/g and fermentation efficiency of  $60.980 \pm 0.237$  %. Jing-Ping Ge also reported the same yield of ethanol i.e., 0.31 g/g with corncob hydrolysate using *Candida shehatae* ACCC 20335 (Jing-Ping et al. 2011). According to Tanimura, *Candida shehatae* strain ATY839 has produced 16.8 g/L i.e., 71.6 % of the maximum theoretical ethanol yield at 24 h (Ayumi et al. 2012).



#### Fermentation of enzymatic hydrolysate

The fermentability of wheat straw hemicellulosic enzymatic hydrolysate was also evaluated by wild and mutant

**Table 2 Concentration of sugars (g/L) in acid and enzymatic hydrolysates**

S. no.	Type of hydrolysis	Time of incubation (h)	g/L	Saccharification (%)
1.	Enzymatic hydrolysis	18	$16.67 \pm 0.77$	$18.75 \pm 0.56$
		24	$23 \pm 0.55$	$28.75 \pm 0.32$
		48	$35.73 \pm 0.39$	$35.73 \pm 0.65$
2.	Dilute acid hydrolysis (2 %)	–	$39.3 \pm 0.46$	$41.79 \pm 0.62$







strains of *Pichia stipitis* and *Candida shehatae*. Among the wild and mutant strains, PSUV9 showed highest concentration of ethanol ( $12.15 \pm 0.57$  g/L) after 36 h (Fig. 4). However, the ethanol productivity was almost static after 36 h of incubation. The fermentation efficiency was  $87.36 \pm 0.37$  % and the resulting yield of ethanol was equivalent to  $0.450 \pm 0.009$  g/g based on the total fermentable sugars (35.73 g/L) of the hydrolysate. Chandrasekhar Gajula produced a maximum ethanol yield of  $0.44$  g/g with *P. stipitis* NCIM 3498 respectively in batch fermentation conditions using ground nut shell enzyme hydrolysate (Chandrasekhar et al. 2011). We assume that the maximum yield of ethanol in this study is attributed to the occurrence and fermentation of pentoses and some

amount of hexoses in the hydrolysate. Next to PSUV9, CSEB7 showed a maximum ethanol concentration of  $9.55 \pm 0.47$  g/L which accounts for  $0.440 \pm 0.001$  g/g ethanol yield and  $86.09 \pm 0.31$  % of fermentation efficiency followed by PSEB4 with the ethanol concentration of  $9.54 \pm 0.61$  g/L of ethanol which is equivalent to yield of  $0.430 \pm 0.002$  g/g and efficiency of  $84.28 \pm 0.81$  %.

The parent strains of *Pichia stipitis* and *Candida shehatae* have produced  $8.28 \pm 0.54$  and  $7.92 \pm 0.89$  g/L of ethanol which were equivalent to yields of  $0.380 \pm 0.006$  and  $0.370 \pm 0.002$  g/g respectively. It is clearly evident that the yield of ethanol was higher in enzymatic hydrolysate than acid hydrolysate using both wild and mutant strains and this poor fermentability of the acid hydrolysate might



**Table 3** Kinetic parameters for ethanol production from acid and enzymatic hydrolysates using the two best mutant strains (PSUV9, CSEB7) and parent strains (PSP, CSP) of *P. stiptis* NCIM-3498 and *Candida shehatae* NCIM 3501

Parameters	<i>Pichia stiptis</i>				<i>Candida shehatae</i>			
	Acid hydrolysate (48 h)		Enzymatic hydrolysate (36 h)		Acid hydrolysate (48 h)		Enzymatic hydrolysate (36 h)	
	Parent strain (PSP)	Mutant strain (PSUV9)	Parent strain (PSP)	Mutant strain (PSUV9)	Parent strain (CSP)	Mutant strain (CSEB7)	Parent strain (CSP)	Mutant strain (CSEB7)
Ethanol (g/L)	9.61 ± 0.39	11.93 ± 0.38	8.28 ± 0.54	12.15 ± 0.57	8.35 ± 0.36	9.98 ± 0.81	7.92 ± 0.89	9.55 ± 0.47
Sugar utilized (g/L)	29.20 ± 0.25	30.80 ± 0.93	21.84 ± 0.27	27.27 ± 0.42	26.85 ± 0.91	28.26 ± 0.52	21.35 ± 0.40	21.75 ± 0.64
Yield (g/g)	0.330 ± 0.008	0.390 ± 0.008	0.380 ± 0.006	0.450 ± 0.009	0.310 ± 0.007	0.350 ± 0.005	0.370 ± 0.002	0.440 ± 0.001
Productivity (g/L/h)	0.200 ± 0.015	0.240 ± 0.004	0.230 ± 0.005	0.330 ± 0.011	0.170 ± 0.007	0.200 ± 0.012	0.220 ± 0.007	0.260 ± 0.002
% conversion efficiency	64.53 ± 0.24	75.95 ± 0.26	74.34 ± 0.22	87.36 ± 0.37	60.98 ± 0.23	69.24 ± 0.18	72.74 ± 0.39	86.09 ± 0.31

be due to the presence of some toxic components which remained after detoxification, affected the fermentation activity of the yeast.

**Statistical evaluation**

The results of paired t test (Table 4) show that there is significant difference among all the pairs of wild and mutant strains. Since the pair 1(PSP&PSUV9) has more significance compared with the other pairs with t value (125.03) and P value (<0.0001), PSUV9 was found to be the best mutant among all the mutants.

**The stability of the mutant *Pichia stipitis* PSUV9 and *Candida shehatae* CSEB7**

The stability of the mutants *Pichia stipitis* PSUV9 and *Candida shehatae* CSEB7 for increased ethanol production was determined by successive subculturing on MGYE plates for 19 generations (generation time of 48 h). After each subculture, the mutants were tested for

their ability to produce consistent levels of ethanol in acid and enzymatic hydrolysates of wheat straw. The mutants maintained the consistent yields after 19 fermentation cycles indicating that the mutation is stable (Fig. 5).

**Conclusions**

Strain improvement by mutation is one of the best methods to increase the ethanol yield and in this case, we were able to obtain two strains capable of producing significantly higher ethanol yields than the wild strains. The mutant strains PSUV9 and CSEB7 showed higher ethanol production rates from wheat straw as compared to the wild strains. This research demonstrates the utility of random mutagenesis to generate advantageous strains of *P. stipitis* and *C. shehatae*. However, some significant improvements with regard to the two pentose fermenting yeast strains have been reached and these improvements offer new possibilities for further optimization.

**Table 4 Statistical evaluation (paired samples test) of ethanol production (g/L) in acid and enzymatic hydrolysates of wheat straw by wild type mutants of *Pichia Stipitis* and *Candida shehatae***

	Paired differences (dependent sample t test)					
	95 % Confidence interval of the difference					
	Mean	Mean differences	SD	t	P value	Significant Y/N
Pair 1	5.31 7.12	1.81	0.02516	125.03	<0.0001	Y****
Pair 2	5.31 5.07	0.24	0.03214	12.572	0.0063	Y**
Pair 3	4.96 4.23	0.73	0.04509	28	0.0012	Y**
Pair 4	4.96 5.73	0.77	0.0450	29.064	0.0012	Y**
Pair 5	6.91 8.26	1.35	0.0208	112.88	0.0001	Y****
Pair 6	6.91 6.04	0.87	0.03214	46.150	0.0005	Y***
Pair 7	5.93 5.11	0.82	0.02081	67.674	0.0002	Y***
Pair 8	5.93 6.88	0.95	0.06658	25.580	0.00015	Y**

Pair 1: Ethanol production (PSP vs PSUV9) in dilute acid hydrolysate of wheat straw

Pair 2: Ethanol production (PSP vs PSEB5) in dilute acid hydrolysate of wheat straw

Pair 3: Ethanol production (CTP vs CTUV4) in dilute acid hydrolysate of wheat straw

Pair 4: Ethanol production (CTP vs CTEB7) in dilute acid hydrolysate of wheat straw

Pair 5: Ethanol production (PSP vs PSUV9) in enzymatic hydrolysate of wheat straw


Pair 6: Ethanol production (PSP vs PSEB5) in enzymatic hydrolysate of wheat straw

Pair 7: Ethanol production (CTP vs CTUV4) in enzymatic hydrolysate of wheat straw

Pair 8: Ethanol production (CTP vs CTEB7) in enzymatic hydrolysate of wheat straw

Y/N yes/no

\*\*\*\*P ≤ 0.0001; \*\*\*0.0001 > P < 0.0009; \*\*P > 0.0009



**Fig. 5** Stability of the production of ethanol by mutant *Pichia stipitis* PSUV9 in acid and enzymatic hydrolysates

#### Authors' contributions

SK designed and carried out the experiments, performed statistical analysis and drafted the manuscript. SPG participated in design of experiments. JG helped in drafting the manuscript. LV participated in design of the study and revision of the manuscript critically. All authors read and approved the final manuscript.

#### Competing interests

The authors declare that they have no competing interests.

Received: 6 September 2015 Accepted: 5 September 2016

Published online: 13 September 2016

#### References

- Anuj KC, Chandrasekhar G, Radhika K, Ravinder R, Ravindra P (2011) Bioconversion of pentose sugars into ethanol: a review and future directions. *BMBR* 6(1):8–20
- Aristidou A, Penttila M (2000) Metabolic engineering applications to renewable resource utilization. *Curr Opin Biotechnol* 11:187–198
- Ayumi T, Toshihide N, Itsuki W, Jun O, Jun S (2012) Isolation of a novel strain of *Candida shehatae* for ethanol production at elevated temperature. *SpringerPlus* 1:27
- Bailey MJ, Biely P, Poutanen K (1992) Interlaboratory testing of methods for assay of xylanase activity. *J Biotechnol* 23:257–270
- Borbala E, Balazs F, Mats G, Guido Z (2012) Separate hydrolysis and co-fermentation for improved xylose utilization in integrated ethanol production from wheat meal and wheat straw. *Biotechnol Biofuels* 5:12
- Bothast RJ, Nichols NN, Dien BS (2002) Biocatalysts for production of bioethanol. In: Society of industrial microbiology annual meeting 12-1
- Canilha L, Carvalho W, Felipe MGA, Silva JBA (2008) Xylitol production from wheat straw hemicellulosic hydrolysate: hydrolysate detoxification and carbon source used for inoculum preparation. *Braz J Microbiol* 39:333–336
- Chandel AK, Chan EC, Rudravaram R, Narasu ML, Rao LV, Ravindra P (2007) Economics and environmental impact of bioethanol production technologies: an appraisal. *Biotechnol Mol Biol Rev* 2:14–32
- Chandrasekhar G, Anuj Kumar C, Radhika K, Ravinder R, Ravindra P, Lakshmi Narasu M (2011) Fermentation of groundnut shell enzymatic hydrolysate for fuel ethanol production by free and sorghum stalks immobilized cells of *Pichia stipitis* NCIM 3498. *Int J Chem React Eng* 9:Article A61
- Chang V, Holtzapfel M (2000) Fundamental factors affecting biomass enzymatic reactivity. *Appl Biochem Biotechnol* 84–86(1–9):5–37
- Delgenes J, Moletta R, Navarro J (1990) Acid hydrolysis of wheat straw and process considerations for ethanol fermentation by *Pichia stipitis* Y7124. *Proc Biochem* 25:132–135
- Eken-Sarakoglu N, Arslan Y (2000) Comparison of different pretreatments in ethanol fermentation using corn cob hemicellulosic hydrolysate by *Pichia stipitis* and *Candida shehatae*. *Biotechnol Lett* 22(10):855–858
- Gong CS, Claypool TA, McCracken KD, Mann CM, Ueng PP, Tsao GT (1983) Conversion of pentoses by yeast. *Biotechnol Bioeng* 25:85–104
- Jing-Ping G, Bai-Yan C, Guo-Ming L, Hong-Zhi L, Bao-Zhu F, Gang S, Xiao-Feng Y, Wen-Xiang P (2011) Comparison of different detoxification methods for corn cob hemicellulose hydrolysate to improve ethanol production by *Candida shehatae* ATCC 20335. *Afr J Microbiol Res* 5(10):1163–1168
- Joanna C, Ewelina D (2003) Respiratory deficient mutants of xylose-fermenting yeast obtainment and features. *Electr J Pol Agric Univ* 6(2):1
- José CD, Joana P, Susana MP, Lina BH, Belina R, Céu MS (2010) Ethanol production from enzymatically pretreated wheat straw. In: Proceedings of the second international conference of IAMAW, IZMIR, Turkey
- Junhua Z, Matti S, Maija T, Liisa V (2011) The role of acetyl xylan esterase in the solubilization of xylan and enzymatic hydrolysis of wheat straw and giant reed. *Biotechnol Biofuels* 4:60
- Koti S, Prashanthi S, Gentela J, Kothagauni S, Jagavathi S, Venkateswar Rao L (2012) Optimization of pretreatment of wheat straw using alkali and biphasic hydrolysis. *Dyn Biochem Process Biotechnol Mol Biol* 6(1):91–94
- Lee YY, Iyer P, Torget RW (1999) Dilute-acid hydrolysis of lignocellulosic biomass. *Adv Biochem Eng Biotechnol* 65:93–115
- Li Y, J-y Park, Shiroma R, Ike M, Tokuyasu K (2012) Improved ethanol and reduced xylitol production from glucose and xylose mixtures by the mutant strain of *Candida shehatae* ATCC 22984. *Appl Biochem Biotechnol* 166:1781–1790
- Miller GL (1959) Use of dinitrosalicylic acid reagent for determination of reducing sugars. *Anal Chem* 31:426–428
- Mosier NS, Hendrickson R, Dreschel R (2003) Principles and economics of pretreating cellulose in water for ethanol production. Paper No. 103, BIOT Division, 225th American chemical society meeting, New Orleans, March
- Nigam JN (2001) Ethanol production from hardwood spent sulfite liquor using an adapted strain of *Pichia stipitis*. *J Ind Microbiol Biotechnol* 26(3):145–150
- Nigam JN (2002) Bioconversion of water-hyacinth (*Eichhornia crassipes*) hemicellulose acid hydrolysate to motor fuel ethanol by xylose-fermenting yeast. *J Biotechnol* 97:107–116
- Roberto IC, Felipe MGA, Lacys LS, Silva SS, Mancilha IM (1991) Utilization of sugarcane bagasse hemicellulosic hydrolysate by *Candida guilliermondii* for xylitol production. *Biores Technol* 36:271–275
- Sai Prashanthi G, Sravanthi K, Srilekha Yadav K, Venkateshwar S, Venkateswar Rao L (2013) Evaluation of pretreatment methods for enzymatic saccharification of wheat straw for bioethanol production. *Carbohydr Polym* 91(2):646–650
- Shi NQ, Davis FB, Sherman JC, Jeffries TW (1999) Disruption of the cytochrome c gene in xylose-utilizing yeast *Pichia stipitis* leads to higher ethanol production. *Yeast* 15:1021–1030



- Singleton VL, Joseph A, Rossi JR (1965) Colorometry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am J Enol Vitic* 16:144–153
- Srilekha Yadav K, Naseeruddin S, Sai Prashanthi G, Sateesh L, Venkateswar Rao L (2011) Bioethanol fermentation of concentrated rice straw hydrolysate using co-culture of *Saccharomyces cerevisiae* and *Pichia stipitis*. *Biores Technol* 102:6473–6478
- Stephenson PG, Moore M, Terry MJ, Zubkov MV, Bibby TS (2011) Improving photosynthesis for algal biofuels: toward a green revolution. *Trends Biotechnol* 29:615–623
- Sues A, Millati R, Edebo L (2005) Ethanol production from hexoses, pentoses, and dilute-acid hydrolyzate by *Mucor indicus*. *FEMS Yeast Res* 5:669–676
- Winston F, Ausube FM (1990) *Current protocols in molecular biology*, supplement 12. Wiley, New York, pp 13.3.1–13.3.4
- Yan L, Shuzo T (2006) Ethanol fermentation from biomass resources: current state and prospects. *Appl Microbiol Biotechnol* 69(6):627–642

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

---

Submit your next manuscript at ▶ [springeropen.com](http://springeropen.com)

---