

Review began 09/17/2021
Review ended 09/19/2021
Published 09/21/2021

© Copyright 2021

Labib et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Acne Keloidalis Nuchae: A Staged Reconstruction

Amir Labib¹, Louay Salfity², Barry Powell²

1. Plastic Surgery, St. George's University Hospitals NHS Foundation Trust, London, GBR 2. Plastic and Reconstructive Surgery, St. George's University Hospitals NHS Foundation Trust, London, GBR

Corresponding author: Amir Labib, amirlabib2001@gmail.com

Abstract

Acne keloidalis nuchae (AKN) is a chronic scarring folliculitis that affects the occipital area. It is characterized by papule and pustule formation which eventually leads to tumor-like mass. Early cases can be managed by medical treatment that may require months, and, usually, lesions recur. In more severe cases, surgical excision is the mainstay of treatment. In our case, we proposed a staged approach when dealing with advanced stages of AKN. This includes deep excision of the lesion down to deep subcutaneous tissue with application of negative-pressure wound therapy (NPWT) for a week followed by resurfacing of the resultant defect with a split-thickness skin graft and NPWT for another week. This approach achieved quicker wound healing with no recurrence compared with other techniques such as healing by secondary intention. To our knowledge, this case is one of the most extensive cases published in the literature.

Categories: Dermatology, Pathology, Plastic Surgery

Keywords: treatment, therapy, npwt, hair disorder, scalp, acne keloidalis nuchae

Introduction

Acne keloidalis nuchae (AKN) is a chronic form of folliculitis and perifolliculitis. It affects the nape area and results in multiple itchy papules that easily get infected resulting in cicatricial alopecia and keloid-like plaque [1]. AKN is a misnomer as acne is not the cause and there is no true keloid formation. Moreover, the concomitant presence of keloid elsewhere is quite rare. AKN is more common among males of African descent. It was first described by Kaposi in 1869 as “dermatitis papillaris capillitii” [2]. Many theories have been postulated for its etiology, with some describing it as a mechanical form of folliculitis. Other theories relate it to metabolic syndrome [3].

Multiple modalities of treatment have been described for AKN including topical or intralesional steroid, retinoid, a prolonged course of oral antibiotics, especially rifampicin and clindamycin, and radiation. Surgical options include excision and direct closure, excision with healing by secondary intention, excision and immediate grafting, and use of tissue expander. Yet, achieving acceptable results and disease clearance is challenging [4].

Here, we present a two-stage approach in a severe case that showed good disease control with no evidence of recurrence after a long period of follow-up.

Case Presentation

A 58-year-old, white, non-smoker gentleman with a history of type 2 diabetes mellitus presented to the Plastic Surgery Department of St. Georges's Hospital with a 12-month history of small papules in the nape area which he started to notice after changing his barber. His general physician had tried a one-month course of oral antibiotics with no response. On assessment, almost the entire occipital scalp was replaced by a huge swelling with secondary infection (Figure 1).

How to cite this article

Labib A, Salfity L, Powell B (September 21, 2021) Acne Keloidalis Nuchae: A Staged Reconstruction. Cureus 13(9): e18173. DOI 10.7759/cureus.18173



FIGURE 1: Preoperative photo showing the swelling of the entire occipital area.

An oral course of rifampicin and clindamycin was prescribed resulting in good control of secondary infection. Moreover, dapsone and isotretinoin were tried based on the advice of a dermatologist for one year with little benefit. The patient opted for surgical intervention to resolve the disfiguring swelling in the occipital area. The operation was done in two stages. The first stage included excision of the entire lesion and application of negative-pressure wound therapy (NPWT). One week later, the patient underwent a split-thickness skin graft (SSG) and NPWT was reapplied. The postoperative period was uneventful except for the need for postoperative blood transfusion due to significant blood loss intraoperatively and over granulation tissue which was managed by topical steroid cream (Figure 2).



FIGURE 2: Second-week postoperative photo.

Eventually, all areas healed well within less than eight weeks of graft placement with no recurrence after follow-up for eight months (Figure 3).



FIGURE 3: Eighth-month postoperative photo.

Discussion

AKN causes significant morbidity among affected individuals. Scarring, alopecia, and bad smell from recurrent bacterial infection are the most common complaints. Although the pathophysiology is unknown, possible etiologies include close shaving of hair in the nape area, constant mechanical irritation from shirt collars, and chronic low-grade bacterial infection. It may also present as a cutaneous manifestation of

metabolic syndrome [3,5].

In early disease stages, lifestyle modification such as weight loss, avoidance of low haircuts, and avoidance of high-collared shirts along with medical treatment may be helpful. Medical treatment includes topical or intralesional steroids, topical or oral antibiotics, and/or oral isotretinoin. However, more advanced cases rarely respond to these measures. Radiotherapy has a limited role as it may increase the risk of skin cancers and head and neck malignancy, especially in large lesions that may necessitate larger doses of radiation [6].

Surgical excision is the mainstay in the management of advanced cases. Studies have suggested that subfollicular destruction decreases the risk of recurrence; hence, surgeons should aim deep excision down to the deep subcutaneous tissue or even the fascial layer [7]. Reconstructive options after excision include primary closure, secondary intention, or resurfacing by SSG. Primary closure is difficult to achieve in large areas as it may restrict head movement and lead to wounding dehiscence and widening of the scar later on. A secondary intention is another valid option after surgical excision, but prolonged healing of up to 12 weeks in severe cases remains an obstacle [4]. On the other hand, primary resurfacing by SSG carries a higher risk of graft failure as the wound bed may not be suitable for immediate grafting. Pestalardo et al. [8] reported the use of tissue expander in the initial stage, followed by excision of the lesion and advancement of the expanded flap to cover the resultant defect. They managed to control disease progression with no recurrence. Nevertheless, tissue expander complications such as infection and exposure are limiting factors and have been reported [8].

We propose a staged approach that includes primary deep excision down to the deep subcutaneous tissue along with the application of NPWT for one week, followed by resurfacing using SSG and NPWT for another week. We managed to achieve 80% wound healing by two weeks with small areas of over granulation which were managed by topical steroids. Complete wound healing was achieved in less than two months. Further, follow-up at eight months showed no evidence of any new lesions.

Following this staged approach saves the need for prolonged complicated dressing with the discomfort of oozing from a large wound, as in the case of secondary intention. It also improves the status of the wound bed which further increases graft uptake. One of the issues that we need to highlight in our case is that scalpel excision may lead to significant blood loss. Therefore, excision using electrosurgical cautery may be useful [9].

Conclusions

AKN is a challenging condition to treat. Early stages can be managed by medical treatment while surgical excision is the treatment of choice for advanced disease. A staged excision is a useful tool in the armamentarium for the management of AKN, especially in extensive cases. It achieves quicker healing compared with secondary intention and decreases the chances of disease recurrence.

Additional Information

Disclosures

Human subjects: Consent was obtained or waived by all participants in this study. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

References

1. Califano J, Miller S, Frodel J: Treatment of occipital acne keloidalis by excision followed by secondary intention healing. *Arch Facial Plast Surg*. 1999, 1:308-11. [10.1001/archfaci.1.4.308](https://doi.org/10.1001/archfaci.1.4.308)
2. Vasily DB, Breen PC, Miller OF 3rd: Acne keloidalis nuchae: report and treatment of a severe case. *J Dermatol Surg Oncol*. 1979, 5:228-30. [10.1111/j.1524-4725.1979.tb00648.x](https://doi.org/10.1111/j.1524-4725.1979.tb00648.x)
3. Verma SB, Wollina U: Acne keloidalis nuchae: another cutaneous symptom of metabolic syndrome, truncal obesity, and impending/overt diabetes mellitus?. *Am J Clin Dermatol*. 2010, 11:433-6. [10.2165/11537000-000000000-00000](https://doi.org/10.2165/11537000-000000000-00000)
4. Maranda EL, Simmons BJ, Nguyen AH, Lim VM, Keri JE: Treatment of acne keloidalis nuchae: a systematic review of the literature. *Dermatol Ther (Heidelb)*. 2016, 6:363-78. [10.1007/s15555-016-0134-5](https://doi.org/10.1007/s15555-016-0134-5)
5. D'Souza P, Iyer VK, Ramam M: Familial acne keloidalis. *Acta Derm Venereol*. 1998, 78:382. [10.1080/000155598443132](https://doi.org/10.1080/000155598443132)
6. Kantor GR, Ratz JL, Wheeland RG: Treatment of acne keloidalis nuchae with carbon dioxide laser. *J Am Acad Dermatol*. 1986, 14:263-7. [10.1016/s0190-9622\(86\)70031-5](https://doi.org/10.1016/s0190-9622(86)70031-5)
7. Dinehart SM, Herzberg AJ, Kerns BJ, Pollack SV: Acne keloidalis: a review. *J Dermatol Surg Oncol*. 1989, 15:642-7. [10.1111/j.1524-4725.1989.tb03603.x](https://doi.org/10.1111/j.1524-4725.1989.tb03603.x)
8. Pestalardo CM, Cordero A Jr, Ansorena JM, Bestue M, Martinho A: Acne keloidalis nuchae. Tissue expansion treatment. *Dermatol Surg*. 1995, 21:723-4. [10.1111/j.1524-4725.1995.tb00278.x](https://doi.org/10.1111/j.1524-4725.1995.tb00278.x)

9. Beckett N, Lawson C, Cohen G: Electrosurgical excision of acne keloidalis nuchae with secondary intention healing. *J Clin Aesthet Dermatol*. 2011, 4:36-9.