

Corrigendum

Corrigendum to “Prospective Analysis of Lipid Variations in Hyperthyroid Subjects from Lahore, Pakistan”

Muhammad Amir Iqbal ¹, **Shaaf Ahmad** ², **Tamseela Mumtaz**^{1,3}, **Zahra Naseem**¹,
Javeria Malik¹, **Husna Ahmad** ¹ and **Nabila Roohi** ¹

¹*Institute of Zoology, University of the Punjab, Canal Road, Lahore, Punjab 54590, Pakistan*

²*King Edward Medical University/Mayo Hospital, Hospital Road, Lahore, Punjab 54000, Pakistan*

³*Department of Zoology, Government College for Women University, Faisalabad, Punjab, Pakistan*

Correspondence should be addressed to Nabila Roohi; nabilaruhi@gmail.com

Received 7 December 2021; Accepted 7 December 2021; Published 6 January 2022

Copyright © 2022 Muhammad Amir Iqbal et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Prospective Analysis of Lipid Variations in Hyperthyroid Subjects from Lahore, Pakistan” [1], the authors have identified minor errors in Figure 1 and Table 1. The corrected Figure 1 and Table 1 are as follows:




FIGURE 1: Comparison of the levels (mean \pm SEM) of different study parameters (a-f) in control (C), subclinical (S), overt (O), and follow-up (F) groups showing significant differences at * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$.

TABLE 1: A comprehensive presentation of thyroid and lipid profile in studied groups.

Parameters	Control	Subclinical	Overt	Follow-up	P-value
FT ₄ (pmol/L)	17.53 ± 0.36	19.36 ± 0.93	45.65 ± 2.49	14.66 ± 1.08	<0.0001
FT ₃ (pmol/L)	3.91 ± 0.09	4.62 ± 0.20	28.77 ± 1.94	4.96 ± 0.16	<0.0001
TSH (mIU/L)	1.97 ± 0.15	0.09 ± 0.01	0.04 ± 0.007	0.72 ± 0.214	<0.0001
TC (mg/dL)	157.9 ± 2.39	162.5 ± 8.01	101.8 ± 4.84	149.5 ± 7.53	<0.0001
TG (mg/dL)	96.18 ± 3.69	109.6 ± 10.85	87.00 ± 3.57	122.4 ± 8.62	0.03
HDL-C (mg/dL)	62.88 ± 1.56	55.85 ± 3.48	44.00 ± 2.57	57.00 ± 2.57	<0.0001
LDL-C (mg/dL)	109.0 ± 2.64	119.1 ± 6.35	98.68 ± 2.51	123.8 ± 7.12	0.0002
Apo-B (μg/mL)	1310 ± 104.4	1122 ± 206.1	720.4 ± 28.18	1171 ± 97.28	0.0002
Apo-A (μg/mL)	1448 ± 50.74	1360 ± 79.76	1218 ± 58.87	1493 ± 102.8	0.0164

Values are means ± SEM. FT₄: free tetra-iodothyronine; FT₃: free tri-iodothyronine; TSH: thyroid-stimulating hormone; TC: total cholesterol; TG: triglycerides; HDL-C: high-density lipoprotein-C; LDL-C: low-density lipoprotein-C; Apo-B: apolipoprotein-B; Apo-A: apolipoprotein-A.

References

- [1] M. A. Iqbal, S. Ahmad, T. Mumtaz et al., "Prospective Analysis of Lipid Variations in Hyperthyroid Subjects from Lahore, Pakistan," *BioMed Research International*, vol. 2021, Article ID 9936782, 2021.