




# Complete Genome Sequence of the Plant Pathogen *Ralstonia solanacearum* Strain CIAT-078, Isolated in Colombia, Obtained Using Oxford Nanopore Technology

Diana López-Alvarez,<sup>a,b</sup> Ana M. Leiva,<sup>a</sup> Israel Barrantes,<sup>c</sup> Juan M. Pardo,<sup>a</sup> Viviana Dominguez,<sup>a</sup>  Wilmer J. Cuellar<sup>a</sup>

<sup>a</sup>International Center for Tropical Agriculture (CIAT), Cali, Colombia

<sup>b</sup>Faculty of Agricultural Sciences, National University of Colombia, Palmira, Colombia

<sup>c</sup>Institute for Biostatistics and Informatics in Medicine and Ageing Research, Rostock, Germany

**ABSTRACT** Moko is one of the main diseases affecting banana and plantain in Colombia. Here, we report the genome sequence of the causal agent, the bacterium *Ralstonia solanacearum* (Smith) strain CIAT-078, collected in 2004 from affected plantains in central-west Colombia. The assembled genome was obtained using Oxford Nanopore Technology.

*Ralstonia solanacearum* (Smith) belongs to a species complex of soilborne phytopathogenic bacteria that colonize the xylem tissue of hundreds of plant species worldwide (1). It is classified using molecular and biological methods into three species, four phylotypes, and several races; isolates belonging to phylotype II and race 2 cause Moko disease in banana and plantains (2, 3). Moko has been observed in Colombia since 1954 (4); the oldest leaves of affected plants show yellowing and wilting, become necrotic, and eventually collapse. Most importantly, the fruit pulp becomes discolored, causing major commercial losses (5, 6).

For genome sequencing, we selected CIAT-078, a pathogenic strain collected in 2004 from Moko-affected fields (Quindío, Colombia) that was previously characterized at the pathogenicity and biochemical level (7, 8) and used in routine screening for resistance assays (6). CIAT-078 was reactivated in semiselective medium South Africa (SMSA) (9) for 4 days and then in nutrient agar (Difco, USA) for 48 hours at 28°C. The bacteria were grown in LB medium prior to DNA extraction (Puregene Yeast/Bact. kit; Qiagen), and pathogenicity was confirmed by infecting ‘Dominico Harton’ plantains (6). Libraries were prepared from 1 µg of DNA using ligation sequencing kit 1D (catalog number SQK-LSK109) and sequenced using R9.4 chemistry (FLO-MIN106D) (10). Default settings were used for all software unless specified. Raw signals were base called using Guppy v3.4.3, and a total of 291,746 raw reads ( $N_{50}$  4.27 kb) were generated. An assembly of quality-controlled reads (fastq pass) was done with Minimap2 (11) and Racon v1.4.7 (12) using strain UW163 as the reference (GenBank accession numbers [NZ\\_CP012939](https://www.ncbi.nlm.nih.gov/nuccore/NZ_CP012939) and [NZ\\_CP012940](https://www.ncbi.nlm.nih.gov/nuccore/NZ_CP012940)). UW163 was selected due to the quality of its assembly, taxonomic classification (phylotype II), and host plant (plantain) (13). A second assembly was carried out using the UW163-based assembly as a reference. Assembly metrics were calculated with Qualimap v2.2 (14), and the genome was annotated using the Prokaryotic Genome Annotation Pipeline (15).

The circular chromosome of CIAT-078 consists of 3,481,951 base pairs (bp) (G+C content of 66.6%; 70× coverage). It contains 3,238 protein-coding sequences, 50 tRNA genes, 7 rRNA genes, 2 clustered regularly interspaced short palindromic repeat (CRISPR) loci, 5 riboswitches, 3 noncoding RNAs, 1 transfer-messenger RNA gene, and 4 AL1L pseudoknots. The circular megaplasmid consists of 1,907,373 bp (G+C content of 66.8%; 59× coverage) and contains 1,543 protein-coding sequences, 1 tRNA gene,

**Citation** López-Alvarez D, Leiva AM, Barrantes I, Pardo JM, Dominguez V, Cuellar WJ. 2020. Complete genome sequence of the plant pathogen *Ralstonia solanacearum* strain CIAT-078, isolated in Colombia, obtained using Oxford Nanopore Technology. *Microbiol Resour Announc* 9:e00448-20. <https://doi.org/10.1128/MRA.00448-20>.

**Editor** Christina A. Cuomo, Broad Institute

**Copyright** © 2020 López-Alvarez et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Wilmer J. Cuellar, [w.cuellar@cgiar.org](mailto:w.cuellar@cgiar.org).

**Received** 21 April 2020

**Accepted** 11 May 2020

**Published** 28 May 2020

and 2 riboswitches. The genes *rplB*, *mutS*, and *egl*, used to classify the bacteria at the phylotype and sequevar levels (2, 3), were located at positions 1381053 (*rpl*) and 2635266 (*mutS*) of the chromosome and position 1808381 (*egl*) of the megaplasmid. A maximum likelihood phylogeny of these genes grouped CIAT-078 within phylotype IIB sequevar 4 isolates, a subgroup representative of central-west Colombia (16). DUF3313, a recently identified sequence for diagnostics (6), was located at position 3277102 of the chromosome. The complete genome sequence of CIAT-078 will contribute to comparative genome and functional studies.

**Data availability.** The genome sequence of CIAT-078 was deposited in GenBank under accession numbers [CP051295](#) and [CP051296](#). Raw reads were deposited under SRA accession number [SRP250670](#) (BioProject number [PRJNA608676](#)).

## ACKNOWLEDGMENTS

We are grateful to the staff from the Virology and Crop Protection group at CIAT for technical support.

This work was undertaken as part of, and was cofunded by, the CGIAR Research Program on Roots, Tubers, and Bananas and was supported by CGIAR Trust Fund contributors (<https://www.cgiar.org/funders>).

## REFERENCES

- Mansfield J, Genin S, Magori S, Citovsky V, Sriariyanum M, Ronald P, Dow M, Verdier V, Beer SV, Machado MA, Toth I, Salmond G, Foster GD. 2012. Top 10 plant pathogenic bacteria in molecular plant pathology. *Mol Plant Pathol* 13:614–629. <https://doi.org/10.1111/j.1364-3703.2012.00804.x>.
- Fegan M, Prior P. 2005. How complex is the *Ralstonia solanacearum* species complex, p 449–461. In Allen C, Prior P, Hayward AC (ed), *Bacterial wilt disease and the Ralstonia solanacearum* species complex. APS Press, St. Paul, MN.
- Prior P, Ailloud F, Dalsing BL, Remenant B, Sanchez B, Allen C. 2016. Genomic and proteomic evidence supporting the division of the plant pathogen *Ralstonia solanacearum* into three species. *BMC Genomics* 17:90. <https://doi.org/10.1186/s12864-016-2413-z>.
- Castañeda-Sanchez DA, Espinosa-Orrego JA. 2005. Comportamiento e impacto de la enfermedad de moko en la zona de Urabá (Colombia), en las últimas tres décadas y media y propuesta de un índice de riesgo de la enfermedad. *Rev Fac Nac Agron Medellín* 58:2587–2599.
- Ploetz RC. 2008. Black Sigatoka and Moko: impact and spread of two destructive banana diseases in the Caribbean Basin, p 94–102. In *Proceedings of Caribbean Food Crops Society*, Miami, FL. <https://doi.org/10.22004/ag.econ.256503>.
- Pardo JM, López-Alvarez D, Ceballos G, Alvarez E, Cuellar WJ. 2019. Detection of *Ralstonia solanacearum* phylotype II, race 2 causing Moko disease and validation of genetic resistance observed in the hybrid plantain FHIA-21. *Trop Plant Pathol* 44:371–379. <https://doi.org/10.1007/s40858-019-00282-3>.
- Gómez-Calvo EA, Alvarez E, Llano G. 2005. Identificación y caracterización de cepas de *Ralstonia solanacearum* raza 2, agente causante del Moko de plátano en Colombia. *Fitopatol Colomb* 28:71–75.
- Alvarez E, Pantoja A, Gañan L, Ceballos G. 2015. Current status of Moko disease and black Sigatoka in Latin America and the Caribbean, and options for managing them. International Center for Tropical Agriculture (CIAT), Cali, Colombia.
- Engelbrecht MC. 1994. Modification of a semi-selective medium for the isolation and quantification of *Pseudomonas solanacearum*. *ACIAR Bacterial Wilt Newsl* 10:3–5.
- Leiva AM, Siriwan W, Lopez-Alvarez D, Barrantes I, Hemniam N, Saokham K, Cuellar WJ. 2020. Nanopore-based complete genome sequence of Sri Lankan cassava mosaic virus (Geminivirus) from Thailand. *Microbiol Resour Announc* 9:e01274-19. <https://doi.org/10.1128/MRA.01274-19>.
- Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100. <https://doi.org/10.1093/bioinformatics/bty191>.
- Vaser R, Sović I, Nagarajan N, Šikić M. 2017. Fast and accurate de novo genome assembly from long uncorrected reads. *Genome Res* 27:737–746. <https://doi.org/10.1101/gr.214270.116>.
- Ailloud F, Lowe TM, Robène I, Cruveiller S, Allen C, Prior P. 2016. *In planta* comparative transcriptomics of host-adapted strains of *Ralstonia solanacearum*. *PeerJ* 4:e1549. <https://doi.org/10.7717/peerj.1549>.
- Okonechnikov K, Conesa A, García-Alcalde F. 2016. Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* 32:292–294. <https://doi.org/10.1093/bioinformatics/btv566>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Ramírez M, Moncada RN, Villegas-Escobar V, Jackson RW, Ramírez CA. 2020. Phylogenetic and pathogenic variability of strains of *Ralstonia solanacearum* causing moko disease in Colombia. *Plant Pathol* 69:360–369. <https://doi.org/10.1111/ppa.13121>.