

DNA Recombination and Natural Selection Pressure Sustain Genetic Sequence Diversity of the Feline MHC Class I Genes

Naoya Yuhki and Stephen J. O'Brien

From the Laboratory of Viral Carcinogenesis, National Cancer Institute, Frederick, Maryland 21701

Summary

Sequence comparisons of seven distinct MHC class I cDNA clones revealed that feline class I molecules have a remarkable similarity to human HLA genes in their organization of functional domains as well as in the nonrandom partitioning of genetic variability according to the functional constraints ascribed to different regions of the MHC molecule. The distribution of the pattern of sequence polymorphism in the cat as compared with genetic diversity of human and mouse class I genes provides evidence for four coordinate factors that contribute to the origin and sustenance of abundant allele diversity that characterizes the MHC in the species. These include: (a) a gradual accumulation of spontaneous mutational substitution over evolutionary time; (b) selection against mutational divergence in regions of the class I molecule involved in T cell receptor interaction and also in certain regions that interact with common features of antigens; (c) positive selection pressure in favor of persistence of polymorphism and heterozygosity at 57 nucleotide residues that comprise the antigen recognition site; and (d) periodic intragenic (interallelic) and intergenic recombination within the class I genes. We describe a highly conserved 23-bp nucleotide sequence within the coding region of the first α -helix that separates two relatively polymorphic segments located in the $\alpha 1$ domain that may act as a template or "hot spot" for homologous recombination between class I alleles.

The MHC genes encode two classes of cell surface molecules that can present immunological peptides to TCRs (class I and class II molecules) (1, 2). These molecules have extremely polymorphic features in most species (1). According to the x-ray crystallographic model of the human HLA-A2 class I molecule (3), the majority of polymorphic residues of this molecule in human and mouse are located on the molecular surface that comprises the putative antigen binding site (4). This finding, together with the evidence that HLA class I molecules have differential binding capacities for immunological peptides (5), suggests that the polymorphism of MHC class I molecules reflects the capacity of these molecules to bind a wide spectrum of immunological determinants.

Several competing hypotheses have been proposed to explain the evolutionary origin of the high degree of polymorphism; these include (a) a high rate of mutation (6), (b) interlocus recombination or gene conversion (7, 8), (c) overdominant selection (9–13), (d) frequency-dependent selection (11), or (e) some combinations of the above factors. Pursuant to this question, recent comparative analyses of sequence diversity within human and chimpanzee coding regions have revealed that a large fraction of polymorphic residues were found in both human and chimpanzee MHC alleles (14–18). Of 198 different polymorphic sites found in the chimpanzee, 167

(84%) were also found in one or more HLA molecules, suggesting that most of the modern sequence diversity predates the divergence of these two species over 5 million years ago. A similar analysis revealed that murine class II alleles present today in rodents have shared primitive sequence stretches that must have occurred before the divergence of mouse and rat over 10 million years ago (19). The accumulation and persistence of extensive polymorphism for such extended periods of evolutionary time would preclude the necessity for increased mutation rate as an explanation for abundant diversity.

A different approach by Hughes and Nei (12, 13) tested the increased mutation vs. selection hypotheses for MHC polymorphism by analyzing the theoretical predictions of the two competing hypotheses in the context of the functional domains of the MHC molecule. These investigators examined the ratio of synonymous vs. nonsynonymous nucleotide substitutions in the well-defined antigen recognition site (ARS)¹ of human and murine class I allelic transcripts. They reasoned that, if the ARS was the object of overdominant selection in light of its function to recognize, bind, and present

¹ Abbreviations used in this paper: ARS, antigen recognition site; FeLV, feline leukemia virus; FIV, feline immunodeficiency virus.


```

250 P S G E Q R Y T C H V Q H K G L P E P 269
748 CCT TCT GGA GAG CAG CAG TAC ACC TGC CAT GTG CAG CAC AAG GGG CTG CCC GAG CCC 807
FLA110
FLA124
FLA189
FLA198
FLA223
FLA1
FLA11
FLA12
FLA13
FLA14
FLA15
FLA16
FLA17
FLA18
FLA19
FLA20
FLA21
FLA22
FLA23
FLA24
FLA25
FLA26
FLA27
FLA28
FLA29
FLA30
FLA31
FLA32
FLA33
FLA34
FLA35
FLA36
FLA37
FLA38
FLA39
FLA40
FLA41
FLA42
FLA43
FLA44
FLA45
FLA46
FLA47
FLA48
FLA49
FLA50
FLA51
FLA52
FLA53
FLA54
FLA55
FLA56
FLA57
FLA58
FLA59
FLA60
FLA61
FLA62
FLA63
FLA64
FLA65
FLA66
FLA67
FLA68
FLA69
FLA70
FLA71
FLA72
FLA73
FLA74
FLA75
FLA76
FLA77
FLA78
FLA79
FLA80
FLA81
FLA82
FLA83
FLA84
FLA85
FLA86
FLA87
FLA88
FLA89
FLA90
FLA91
FLA92
FLA93
FLA94
FLA95
FLA96
FLA97
FLA98
FLA99
FLA100
FLA101
FLA102
FLA103
FLA104
FLA105
FLA106
FLA107
FLA108
FLA109
FLA110
FLA111
FLA112
FLA113
FLA114
FLA115
FLA116
FLA117
FLA118
FLA119
FLA120
FLA121
FLA122
FLA123
FLA124
FLA125
FLA126
FLA127
FLA128
FLA129
FLA130
FLA131
FLA132
FLA133
FLA134
FLA135
FLA136
FLA137
FLA138
FLA139
FLA140
FLA141
FLA142
FLA143
FLA144
FLA145
FLA146
FLA147
FLA148
FLA149
FLA150
FLA151
FLA152
FLA153
FLA154
FLA155
FLA156
FLA157
FLA158
FLA159
FLA160
FLA161
FLA162
FLA163
FLA164
FLA165
FLA166
FLA167
FLA168
FLA169
FLA170
FLA171
FLA172
FLA173
FLA174
FLA175
FLA176
FLA177
FLA178
FLA179
FLA180
FLA181
FLA182
FLA183
FLA184
FLA185
FLA186
FLA187
FLA188
FLA189
FLA190
FLA191
FLA192
FLA193
FLA194
FLA195
FLA196
FLA197
FLA198
FLA199
FLA200
FLA201
FLA202
FLA203
FLA204
FLA205
FLA206
FLA207
FLA208
FLA209
FLA210
FLA211
FLA212
FLA213
FLA214
FLA215
FLA216
FLA217
FLA218
FLA219
FLA220
FLA221
FLA222
FLA223
FLA224
FLA225
FLA226
FLA227
FLA228
FLA229
FLA230
FLA231
FLA232
FLA233
FLA234
FLA235
FLA236
FLA237
FLA238
FLA239
FLA240
FLA241
FLA242
FLA243
FLA244
FLA245
FLA246
FLA247
FLA248
FLA249
FLA250
FLA251
FLA252
FLA253
FLA254
FLA255
FLA256
FLA257
FLA258
FLA259
FLA260
FLA261
FLA262
FLA263
FLA264
FLA265
FLA266
FLA267
FLA268
FLA269
FLA270
FLA271
FLA272
FLA273
FLA274
FLA275
FLA276
FLA277
FLA278
FLA279
FLA280
FLA281
FLA282
FLA283
FLA284
FLA285
FLA286
FLA287
FLA288
FLA289
FLA290
FLA291
FLA292
FLA293
FLA294
FLA295
FLA296
FLA297
FLA298
FLA299
FLA300
FLA301
FLA302
FLA303
FLA304
FLA305
FLA306
FLA307
FLA308
FLA309
FLA310
FLA311
FLA312
FLA313
FLA314
FLA315
FLA316
FLA317
FLA318
FLA319
FLA320
FLA321
FLA322
FLA323
FLA324
FLA325
FLA326
FLA327
FLA328
FLA329
FLA330
FLA331
FLA332
FLA333
FLA334
FLA335
FLA336
FLA337
FLA338
FLA339
FLA340
FLA341
FLA342
FLA343
FLA344
FLA345
FLA346
FLA347
FLA348
FLA349
FLA350
FLA351
FLA352
FLA353
FLA354
FLA355
FLA356
FLA357
FLA358
FLA359
FLA360
FLA361
FLA362
FLA363
FLA364
FLA365
FLA366
FLA367
FLA368
FLA369
FLA370
FLA371
FLA372
FLA373
FLA374
FLA375
FLA376
FLA377
FLA378
FLA379
FLA380
FLA381
FLA382
FLA383
FLA384
FLA385
FLA386
FLA387
FLA388
FLA389
FLA390
FLA391
FLA392
FLA393
FLA394
FLA395
FLA396
FLA397
FLA398
FLA399
FLA400
FLA401
FLA402
FLA403
FLA404
FLA405
FLA406
FLA407
FLA408
FLA409
FLA410
FLA411
FLA412
FLA413
FLA414
FLA415
FLA416
FLA417
FLA418
FLA419
FLA420
FLA421
FLA422
FLA423
FLA424
FLA425
FLA426
FLA427
FLA428
FLA429
FLA430
FLA431
FLA432
FLA433
FLA434
FLA435
FLA436
FLA437
FLA438
FLA439
FLA440
FLA441
FLA442
FLA443
FLA444
FLA445
FLA446
FLA447
FLA448
FLA449
FLA450
FLA451
FLA452
FLA453
FLA454
FLA455
FLA456
FLA457
FLA458
FLA459
FLA460
FLA461
FLA462
FLA463
FLA464
FLA465
FLA466
FLA467
FLA468
FLA469
FLA470
FLA471
FLA472
FLA473
FLA474
FLA475
FLA476
FLA477
FLA478
FLA479
FLA480
FLA481
FLA482
FLA483
FLA484
FLA485
FLA486
FLA487
FLA488
FLA489
FLA490
FLA491
FLA492
FLA493
FLA494
FLA495
FLA496
FLA497
FLA498
FLA499
FLA500
FLA501
FLA502
FLA503
FLA504
FLA505
FLA506
FLA507
FLA508
FLA509
FLA510
FLA511
FLA512
FLA513
FLA514
FLA515
FLA516
FLA517
FLA518
FLA519
FLA520
FLA521
FLA522
FLA523
FLA524
FLA525
FLA526
FLA527
FLA528
FLA529
FLA530
FLA531
FLA532
FLA533
FLA534
FLA535
FLA536
FLA537
FLA538
FLA539
FLA540
FLA541
FLA542
FLA543
FLA544
FLA545
FLA546
FLA547
FLA548
FLA549
FLA550
FLA551
FLA552
FLA553
FLA554
FLA555
FLA556
FLA557
FLA558
FLA559
FLA560
FLA561
FLA562
FLA563
FLA564
FLA565
FLA566
FLA567
FLA568
FLA569
FLA570
FLA571
FLA572
FLA573
FLA574
FLA575
FLA576
FLA577
FLA578
FLA579
FLA580
FLA581
FLA582
FLA583
FLA584
FLA585
FLA586
FLA587
FLA588
FLA589
FLA590
FLA591
FLA592
FLA593
FLA594
FLA595
FLA596
FLA597
FLA598
FLA599
FLA600
FLA601
FLA602
FLA603
FLA604
FLA605
FLA606
FLA607
FLA608
FLA609
FLA610
FLA611
FLA612
FLA613
FLA614
FLA615
FLA616
FLA617
FLA618
FLA619
FLA620
FLA621
FLA622
FLA623
FLA624
FLA625
FLA626
FLA627
FLA628
FLA629
FLA630
FLA631
FLA632
FLA633
FLA634
FLA635
FLA636
FLA637
FLA638
FLA639
FLA640
FLA641
FLA642
FLA643
FLA644
FLA645
FLA646
FLA647
FLA648
FLA649
FLA650
FLA651
FLA652
FLA653
FLA654
FLA655
FLA656
FLA657
FLA658
FLA659
FLA660
FLA661
FLA662
FLA663
FLA664
FLA665
FLA666
FLA667
FLA668
FLA669
FLA670
FLA671
FLA672
FLA673
FLA674
FLA675
FLA676
FLA677
FLA678
FLA679
FLA680
FLA681
FLA682
FLA683
FLA684
FLA685
FLA686
FLA687
FLA688
FLA689
FLA690
FLA691
FLA692
FLA693
FLA694
FLA695
FLA696
FLA697
FLA698
FLA699
FLA700
FLA701
FLA702
FLA703
FLA704
FLA705
FLA706
FLA707
FLA708
FLA709
FLA710
FLA711
FLA712
FLA713
FLA714
FLA715
FLA716
FLA717
FLA718
FLA719
FLA720
FLA721
FLA722
FLA723
FLA724
FLA725
FLA726
FLA727
FLA728
FLA729
FLA730
FLA731
FLA732
FLA733
FLA734
FLA735
FLA736
FLA737
FLA738
FLA739
FLA740
FLA741
FLA742
FLA743
FLA744
FLA745
FLA746
FLA747
FLA748
FLA749
FLA750
FLA751
FLA752
FLA753
FLA754
FLA755
FLA756
FLA757
FLA758
FLA759
FLA760
FLA761
FLA762
FLA763
FLA764
FLA765
FLA766
FLA767
FLA768
FLA769
FLA770
FLA771
FLA772
FLA773
FLA774
FLA775
FLA776
FLA777
FLA778
FLA779
FLA780
FLA781
FLA782
FLA783
FLA784
FLA785
FLA786
FLA787
FLA788
FLA789
FLA790
FLA791
FLA792
FLA793
FLA794
FLA795
FLA796
FLA797
FLA798
FLA799
FLA800
FLA801
FLA802
FLA803
FLA804
FLA805
FLA806
FLA807
FLA808
FLA809
FLA810
FLA811
FLA812
FLA813
FLA814
FLA815
FLA816
FLA817
FLA818
FLA819
FLA820
FLA821
FLA822
FLA823
FLA824
FLA825
FLA826
FLA827
FLA828
FLA829
FLA830
FLA831
FLA832
FLA833
FLA834
FLA835
FLA836
FLA837
FLA838
FLA839
FLA840
FLA841
FLA842
FLA843
FLA844
FLA845
FLA846
FLA847
FLA848
FLA849
FLA850
FLA851
FLA852
FLA853
FLA854
FLA855
FLA856
FLA857
FLA858
FLA859
FLA860
FLA861
FLA862
FLA863
FLA864
FLA865
FLA866
FLA867
FLA868
FLA869
FLA870
FLA871
FLA872
FLA873
FLA874
FLA875
FLA876
FLA877
FLA878
FLA879
FLA880
FLA881
FLA882
FLA883
FLA884
FLA885
FLA886
FLA887
FLA888
FLA889
FLA890
FLA891
FLA892
FLA893
FLA894
FLA895
FLA896
FLA897
FLA898
FLA899
FLA900
FLA901
FLA902
FLA903
FLA904
FLA905
FLA906
FLA907
FLA908
FLA909
FLA910
FLA911
FLA912
FLA913
FLA914
FLA915
FLA916
FLA917
FLA918
FLA919
FLA920
FLA921
FLA922
FLA923
FLA924
FLA925
FLA926
FLA927
FLA928
FLA929
FLA930
FLA931
FLA932
FLA933
FLA934
FLA935
FLA936
FLA937
FLA938
FLA939
FLA940
FLA941
FLA942
FLA943
FLA944
FLA945
FLA946
FLA947
FLA948
FLA949
FLA950
FLA951
FLA952
FLA953
FLA954
FLA955
FLA956
FLA957
FLA958
FLA959
FLA960
FLA961
FLA962
FLA963
FLA964
FLA965
FLA966
FLA967
FLA968
FLA969
FLA970
FLA971
FLA972
FLA973
FLA974
FLA975
FLA976
FLA977
FLA978
FLA979
FLA980
FLA981
FLA982
FLA983
FLA984
FLA985
FLA986
FLA987
FLA988
FLA989
FLA990
FLA991
FLA992
FLA993
FLA994
FLA995
FLA996
FLA997
FLA998
FLA999
FLA1000

```

```

1216 GTCTTACTTCATGTTGGGCTGAGTACTGAGCGCTACTCGCTACTGAAATGAAATCTGATGAAGATTGTGTT 1294
FLA110
FLA124
FLA189
FLA198
FLA223
FLA1
FLA11
FLA12
FLA13
FLA14
FLA15
FLA16
FLA17
FLA18
FLA19
FLA20
FLA21
FLA22
FLA23
FLA24
FLA25
FLA26
FLA27
FLA28
FLA29
FLA30
FLA31
FLA32
FLA33
FLA34
FLA35
FLA36
FLA37
FLA38
FLA39
FLA40
FLA41
FLA42
FLA43
FLA44
FLA45
FLA46
FLA47
FLA48
FLA49
FLA50
FLA51
FLA52
FLA53
FLA54
FLA55
FLA56
FLA57
FLA58
FLA59
FLA60
FLA61
FLA62
FLA63
FLA64
FLA65
FLA66
FLA67
FLA68
FLA69
FLA70
FLA71
FLA72
FLA73
FLA74
FLA75
FLA76
FLA77
FLA78
FLA79
FLA80
FLA81
FLA82
FLA83
FLA84
FLA85
FLA86
FLA87
FLA88
FLA89
FLA90
FLA91
FLA92
FLA93
FLA94
FLA95
FLA96
FLA97
FLA98
FLA99
FLA100
FLA101
FLA102
FLA103
FLA104
FLA105
FLA106
FLA107
FLA108
FLA109
FLA110
FLA111
FLA112
FLA113
FLA114
FLA115
FLA116
FLA117
FLA118
FLA119
FLA120
FLA121
FLA122
FLA123
FLA124
FLA125
FLA126
FLA127
FLA128
FLA129
FLA130
FLA131
FLA132
FLA133
FLA134
FLA135
FLA136
FLA137
FLA138
FLA139
FLA140
FLA141
FLA142
FLA143
FLA144
FLA145
FLA146
FLA147
FLA148
FLA149
FLA150
FLA151
FLA152
FLA153
FLA154
FLA155
FLA156
FLA157
FLA158
FLA159
FLA160
FLA161
FLA162
FLA163
FLA164
FLA165
FLA166
FLA167
FLA168
FLA169
FLA170
FLA171
FLA172
FLA173
FLA174
FLA175
FLA176
FLA177
FLA178
FLA179
FLA180
FLA181
FLA182
FLA183
FLA184
FLA185
FLA186
FLA187
FLA188
FLA189
FLA190
FLA191
FLA192
FLA193
FLA194
FLA195
FLA196
FLA197
FLA198
FLA199
FLA200
FLA201
FLA202
FLA203
FLA204
FLA205
FLA206
FLA207
FLA208
FLA209
FLA210
FLA211
FLA212
FLA213
FLA214
FLA215
FLA216
FLA217
FLA218
FLA219
FLA220
FLA221
FLA222
FLA223
FLA224
FLA225
FLA226
FLA227
FLA228
FLA229
FLA230
FLA231
FLA232
FLA233
FLA234
FLA235
FLA236
FLA237
FLA238
FLA239
FLA240
FLA241
FLA242
FLA243
FLA244
FLA245
FLA246
FLA247
FLA248
FLA249
FLA250
FLA251
FLA252
FLA253
FLA254
FLA255
FLA256
FLA257
FLA258
FLA259
FLA260
FLA261
FLA262
FLA263
FLA264
FLA265
FLA266
FLA267
FLA268
FLA269
FLA270
FLA271
FLA272
FLA273
FLA274
FLA275
FLA276
FLA277
FLA278
FLA279
FLA280
FLA281
FLA282
FLA283
FLA284
FLA285
FLA286
FLA287
FLA288
FLA289
FLA290
FLA291
FLA292
FLA293
FLA294
FLA295
FLA296
FLA297
FLA298
FLA299
FLA300
FLA301
FLA302
FLA303
FLA304
FLA305
FLA306
FLA307
FLA308
FLA309
FLA310
FLA311
FLA312
FLA313
FLA314
FLA315
FLA316
FLA317
FLA318
FLA319
FLA320
FLA321
FLA322
FLA323
FLA324
FLA325
FLA326
FLA327
FLA328
FLA329
FLA330
FLA331
FLA332
FLA333
FLA334
FLA335
FLA336
FLA337
FLA338
FLA339
FLA340
FLA341
FLA342
FLA343
FLA344
FLA345
FLA346
FLA347
FLA348
FLA349
FLA350
FLA351
FLA352
FLA353
FLA354
FLA355
FLA356
FLA357
FLA358
FLA359
FLA360
FLA361
FLA362
FLA363
FLA364
FLA365
FLA366
FLA367
FLA368
FLA369
FLA370
FLA371
FLA372
FLA373
FLA374
FLA375
FLA376
FLA377
FLA378
FLA379
FLA380
FLA381
FLA382
FLA383
FLA384
FLA385
FLA386
FLA387
FLA388
FLA389
FLA390
FLA391
FLA392
FLA393
FLA394
FLA395
FLA396
FLA397
FLA398
FLA399
FLA400
FLA401
FLA402
FLA403
FLA404
FLA405
FLA406
FLA407
FLA408
FLA409
FLA410
FLA411
FLA412
FLA413
FLA414
FLA415
FLA416
FLA417
FLA418
FLA419
FLA420
FLA421
FLA422
FLA423
FLA424
FLA425
FLA426
FLA427
FLA428
FLA429
FLA430
FLA431
FLA432
FLA433
FLA434
FLA435
FLA436
FLA437
FLA438
FLA439
FLA440
FLA441
FLA442
FLA443
FLA444
FLA445
FLA446
FLA447
FLA448
FLA449
FLA450
FLA451
FLA452
FLA453
FLA454
FLA455
FLA456
FLA457
FLA458
FLA459
FLA460
FLA461
FLA462
FLA463
FLA464
FLA465
FLA466
FLA467
FLA468
FLA469
FLA470
FLA471
FLA472
FLA473
FLA474
FLA475
FLA476
FLA477
FLA478
FLA479
FLA480
FLA481
FLA482
FLA483
FLA484
FLA485
FLA486
FLA487
FLA488
FLA489
FLA490
FLA491
FLA492
FLA493
FLA494
FLA495
FLA496
FLA497
FLA498
FLA499
FLA500

```

Figure 1. Nucleotide sequences of seven distinct feline class I cDNA clones. The complete sequences are shown for the FLA110 clone. The dashed line and single letter indicate the identical and distinct nucleotide from the FLA110 clone. The deduced amino acid sequence of the FLA110 cDNA clone is shown above the nucleotide sequence of FLA110. Asterisk, plus, and dot symbols above the sequence of FLA110 indicate residues toward the antigen binding site of antigen recognition, on an α -helix pointing up, and on an α -helix pointing away from the antigen binding site and TCR recognition site, respectively, according to the homologous structure of the human HLA-A2 molecule (3, 4).

various peptides to TCRs, then the nonsynonymous (amino acid altering) substitutions would be more abundant than synonymous nucleotide substitutions in the same region. Pairwise comparisons of 12 human and 9 murine class I cDNA allelic transcripts revealed a marked excess in nonsynonymous substitutions compared with synonymous changes in the ARS. Since a model of increased spontaneous mutation under a presumption of selective neutrality would not favor either type of nucleotide substitutions, they interpreted their results as strong evidence against the increased mutation explanation and in favor of overdominant selection.

Parham et al. (15), in an extensive comparative analysis of 40 human HLA class I transcripts and 9 chimpanzee ChLA class I transcripts, presented further evidence for the ancient origin of allelic diversity. Their results also revealed a "patchwork" pattern of diversity between allelic transcripts where common stretches of sequence identity between transcripts were apparent. These results suggested that interallelic recombination likely played a major role in generation of novel class I alleles in human-ape evolution. Intergenic recombination, sometimes referred to as "gene conversion" in the context of MHC diversity observed in rodent class I genes, was thought to be of minor importance since transcripts of each human locus, i.e., HLA-A, -B, and -C, each had numerous shared derived (or synapomorphic) nucleotide residues that were characteristic of the three loci (14, 15). The logic argued that if interlocus exchanges were frequent then diversity between loci would be "homogenized", but that was clearly not the case in man. In contrast, the rodent class I locus were highly homogenized apparently by the inter-locus sequence exchanges (7, 20).

We present in this paper a reexamination of these hypotheses using a new set of class I MHC transcripts derived from the domestic cat. This species has several advantages for studies of MHC diversity. Cats have a moderately polymorphic MHC, called FLA, that has been studied from both serological (21) and molecular perspectives (22–24). In addition, the cat is afflicted by epidemics of two pathological RNA viruses: feline leukemia virus, FeLV (25); and feline immunodeficiency virus, FIV (26, 27). Finally, the cat and its free-ranging relatives have been the object of ongoing studies of population genetics and molecular evolution (28–30). The present results affirm and extend, in a third mammalian order, the critical roles of positive selection and DNA recombination in the origins and maintenance of genetic diversity of the MHC. In addition, although locus specificity of FLA class I transcripts are evident in the conserved transmembrane and cytoplasmic domains, occasional mosaic sequences indicate that inter-locus sequence exchange has played a notable role in the evolution of feline MHC class I genes.

Materials and Methods

Total RNA was isolated from three unrelated domestic cat spleen cell preparations by the hot phenol-guanidium isothiocyanate method (31). Poly(A)⁺ RNAs were isolated by oligo(dT)-cellulose columns (32). cDNAs were synthesized by a modification of the method of Gubler and Hoffman (33). These cDNAs were ligated

with a λ ZAP II vector (34) and in vitro packaged using a Giga-pack plus reagent (Stratagene). 2×10^5 unamplified plaques from each library were screened with a ³²P-labeled feline class I cDNA insert (pFLA24) (23) at low stringency. Positive single plaques were isolated by the second screening and cDNA clones were autoexcised into phagemid forms by the infection of R408 helper phages (34). The phagemid clone DNAs were electrophoresed on 1% agarose gel, stained with ethidium bromide and the sizes were measured by comparing them with the full-length cDNA clone, pFLA24. The full-length or nearly full-length clones were then transfected into JM101 *Escherichia coli* cells, superinfected with M13K07 helper phages or the cDNAs were subcloned into M13 vectors and single-strand DNAs for sequence analysis were purified (35). The sequences of each clone were determined by a dideoxynucleotide sequencing method using [³⁵S]dATP (New England Nuclear, Boston, MA) and T7 DNA polymerase (36) (U.S. Biochemical Corp., Cleveland, OH), and seven sequencing primers for the conserved sequences of two previously isolated feline class I cDNA clones (23). These sequences were compared using the Wilbur and Lipman algorithm (37).

The names of two previously described transcripts, FLA2, FLA24 (23), were changed to FLAB2 and FLAA24, respectively, to reflect locus-specificity in this paper. The seven cDNA clones described here were isolated from libraries made from four different cats as follows: cat no. 1, FLAB2 and FLAA24; cat no. 2, FLAA10, FLAA23 and FLAB9; cat no. 3, FLAA1; and cat no. 4, FLAX8.

Results

Five distinct full-length or nearly full-length feline MHC class I cDNA clones were identified from three unrelated cat spleen cDNA libraries by a screen of over 60 cDNA clones. Together with two previously described, distinct feline MHC class I cDNA clones (23), we determined the nucleotide sequence of seven distinct cDNA transcripts in order to examine the diversity and the evolution of class I molecules. MHC class I genes had been reported to consist of eight coding exons which encode the following protein domains: (1) leader sequence; (2) and (3) $\alpha 1$ and $\alpha 2$ polymorphic extracellular domains which are involved in antigen and TCR recognitions; (4) $\alpha 3$ extracellular domain; (5) transmembrane domain; and (6–8) three cytoplasmic domains. According to the x-ray crystallographic analysis of human HLA-A2 class I molecules, Bjorkman et al. (4) have described 57 amino acid residues located in the $\alpha 1$ and $\alpha 2$ domains that are involved in antigen recognition and are termed the antigen recognition site (ARS). Further, the ARS occurs in two α -helix stretches that are separated by one platform of β -pleated sheets in the class I molecule. In the present analysis we compare the complete sequences for the seven feline cDNA clones (Figs. 1 and 2). The regions of protein topography (α helices and β -pleated sheets) are indicated in the sequence of Fig. 2 according to the structure of the HLA-A2 molecule.

We first examined the first two extracellular domains. Analysis of the pattern of sequence polymorphism in FLA transcripts revealed that, as in the human and mouse, there are clearly recognizable residues that are invariant and others that are highly polymorphic. Further, the variant and invariant amino acids are generally at the same positions in the three species. Table 1 lists 32 residues that are invariant among 40




Figure 2. Amino acid sequences of the seven distinct feline class I molecules. The complete sequence of FLAA10 is shown. Single letters and dashed lines indicate the different residues and the identical residues with those of FLAA10, respectively. Two α -helix coding regions and eight β -sheet coding regions are indicated below the sequences. The same definitions as for Fig. 1 are used for the asterisk, plus, and dot symbols above the amino acid sequence. H and M above the number indicate polymorphic sites defined in human and mouse class I molecules respectively. Highly polymorphic sites of feline class I molecules are defined as the site that showed over three different residues and are indicated as underlines of the FLAA10 sequence.

human transcripts. Of these, 31 are also invariant in cats and 27 of the 32 conserved residues have identical amino acids in human and feline transcripts. Four cat residues were different from the human residues but were still conserved within the feline molecules (L5, I73, N158, A169). The single highly polymorphic feline residue that was invariant in humans (Q155) is also polymorphic in mouse H-2 transcripts (23). The majority of highly polymorphic residues of feline class I molecules (9 of 10 highly polymorphic sites which show over three different residues) are located on the site facing the antigen binding groove according to the model of the HLA-A2 molecule.

The pattern and distribution of sequence substitution between allelic transcripts can be used to test for the role of mutation versus the selection of MHC diversity based upon the procedures of Hughes and Nei (12, 13). If the MHC molecules are subject to positive selection, and if the 57 residues of the ARS that participate directly in antigen recognition are the objects of selective pressure, the rate of nonsynonymous mutation will be higher than that of synonymous substitution in ARS. The logic is predicated upon the theoretical notion that positive selection would enhance the rate of "sen-

sible" codon substitution compared with that for synonymous substitutions because synonymous mutations are more or less selectively neutral. By contrast, if high polymorphism is a consequence of increased mutation, the rates of synonymous vs. nonsynonymous mutation would be equivalent in all regions. In a comparison of FLA transcripts, there occurred nearly twice as many nonsynonymous substitutions as synonymous substitutions in the ARS region (Table 2). Of the 21 pairwise comparisons between the 7 FLA transcripts, in every case there was a greater number of nonsynonymous substitutions in this functional region. By contrast, the same analysis of the 125 residues that are in the $\alpha 1$ and $\alpha 2$ domains, but are not in the ARS, resulted in the opposite observation, namely that the frequency of synonymous substitutions were greater by nearly twofold. These findings are in precise agreement with the same analysis of human and murine nucleotide substitutions (Table 2). Taken together the data provide strong evidence for the role of overdominant selection or heterozygote advantage in the ARS region of the MHC molecules in the three species.

A close examination of the highly polymorphic region in the $\alpha 1$ domain that encodes the first α -helix (amino acid

Table 1. Highly Conserved and Polymorphic Residues in the $\alpha 1$ and $\alpha 2$ Domains of Human and Cat Class I Molecules

	Criteria	Human	Cat
Highly conserved residues	Recognition of common features of Ag	M5, Y7, F22, G26, Y59, Y84, T143, K146, Y159, Y171	<u>L5</u> , Y7, F22, G26, Y59, Y84, T143, K146, Y159, Y171
	Formation of conserved faces of α -helixes	P57, E58, D61, T64, Q72, R75, L78, Y84, K146, A150, E154, Q155, G162, V165	P57, E58, D61, T64, Q72, R75, L78, Y84, K146, A150, E154, (G162), V165
		T73, A158, R169	<u>I73</u> , <u>N158</u> , <u>A169</u>
	N-Glycosylation site	N86, Q87, S88	N86, Q87, S88
	Disulfide bonding	C101, C164	C101, C164
Highly polymorphic residues	β -sheets in $\alpha 1$ domain	9	None
	First α -helix	62, 65, 66, 67, 70, 71, 77, 80	63, 67, 77, 79*, 81
	β -sheets in $\alpha 2$ domain	95, 97, 114, 116	97, 104, 116
	Second α -helix	156, 163	155, 156

Probable functional constraints are based on homology to human residues (3, 4). Uniquely conserved residues in feline class I molecules are underlined. One residue (G162) is conserved in six of seven feline molecules. 9 of 10 highly polymorphic residues (over 3 different residues at the site) are located on the site facing the antigen binding groove according to the human HLA-A2 model (with the exception of one residue, 79*).

residues 58–86) demonstrated a patchwork or mosaic pattern between FLA transcripts (Fig. 2). FLAA24, FLAB9, and FLAX8 were identical to FLAA10 in the NH₂-terminal half of the α -helix while the same molecules diverged ap-

preciably in the COOH-terminal half of this region. The remaining three transcripts (FLAA23, FLAA1, and FLAB2) were divergent from FLAA10 in the NH₂-terminal half, while they were identical to FLAA10 in the COOH-terminal

Table 2. Mean Numbers of Nucleotide Substitutions* per 100 Synonymous Sites (dS) and per 100 Nonsynonymous Sites (dN) of MHC Class I Molecules

Species	No. of sequences	No. of comparisons	Antigen recognition sites (ARS) (n = 57)		Remaining codons in exon 2 & 3 (n = 125)	
			dS	dN	dS	dN
Domestic cat	7	21	7.5 ± 2.4	13.60 ± 1.9 [†]	7.34 ± 2.4	4.9 ± 0.8
				0:21		16:5
Human	12	66	6.8 ± 2.3	20.8 ± 2.3 [§]	11.6 ± 2.1	5.2 ± 0.8 [‡]
				0:66		63:3
Mouse	9	36	13.8 ± 3.2	21.4 ± 2.4	8.3 ± 1.6	6.5 ± 0.8
				6:30		29:7

* By the method of Nei and Gojobori (38). For each species the ratio of pairwise sequence comparisons when synonymous substitutions exceeded nonsynonymous substitutions (dS>dN) vs. the reverse situation is presented as dS>dN:dN>dS. Human and mouse data are derived from (12).

[†] Significant at 0.1% level.

[§] Significant at 0.01% level.

region (Fig. 2). When the nucleotide sequences of these same transcripts were examined (Figs. 1 and 3), they revealed highly polymorphic regions in the 5' and 3' ends of the first α -helix region separated by a strongly conserved 23-nucleotide stretch. We tested the sequence divergence of the two divergent segments of the α -helix for evidence of recombination vs. mutation origins using the algorithm of Stephens (39). This method effectively discriminates random mutational substitution vs. block changes that occur by recombination between divergent sequence. In comparison of the 5' and 3' segments of FLAA10 transcript to homologous regions of each of the other transcripts, the results affirmed divergent clustering ($p \leq 0.05$) in 11 of 12 comparisons lending strong credence to the recombinational origin of the FLA transcripts. Additional vestiges of recombination were apparent in comparisons of the $\alpha 2$ domains where the cDNAs did not conform to partitioning of transcripts evident in the $\alpha 1$ domain (Fig. 2 and Table 3).

The 3' domains of the mammalian MHC class I molecule, ($\alpha 3$ extracellular, transmembrane, and cytoplasmic) are generally conserved in their sequence and have been used to assign locus specificity of class I transcripts (15). The seven FLA transcripts were compared in this region and fell into three discrete classes (Table 3 and Fig. 4). The first group (FLAA10, A24, A23, and A1) were nearly identical throughout the three 3' domains (only three nucleotide substitutions in 441 nucleotides), but quite divergent from the remaining transcripts. The second group contained FLAB2 and B9 which were similar, though not identical, in the TM and CY domains but more divergent in the $\alpha 3$ extracellular domain. Based upon the precedence in humans whereby alleles of the same class I B and C loci vary by a maximum of eight nucleotide substitutions (15), we hypothesize that these two classes of transcripts are products of different loci, designated A and B, respectively. Support for this assignment is obtained because FLAB9, A10, and A23 were derived from a single individual cat. In that animal, we suggest the A locus expressed A10 plus A23 and the B locus expressed B9. The last transcript, FLAX8, has a transmembrane domain sequence that was nearly identical to the B locus FLAB9 (one substitution), a cytoplasmic domain that was more similar to the A locus FLAA1 and A23 sequences (3 substitutions), and an $\alpha 3$ domain that was equidistant from all other transcripts. The mosaic sequence structure of the FLAX8 transcript makes it difficult to assign it unequivocally to either locus. An illustration of the shared domains of the seven transcripts and their locus designation is presented in Fig. 4.

A reexamination of the mosaic sequence structure in the α -helix region (Fig. 3) armed with the knowledge of locus specificity of the transcripts is illuminating. The FLAA10 specific 5' and 3' sequences are found in transcripts from both A and B loci. If our locus designations are correct, this indicates that inter-allelic exchange occurs not only within, but also between, class I loci. Alternatively, the FLAA10-derived sequences in this region are primitive and the variants are derived by recombination with a different ancestral donor gene sequence. In either case, intra- or intergenic recombination has occurred and the conserved 23 bp intermediate sequence served as a junction for the exchange. Further, the mosaic structure within the $\alpha 2$, transmembrane, and cytoplasmic domains (Fig. 4) would also affirm a role for recombination in deriving these arrangements.

Discussion

The present sequence comparison of seven feline MHC class I transcripts to mouse and human genes revealed a remarkable conservation in the patterns of domain organization and in the sites of intraspecies polymorphisms. With only one exception, the FLA class I domains were precisely the length of the human HLA-A and -C domains (15, 22), and there was a striking conservation of sequence homology for 32 invariant human residues in $\alpha 1$ and $\alpha 2$ domains known to play a critical role in MHC class I function (Table 1). The cat molecules not only were genetically invariant at each of the sites, but they also retained identical residues at 27 of the 32 sites. In addition the feline molecules displayed a clustering of polymorphic residues within the two α -helix and two β -strand regions in a manner similar to the pattern of human and mouse sequences (4). Finally, strong evidence against increased mutation rate and for the force of selection of polymorphic sites in the 57 residues of the antigen recognition site was obtained by the demonstrable excess of codon-altering (as compared with synonymous) substitution rate in this region. An earlier analysis of human and mouse transcripts yielded a very similar result (12, 13). We conclude that the class I genes of the three mammalian species retain ancestral constraints both in their organization of functional domains as well as in their tolerance (or predilection) for polymorphism. These shared primitive (ancestral) constraints likely reflect a functional organization inherited from a common ancestor of the three mammalian orders.

Based on the cumulative results of our studies of the human, mouse, and cat sequences, we can recognize four major factors

FIRST α -HELIX CODING REGION

	22bp	23bp	21bp	LOCUS
FLAA10	CGGAACACCCGGATTACTTGG	ACACCGCACAGATTTCCGAGTG	GACCTGAACACGATGCTCCGC	A
FLAA24	-----G-----	-----C-----	A-----T-T-G----	A
FLAB9	-----G-----	-----T-----	AG---G-G-AC-C--G-G-A	B
FLAX8	-----G-----	-----T-----	A-----C-G--CGCC-----	AB
FLAA23	---G-G--G---ACATGAA-A	-----T-----	-----G-----	A
FLAA1	---G-G--G---ACATGAA-A	-----T-----	-----G-----	A
FLAB2	GA-C-G--G---ACG-GAA-A	---A-----	-----G-----	B

Figure 3. Nucleotide sequences that encode the first long α -helix region. Dashed line indicates identical nucleotide to FLAA10 sequence.

Table 3. The Number of Nucleotide Substitutions

		A10	A24	A23	A1	B9	B2*	X8
Leader	FLAA10		2	3	3	0	—	2
	FLAA24			3	2	0	—	0
	FLAA23				4	2	—	3
	FLAA1					1	—	2
	FLAB9						—	0
	FLAB2							—
α 1 Domain	FLAA10		9	20	21	19	(16)	11
	FLAA24			22	22	17	(20)	10
	FLAA23				1	27	(10)	25
	FLAA1					27	(10)	27
	FLAB9						(26)	16
	FLAB2							(24)
α 2 Domain	FLAA10		16	16	18	28	25	20
	FLAA24			6	8	33	17	13
	FLAA23				2	29	15	9
	FLAA1					27	14	11
	FLAB9						38	23
	FLAB2							20
α 3 Domain	FLAA10		0	1	1	10	7	7
	FLAA24			1	1	10	7	7
	FLAA23				0	8	6	7
	FLAA1					9	6	6
	FLAB9						8	9
	FLAB2							8
TM Domain	FLAA10		0	0	0	8	6	7
	FLAA24			0	0	8	6	7
	FLAA23				0	8	6	7
	FLAA1					8	6	7
	FLAB9						4	1
	FLAB1							3
CY Domain	FLAA10		2	0	0	8	6	7
	FLAA24			2	2	10	10	5
	FLAA23				0	8	8	3
	FLAA1					8	8	3
	FLAB9						4	9
	FLAB2							9
3'UT	FLAA10		4	3	4	20	31	15
	FLAA24			3	3	22	33	13
	FLAA23				3	24	34	12
	FLAA1					21	34	16
	FLAB9						26	28
	FLAB2							42

* B2cDNA clone begins within the α 1 domain.




Figure 4. Schematic features of seven feline class I molecules. FLAA23 and A1 are subtypes that show only two amino acid substitutions (indicated by asterisks).

in the generation and maintenance of diversity. The first is a stochastic accumulation of spontaneous mutations throughout the class I molecule. There is little persuasive evidence for an increased frequency of mutations in these genes (12). The second key factor is a discernable negative selection pressure against increased polymorphism in two important regions; namely, the areas including the conserved outer face of the α -helix that interacts directly with the complementation determining regions (CDR1 and CDR2) of the TCR V gene product (2) and the conserved regions of the antigen binding site that recognize a common feature of immunological peptides (4). Residues from both of these regions are invariant within the three species and in most cases retain identical amino acids in each of these sites (see Table 1).

The third important factor is a positive selective pressure in favor of increased heterozygosity and polymorphism at loci in the 57 nucleotide residues in the ARS. Evidence for positive selection was obtained by comparing the ratio of codon-altering (nonsynonymous) substitution rate to synonymous substitution rate in the same region. As predicted by a model of overdominant selection by Hughes and Nei (12), the extent of nonsynonymous substitution rate in the ARS was significantly higher than the synonymous substitution rate (Table 2). The reverse was true in the neighboring residues not involved in antigen recognition, apparently because those regions are not subject to the intense selection for maintenance of genetic diversity required by the sites of antigen recognition. The findings with FLA class I molecules were in exact agreement with the results obtained by Hughes and Nei (12) who compared human and mouse class I transcripts. The results may also reveal a level of frequency dependent selection; however, it is difficult to implicate this precisely with the present data.

The fourth critical component for generation of MHC diversity is intragenic (interallelic) and intergenic recombination. Evidence for both types of recombination as an origin for allelic diversity was evident in the first α -helix where a patchwork or mosaic pattern of shared amino acid diversity was observed (Figs. 1 and 2). Examination of the seven cat nucleotide sequences revealed a 23-bp segment between recom-

binant stretches that is likely the junction of previous recombinant events that generated the previous transcripts (Fig. 3). In addition, shared stretches of mutational differences in several FLA transcripts in the $\alpha 2$ domain are likely derived by recombination between the $\alpha 1$ and $\alpha 2$ domain as well (Fig. 1). Finally, the mosaic structure in transmembrane and cytoplasmic domains of one FLA transcript are likely derived by intergenic recombination between these two domains (Fig. 1 and 4). Because the mosaic patterns do not partition between putative class I loci, these combined observations would suggest a pivotal role for both interallelic and intergenic recombination in the formulation of FLA class I allele diversity.

The recent extensive studies of human transcripts are resounding in their finding of within-locus association in contrast to between-locus similarity (15, 16). Locus specificity actually extends beyond species boundaries since HLA-A alleles more closely resemble chimpanzee ChLA-A alleles than either group resembles their respective B-locus alleles (15). Locus-specific partitioning of allele sequences would not be consistent with a large role for intergenic recombination because that process would tend to homogenize alleles rather than separate them. Murine class I transcripts tend to be more homogenized and lack clear locus-specific sequence variation

(7, 20), suggesting a historic role for intergenic exchange in rodent gene evaluation. The feline class I sequences presented here support the occurrence of between locus but within coding region recombinants in the generation of class I sequence variations.

A role for recombination in the generation of diversity would probably be adaptive since the types of interallelic recombination or intergenic recombination between functional loci would be able to create numerous polymorphic, but functional, molecules since the functional constraints were already overcome in prototype molecules (15). Such a mechanism might actually be more efficient than a random mutational process for generating new diversity. It is tempting to speculate that certain sequences of notable conservation located between regions of high variation may act as templates for homologous intra- and intergenic recombination particularly for those regions conserved in the three species discussed here. The 23-bp region of the first α -helix would be a possible example of this phenomenon. This sequence may have been an important recombination junction and may still retain the potential to create further polymorphic molecules in the future by homologous recombination.

Received for publication 29 January 1990 and in revised form 2 May 1990.

References

1. Klein, J. 1986. Natural History of the Major Histocompatibility Complex. John Wiley and Sons Inc., New York, NY.
2. Davis, M.M., and P.J. Bjorkman. 1988. T-cell antigen receptor genes and T-cell recognition. *Nature (Lond.)* 334:395.
3. Bjorkman, P.J., M.A. Saper, B. Samraoui, W.S. Bennett, J.L. Strominger, and D.C. Wiley. 1987. Structure of the human class I histocompatibility antigen, HLA-A2. *Nature (Lond.)* 329:506.
4. Bjorkman, P.J., M.A. Saper, B. Samraoui, W.S. Bennett, J.L. Strominger, and D.C. Wiley. 1987. The foreign antigen binding site and T-cell recognition regions of class I histocompatibility antigens. *Nature (Lond.)* 329:512.
5. Chen, B.P., and P. Parham. 1989. Direct binding of influenza peptides to class I HLA molecules. *Nature (Lond.)* 337:743.
6. Klein, J. 1978. H-2 mutations: their genetics and effect on immune functions. *Adv. Immunol.* 26:55.
7. Nathenson, S.G., J. Geliebter, G.M. Pfaffenbach, and R.A. Zeff. 1986. Murine major histocompatibility complex class-I mutants: molecular analysis and structure-function implications. *Annu. Rev. Immunol.* 4:471.
8. Mellor, A.L., E.H. Weiss, K. Ramachandran, and R.A. Flavell. 1983. A potential donor gene for the bm1 gene conversion event in the C57BL mouse. *Nature (Lond.)* 306:792.
9. Doherty, P.C., and R.M. Zinkernagel. 1975. Enhanced immunological surveillance in mice heterozygous at the H-2 gene complex. *Nature (Lond.)* 256:50.
10. Klein, J., and F. Figueroa. 1986. Evolution of the major histocompatibility complex. *CRC Crit. Rev. Immunol.* 6:295.
11. Bodmer, W.F. 1972. Evolutionary significance of the HL-A system. *Nature (Lond.)* 237:139.
12. Hughes, A.L., and M. Nei. 1988. Pattern of nucleotide substitution at major histocompatibility complex class I loci reveals overdominant selection. *Nature (Lond.)* 335:167.
13. Hughes, A.L., and M. Nei. 1989. Nucleotide substitution at major histocompatibility complex class II loci: evidence for overdominant selection. *Proc. Natl. Acad. Sci. USA.* 86:958.
14. Parham, P., C.E. Lomen, D.A. Lawlor, J.P. Ways, N. Holmes, H.L. Coppin, R.D. Salter, A.M. Wan, and P.D. Ennis. 1988. Nature of polymorphism in HLA-A, -B, and -C molecules. *Proc. Natl. Acad. Sci. USA.* 85:4005.
15. Parham, P., D.A. Lawlor, C.E. Lomen, and P.D. Ennis. 1989. Diversity and diversification of HLA-A, -B, -C alleles. *J. Immunol.* 142:3937.
16. Mayer, W.E., M. Jonker, D. Klein, P. Ivanyi, G. van Seventer, and J. Klein. 1988. Nucleotide sequences of chimpanzee MHC class I alleles: Evidence for trans-species mode of evolution. *EMBO (Eur. Mol. Biol. Organ.) J.* 7:2765.
17. Lawlor, D.A., F.E. Ward, P.D. Ennis, A.P. Jackson, and P. Parham. 1988. HLA-A and -B polymorphisms predate the divergence of humans and chimpanzees. *Nature (Lond.)* 335:268.
18. Klein, J., J. Gutknecht, and N. Fischer. 1990. The major histocompatibility complex and human evolution. *Trends Genet.* 6:7.
19. Figueroa, F., E. Gunther, and J. Klein. 1988. MHC polymorphism pre-dating speciation. *Nature (Lond.)* 335:265.
20. Rada, C., R. Lorenzi, S-J. Powis, J. van den Bogaerde, P.

- Parham, and J.C. Howard. 1990. Concerted evolution of class I genes in the major histocompatibility complex of murine rodents. *Proc. Natl. Acad. Sci. USA.* 87:2167.
21. Winkler, C., A. Shultz, S. Cevario, and S.J. O'Brien. 1989. Genetic characterization of FLA, the cat major histocompatibility complex. *Proc. Natl. Acad. Sci. USA.* 86:943.
 22. Yuhki, N., and S.J. O'Brien. 1988. Molecular characterization and genetic mapping of class I and class II MHC genes of the domestic cat. *Immunogenetics.* 16:339.
 23. Yuhki, N., G.F. Heidecker, and S.J. O'Brien. 1989. Characterization of MHC cDNA clones in the domestic cat. Diversity and evolution of class I genes. *J. Immunol.* 142:3676.
 24. Yuhki, N., and S.J. O'Brien. 1990. DNA variation of the mammalian major histocompatibility complex reflects genomic diversity and population history. *Proc. Natl. Acad. Sci. USA.* 87:836.
 25. Cotter, S.M., W.D. Hardy, Jr., and M. Essex. 1975. Association of feline leukemia virus with lymphosarcoma and other disorders in the cat. *J. Am. Vet. Med. Assoc.* 166:449.
 26. Pedersen, N.C., E.W. Ho, M.L. Brown, and J.K. Yamamoto. 1987. Isolation of a T-lymphotropic virus from domestic cats with an immunodeficiency-like syndrome. *Science (Wash. DC).* 235:790.
 27. Olmsted, R.A., A.K. Barnes, J.K. Yamamoto, V.M. Hirsch, R.H. Purcell, and P.R. Johnson. 1989. Molecular cloning of feline immunodeficiency virus. *Proc. Natl. Acad. Sci. USA.* 86:2448.
 28. O'Brien, S.J., D.E. Wildt, D. Goldman, C.R. Merrill, and M. Bush. 1983. The cheetah is depauperate in genetic variation. *Science (Wash. DC).* 221:459.
 29. O'Brien, S.J., M.E. Roelke, L. Marker, A. Newman, C.A. Winkler, D. Meltzer, L. Colly, J.F. Evermann, M. Bush, and D.E. Wildt. 1985. Genetic basis for species vulnerability in the cheetah. *Science (Wash. DC).* 227:1428.
 30. O'Brien, S.J. 1986. Molecular genetics in the domestic cat and its relatives. *Trends Genet.* 2:137.
 31. Maniatis, T., E.-F. Fritsch, and J. Sambrook. 1982. *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York. 545 pp.
 32. Davis, L.G., M.D. Dibner, and J.F. Battey. 1986. *Basic Methods in Molecular Biology*. Elsevier, New York. 139 pp.
 33. Gubler, U., and B.J. Hoffman. 1983. A simple and very efficient method for generating cDNA libraries. *Gene (Amst.).* 25:263.
 34. Short, J.M., J.M. Fernandez, J.A. Sorge, and W.D. Huse. 1988. λ ZAP: a bacteriophage λ expression vector with in vivo excision properties. *Nucleic Acids Res.* 16:7583.
 35. Vieira, J., and J. Messing. 1987. Production of single-stranded plasmid DNA. *Methods Enzymol.* 153:3.
 36. Tabor, S., and C.C. Richardson. 1987. DNA sequence analysis with a modified bacteriophage T7 DNA polymerase. *Proc. Natl. Acad. Sci. USA.* 84:4767.
 37. Wilbur, W.J., and D.J. Lipman. 1983. Rapid similarity searches of nucleic acid and protein data banks. *Proc. Natl. Acad. Sci. USA.* 80:726.
 38. Nei, M., and T. Gojobori. 1986. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Mol. Biol. Evol.* 3:418.
 39. Stephens, J.C. 1985. Statistical methods of DNA sequence analysis: detection of intragenic recombination or gene conversion. *Mol. Biol. Evol.* 2:539.