

1 **Three dominant *Vibrio cholerae* lytic phage all require O1 antigen for infection**

2

3 **Drew A. Beckman and Christopher M. Waters***

4

5 Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing,

6 Michigan, USA, 48824

7

8 Running Title: *V.c.* O-Antigen Mutants Pan-Resistant to ICP Phage

9

10 ***Corresponding Author:**

11 5180 Biomedical and Physical Sciences

12 567 Wilson Road

13 East Lansing, MI 48824

14 Telephone 517-884-5360

15 E-mail: watersc3@msu.edu

16 [Keywords:](#)

17

18 **SUMMARY**

19 In its natural aquatic environment, the bacterial pathogen *Vibrio cholerae*, the causative

20 agent of the enteric disease cholera, is in constant competition with bacterial viruses known as

21 phages. Following ICP3 infection, *V. cholerae* cultures that exhibited phage killing always

22 recovered overnight, and clones isolated from these regrowth populations exhibited complete

23 resistance to subsequent infections. Whole genome sequencing of these resistant mutants

24 revealed seven distinct mutations in genes encoding for enzymes involved in O1 antigen

25 biosynthesis, demonstrating that the O1 antigen is a previously uncharacterized receptor of ICP3.

26 To further elucidate the specificity of the resistance conferred by these mutations, they were

27 challenged with the *V. cholerae*-specific phages ICP1 and ICP2. Despite no prior exposure to
28 these phages, all seven O1 antigen mutants demonstrated pan-resistance to all three ICP phages.
29 Given that the O1 antigen is required for infection of *V. cholerae*, our results show that the ICP
30 phage have evolved to recognize an essential surface molecule for infection, providing a barrier
31 to the evolution of pathogenic *V. cholerae* phage resistance in natural environments.

32

33 **MAIN**

34 **Introduction**

35 The Gram-negative aquatic pathogen *Vibrio cholerae*, the causative agent of the severe
36 enteric diarrheal disease cholera, is primarily found in the saline conditions of estuaries and
37 brackish water (1). Recent estimates from 2015 place the human burden of disease worldwide
38 in the range of 1.3-4.0 million cases and approximately 95,000 deaths annually as a result of
39 rapid dehydration (2). Countries in sub-Saharan Africa and southern Asia are most directly
40 impacted by cholera, but natural and manmade humanitarian disasters have ignited outbreaks
41 in Haiti and Iraq (3, 4). Cholera is undoubtedly worsened by poverty where water treatment
42 systems are suboptimal or non-existent (5).

43 Within its environment, *V. cholerae* is preyed upon by bacterial viruses known as
44 bacteriophages, or phages, and these parasites are significant drivers of *V. cholerae* evolution
45 (6). Lytic phages commandeer the host cell's machinery and replicate to high numbers,
46 eventually lysing the host cell. This predation is a critical driver of co-evolution among *V.*
47 *cholerae* bacteria and their respective phages, applying selective pressure for the evolution of
48 phage defense mechanisms like abortive infection (7), mobile genetic elements (8), and toxin-
49 antitoxin systems (9) in addition to counter-defense systems evolved by the phage (10). Despite
50 this robust phage defense, there are three major *V. cholerae* specific lytic phage families
51 isolated from patient stool samples at the International Center for Diarrheal Disease Research,
52 Bangladesh known as ICP1, ICP2, and ICP3 (11, 12). These three phages, while identified

53 together, are quite diverse; ICP1 is a member of the *Myoviridae* family and has a genome 2-3
54 times larger than ICP2 and ICP3 which are members of the *Podoviridae* family (12). The ICP
55 phages are the three largest of the nine major phage families that have been identified to infect
56 *V. cholerae* (13).

57 Before phage can successfully infect host bacteria, they must first bind to a specific cell
58 surface receptor that determines phage specificity and host range. The receptor targeted by
59 ICP1 to mediate phage invasion and replication is the O1 antigen in the lipopolysaccharide
60 (LPS) of the *V. cholerae* outer membrane (11, 12). Of the over 200 known serogroups, only O1
61 and O139 are pathogenic with the former responsible for the greatest number of cases (11, 14,
62 15). The El Tor biotype of *V. cholerae*, which is responsible for the current seventh cholera
63 pandemic, can be subdivided into serotypes based on methylation of the O1 antigen. The
64 Ogawa serotype is categorized by an additional methyl group on the most distal perosamine
65 monomer while this methyl is absent in the Inaba serotype (4, 14, 15). The principal receptor for
66 ICP2 is the OmpU outer membrane porin (16). However, the receptor for ICP3 remains
67 uncharacterized (3, 17). These structures are excellent targets for phage exploitation because
68 of their key roles in human pathogenesis and antimicrobial resistance. For instance, OmpU is
69 required for efficient colonization of hosts while O1 antigen deficient *V. cholerae* mutants are
70 more sensitive to the antibiotic polymyxin B as well as the bactericidal/permeability increasing
71 (BPI) antimicrobial peptide secreted by the human gut (18, 19).

72 Understanding the interaction between the ICP phages and *V. cholerae* is not only
73 essential for characterizing the evolution and ecology of *V. cholerae* in its natural environments,
74 but also for the effective development of new cholera treatment strategies. Phage therapy is
75 emerging as an effective complement to antibiotic treatment, especially with the rise of antibiotic
76 resistant strains (20–22). The phage tropism for its individual receptor limits the negative effects
77 on the human host and their microbiome (23). Two phage therapy approaches are being
78 developed to treat cholera including the Phi_1 phage (24) and a cocktail of ICP1, ICP2, and

79 ICP3 (3). Therefore, identifying the receptors of these phage has important clinical ramifications.

80 Here, we used a forward genetics approach to identify the previously unknown ICP3
81 receptor by examining several *V. cholerae* escape mutants following ICP3 infection. Our results
82 showed that O1 antigen biosynthesis enzymes were mutated in all ICP3 resistant strains, and
83 surprisingly these mutants exhibited pan-resistance to ICP1, ICP2, and ICP3, demonstrating
84 that the O1 antigen is a conserved receptor for all three ICP phages.

85


86 **Results**

87 **Isolation of ICP3 resistant *V. cholerae***

88 During experiments studying the interaction of ICP3 phage with two newly discovered *V.*
89 *cholerae* phage defense systems (7, 9), we observed that after an initial sharp decrease in the
90 culture optical density upon ICP3 addition to values nearly equivalent to blank media controls,
91 the bacterial cultures always rebounded to densities resembling the uninfected cultures. To
92 establish the growth kinetics of *V. cholerae* in liquid media infected with ICP3 phage, an
93 overnight culture of WT cells was subcultured into ten replicates, five of which were
94 subsequently infected with ICP3 during mid-exponential growth phase at a multiplicity of
95 infection (MOI) of 1.0 and the other five at 0.1 (Fig. 1). The density of the cultures rapidly
96 decreased to initial inoculum levels 0.75 hr or 1.0 hr after ICP3 addition with 1.0 MOI or 0.1
97 MOI, respectively (Fig. 1). However, the OD₆₀₀ of these cultures began to rebound 2 to 3 hr
98 following the maximal drop. 24 hours later, the infected and uninfected cultures exhibited nearly
99 identical optical densities (Fig. 1).

100

101



102

103 **Figure 1: Infected *V. cholerae* cultures spontaneously recover over time.**

104 Growth curves of WT *V. cholerae* cultures challenged with ICP3 phage at varying multiplicities
105 of infection (MOI). ICP3 phage were added at 0 hr as indicated by the arrow. Data points are the
106 mean of five biological replicates in the case of the phage groups (n = 5), and two in the case of
107 the no phage control (n = 2). Error bars represent the standard deviation.

108

109 We hypothesized that the cells able to propagate in the phage-infected cultures were
110 escape mutants resistant to ICP3. We isolated an individual clone from each of the ten
111 independent overnight cultures that had rebounded from ICP3 infection. Each isolate was
112 rechallenged with ICP3 at the original MOI of 1.0 for A-E (Fig. 2A) and 0.1 for F-J (Fig. 2B). Four
113 of the five isolates infected with an MOI of 1.0 were resistant to ICP3 phage predation (Fig. 2A)
114 while all five isolates challenged at an MOI of 0.1 were able to withstand ICP3 reinfection (Fig.
115 2B).

116

117


118

119

120


121

122 **A**



123

124 **B**



125

126 **Figure 2: Recovered isolates demonstrate resistance to ICP3 phage in liquid culture.**

127 From the recovery pools of cultures that rebounded from ICP3 phage infection, individual
128 isolates were obtained and rechallenged in a similar manner. They were either infected at an
129 MOI of 1.0 (A) or 0.1 (B) based on their original recovery pool. ICP3 phage, or SM buffer in the
130 case of the no phage control, was added at 0 hr as indicated by the arrow.

131

132

133

134 **Resistant mutations are in O1 antigen biosynthesis genes**

135 Whole genome sequencing of all nine resistant isolates identified that isolates A-I (excluding the
136 non-resistant isolate B) shared the same 11 bp deletion in the *manB* gene while isolate J had an
137 insertion in the *wbeU* gene (Table 1, Fig. 3). Both genes are part of the O1 antigen biosynthesis
138 operons of *V. cholerae* (14, 25). Given that the identical mutation in *manB* was isolated from 8
139 independent cultures, we hypothesized this mutation was likely present in the original common
140 starting overnight culture. Therefore, to isolate additional resistance mutations, five independent
141 overnight cultures of WT *V. cholerae* were inoculated using individual colonies and subjected to
142 ICP3 infection at an MOI of 1.0. All five cultures exhibited regrowth after the initial population
143 decline. Individual clones from these five populations that were resistant to ICP3 rechallenge
144 were isolated and their genomes were sequenced (isolates K-O). In each isolate, there was only
145 one unique mutation, all of which mapped to the O1 antigen biosynthesis pathway (Table 1, Fig.
146 3). Importantly, no other resistant mutations outside of the O1 biosynthesis genes were isolated.
147 These seven unique ICP3 resistant mutants in the O1 antigen biosynthesis pathway were
148 present in six different genes. Six of these mutations are likely null either by causing frameshift
149 mutations (isolates A, J, L, N, and O) or a significant deletion (isolate K) while isolate M has a
150 missense mutation in *gmd* (Table 1). Given the localization of these resistance-conferring
151 mutations, we conclude that ICP3 requires the O1 antigen for infection of *V. cholerae*.

152

153

154

155

156

157

158

159 **Table 1: Whole Genome Sequencing of Pan-Resistant Isolates Reveals Mutations in the**
 160 **O-Antigen Biosynthesis Pathway**

Isolate	Gene ^a	Position ^b	Location In Gene	Mutation	Protein Product ^c
A, C-I	<i>manB</i> , VC0242	Chr I. 2,696,238	920-930/1392 nt	Δ11 bp	Phosphomannomutase
J	<i>wbeU</i> , <i>rfbU</i> , VC0259	Chr I. 2,681,562	571/1113 nt	+G	Glycosyltransferase / LPS Biosynthesis Protein
K	<i>wbeN</i> , <i>rfbN</i> , VC0251	Chr I. 2,687,710	78-95/2478 nt	Δ18 bp	Acyl-CoA Reductase / Acyl Protein Synthase
L	<i>wbeN</i> , <i>rfbN</i> , VC0251	Chr I. 2,685,438	2367/2478 nt	+5 bp	Acyl-CoA Reductase / Acyl Protein Synthase
M*	<i>gmd</i> , VC0243	Chr I. 2,695,683	91/1122 nt	G→A (H31Y)	GDP-Mannose 4,6-Dehydratase / Oxidoreductase
N	<i>manC</i> , VC0241	Chr I. 2,698,253	315/1398 nt	ΔC	Mannose-1-Phosphate Guanylyltransferase / Mannose-6-Phosphate Isomerase
O	<i>wbeM</i> , VC0250	Chr I. 2,688,092	828/1125 nt	+A	Iron-containing Alcohol Dehydrogenase


161 ^aGene names were determined as part of the *wbe* cluster (14, 15, 25) and the *rfb* cluster (27, 36).

162 ^bNZ_CP046844 (Chr I) and NZ_CP046845 (Chr II) used as reference genomes.

163 ^cPutative enzyme functions based on these reference genomes as well as Pombo *et al*, 2022 where applicable (37).

164 *Demonstrated partial resistance in solid culture.

165



166

167 **Figure 3: O1 antigen biosynthesis pathway genes indicating sites of mutations in**
 168 **resistant isolates.**

169 Diagram depicting the genes of the *wbe* locus (*rfb* locus) in O1 serotypes of *V. cholerae* strains.

170 The block arrows of each gene point in the direction of transcription with gene sizes drawn to
171 scale. Multiple names of each gene are given where applicable. The total length from the start
172 codon of *VC0241* to the stop codon of *VC0263* is provided. Vertical arrows indicate the location
173 of mutations in the individual isolates identified by whole genome sequencing.

174

175

176 **ICP3 resistant mutants are also resistant to ICP1 and ICP2 infection**

177 To investigate if these seven ICP3 resistant mutants demonstrated cross-resistance to
178 ICP1 and ICP2, they were grown in liquid culture to exponential phase and exposed to ICP1,
179 ICP2, or ICP3. In these conditions, all the resistant mutants grew similar to the WT strain in the
180 absence of phage (Fig. 4A) and were resistant to ICP3 infection (Fig. 4D). Interestingly, all
181 seven ICP3 resistant mutants were also completely resistant to ICP1 ($p < 0.0001$) and ICP2 ($p <$
182 0.0001) infection in liquid infection assays (Figs. 4B, C).

183

184

185

186

187

188

189

190


191

192

193

194


195




211 solid medium. An example of this plaquing assay for both the sensitive WT control and Isolate A
212 is shown (Fig. 5A). Of the seven previously resistant *V. cholerae* mutants, six were completely
213 resistant to ICP1 ($p = 0.0005$), ICP2 ($p < 0.0001$), and ICP3 ($p = 0.0012$) showing no plaques at
214 any dilution of phage tested (Figs. 5B-D). The WT and isolate M exhibited identical numbers of
215 plaques, suggesting this isolate does not exhibit equivalent resistance in plaque-based assays
216 to that seen in liquid culture experiments. However, the plaques of ICP1, ICP2, or ICP3 that
217 formed on isolate M were significantly cloudier than the WT strain, suggesting this isolate has
218 partial resistance to phage infection in these conditions (Fig. 5A). Despite no prior exposure to
219 ICP1 and ICP2, these ICP3 resistant isolates were able to survive predation indicating all three
220 phage require the O1 antigen for infection.

221

222 **A**



223



237 Restoration of O1 antigen biosynthesis restores phage susceptibility

238 To ensure that phage resistance was due to the sequenced mutations, two representative

239 isolates, A and J, were chosen and complemented by constructing inducible plasmids encoding

240 *manB* and *wbeU*, respectively (Table I). Liquid infection assays compared the complemented

241 strains and a deficient strain containing an empty vector. ICP1, ICP2, or ICP3 phage were

242 added at time 0 with a MOI of 1.0 (Fig. 6). In each case, the resistant isolates containing the


243 complemented plasmid exhibited significant phage killing after 1 hr whereas the corresponding

244 empty vector control strains were unimpacted by phage addition. In each experiment, there was


245 also a significant growth defect prior to phage infection for isolate J (*wbeU*, +G) with the empty
246 vector compared to the complemented strains while isolate A (*manB*, $\Delta 11$) with the empty vector
247 exhibited decreased growth compared to the complemented strain in 1 of the 3 experiments (p
248 = 0.0585) (Fig. 6). This result suggests that not only do these mutations confer resistance to
249 ICP phage infection, they also negatively impact growth of *V. cholerae* in the conditions
250 examined here.

251

252 **A**




B



253

254

C



255

256 **Figure 6: Complementation of mutated genes restores susceptibility phenotype.**

257 Strains with inducible plasmids containing the *manB* gene (*CmanB*), the *wbeU* gene (*CwbeU*),
258 or the empty vector (EV) control were challenged with ICP1 (A), ICP2 (B), and ICP3 (C) at 0
259 hours. Results are shown as the mean of three replicates with error bars as the standard
260 deviation. Two-Way Repeated Measures ANOVA was performed with Tukey's post-hoc multiple
261 comparisons tests between each strain. Significance is only shown at 0 hr and the final time

262 point. Statistical significance is noted as follows: asterisks (*) indicate significance between
263 *manB*Δ11bp/EV and *manB/CmanB*, pound signs (#) indicate significance between
264 *wbeUins*1bp/EV and *wbeUins*1/*CwbeU*. Levels of significance are denoted by: (ns) not
265 significant, (*) $p < 0.05$, (**) $p < 0.01$, (***) $p < 0.001$, and (****) $p < 0.0001$.

266


267

268 **The ICP resistant mutants exhibit variable sensitivity to polymyxin B**

269 In Gram-negative bacteria, the O antigen provides an outermost protective layer to various
270 stresses and antimicrobials (18). Previous O1 biosynthesis mutants that were resistant to ICP1
271 infection exhibited increased sensitivity to the antimicrobial polymyxin B (14). We therefore
272 performed a minimum inhibitory concentration assay to test if the seven O1 antigen ICP
273 resistant mutants were likewise sensitive to polymyxin B (Fig. 7, Table 2). Indeed, all seven
274 mutants exhibited increased sensitivity to polymyxin B relative to the WT strain, but only isolates
275 K and O were significantly different from the WT strain (Table 2). Isolate M, which was less
276 resistant to ICP infections in EOP assays, was the most like the WT strain with complete
277 resistance to polymyxin B at 8 μg/mL whereas isolates K and O remained completely sensitive
278 at this concentration. These results provide further evidence that these mutations disrupt O1
279 antigen biosynthesis although to varying degrees depending on the specific mutation.

280

281



282

283 **Figure 7: Mutant susceptibility to the antibiotic polymyxin B.**

284 Lines of best fit were calculated using non-linear regression models to determine IC₅₀. Each
285 condition includes three biological replicates, and error bars depict the standard deviation.

286

287 **Table 2: Half-Maximal Inhibitory Concentrations of Phage Resistant Mutants to**

288 **Polymyxin B**

Isolate	IC ₅₀ (95% Confidence Interval) (µg/mL)
WT	19.7 (10.8 – 35.8)
A	10.6 (4.18 – 25.8)
J	6.42 (3.07 – 13.4)
K	3.26 (1.82 – 5.78)
L	8.95 (4.65 – 17.0)
M	15.5 (8.41 – 28.3)
N	13.1 (4.81 – 39.0)
O	4.08 (2.08 – 7.96)


289

290 Discussion

291 Our finding that *V. cholerae* cultures rebound after ICP3 infection through enrichment of
292 escape mutants is consistent with the results of previous studies (3, 16, 26). Interestingly, all of
293 these resistant mutants had defects in genes coding for enzymes that synthesize the O1
294 antigen. A putative pathway for the biosynthesis of the O1 antigen has been proposed by
295 Stroehler *et al*, 1995 (27) with modifications by Chatterjee & Chaudhuri, 2004 (25) and Seed *et*
296 *al*, 2012 (14). Three of our mutant strains in Table 1 had alterations to genes encoding
297 perosamine synthesis enzymes. These include a frameshift deletion in the
298 phosphomannomutase *manB* (Isolate A), a missense mutation in the GDP-mannose
299 dehydratase *gmd* (Isolate M), and a frameshift deletion in the mannose-1 phosphate
300 guanylyltransferase *manC* (Isolate N) (Table 1). Mutations that alter the function of these
301 proteins prevent synthesis of the perosamine monomer in O1 antigen from being produced.
302 *manB* mutations have been linked with *V. cholerae* ICP1 resistance, but not with resistance to
303 ICP2 or ICP3 (26). Likewise, two mutations, one in-frame deletion (Isolate K) and one frameshift
304 deletion (Isolate L), in the *wbeN* acyl-CoA reductase gene conferred pan-resistance to ICP
305 phages (Table 1). The final isolates had a mutation in *wbeU* that is associated with LPS
306 biosynthesis and *wbeM*, whose function in O1 antigen biosynthesis is unclear.

307 Given that the receptor for ICP1 has been previously identified as the O antigen, it is not
308 surprising to see resistance among the isolates generated in this study. However, the
309 requirement of the O1 antigen for infection by ICP2 has not been reported. OmpU was
310 previously identified as a receptor for ICP2 (16), and it was recently determined that the ICP2
311 tail protein Gp23 interacts with OmpU (28, Fig. 8). OmpU was identified as the receptor of ICP2
312 from clinical stool samples by analyzing *V. cholerae* resistant to ICP2. These resistant mutants
313 had mutations in *ompU* or *toxR*, a transcriptional activator of *ompU* (16). We speculate that
314 mutations in the O1 antigen biosynthesis genes were not isolated in this study due to the
315 significant *in vivo* cost for *V. cholerae* associated with the loss of O1 (14). Having multiple

316 receptors is not uncommon for phages, sometimes requiring both to be present for adsorption
317 into the cell. This is exhibited by the *E. coli* phage K20 which utilizes LPS in addition to OmpF
318 for infection perpetuation (29).
319



320

321 **Figure 8: Model showing the known receptors of ICP phage.**

322 This model shows a WT strain of *V. cholerae* on the bottom and a strain with a mutation in the
323 O1 antigen biosynthesis pathway on top. The WT strain is able to produce both the O1 antigen
324 and the Outer Membrane Porin U (OmpU) while the mutant can only produce the later. ICP1 is
325 depicted as larger and a member of the *Myoviridae* family. ICP2 and ICP3 are depicted as
326 smaller and members of the *Podoviridae* family. The three ICP phages and their previously
327 known targets are shown as dark green arrows. The new targets identified by this study are
328 shown by light green arrows.

329

330 This study is the first to report the O1 antigen is a receptor for ICP3 (Fig. 8). A study
331 exploring VP4 phage which is genetically similar to ICP3 described its receptor in this same
332 locus (13, 15). This study found mutations in *manB*, *manC*, *wbeU*, *wbeN*, and *wbeG*, which
333 reinforce our findings here. Our isolation of seven independent mutations in O1 antigen
334 biosynthesis and no other putative receptors may suggest that the O1 antigen is the sole
335 receptor for ICP3. However, given that the target region for disruption of O1 antigen
336 biosynthesis is large (Fig. 3), further studies with more saturating mutagenesis are required to
337 conclusively test this hypothesis.

338 The results here raise an important question as to why these O1 antigen deficient *V.*
339 *cholerae* lineages have not dominated environmentally since they provide pan-resistant to ICP1,
340 ICP2, and ICP3 infection. Since LPS operates as a virulence factor within the human host, it is
341 important for survival and replication in that niche (12). LPS mutants exhibit decreased
342 competitiveness and ability to colonize the human gut compared to their wild type counterparts
343 (30). There is additional evidence that the LPS protects against acid exposure which may be
344 typical during human colonization (30). Furthermore, in our own studies, we observed
345 decreased growth in O1 antigen biosynthesis mutants compared to their complemented
346 counterparts in laboratory growth conditions and increased sensitivity to polymyxin B although it
347 was only statistically significant in two mutants (Figs. 6 and 7). These collective results suggest
348 that emergence of O1 antigen null mutants is not evolutionary favored due to the significant
349 fitness cost of such mutations. This finding is consistent with previous studies that have similarly
350 concluded that although isolation of phage receptor mutants is common during selection for
351 phage resistance in laboratory conditions, such mutants are rarely seen in natural populations
352 (31).

353 The E7946 El Tor strain encodes homopolymer nucleotide tracts in the genes *wbeL* and
354 *manA*, which are required for O1 antigen biosynthesis, that mediate phase variable expression

355 of the O1 antigen leading to ICP1 phage resistance (14). It was postulated that such a system
356 could lead to a strategy in which a segment of the population evolves resistance to ICP1 phage
357 infection while maintaining the genetic potential to restore O1 for infection of the host. We did
358 not isolate any mutations in *manA* or *wbeL*, and an analysis of the sequence of genes in the El
359 Tor strain used here, C6706, showed that the polyA(A₉) tracts were conserved in *manA* but not
360 in *wbeL*. This suggests that the loss of *manA* is not favored upon ICP3 selection.

361 As phages use cell surface proteins and molecules as receptors, an emerging concept is
362 that resistance to phage infection may have the evolutionary tradeoff of antibiotic sensitivity.
363 Such a tradeoff has been demonstrated in the *E. coli* phage U136B which uses the antibiotic
364 efflux protein TolC as a receptor, as mutations that confer resistant to phage infection lead to
365 increased antibiotic sensitivity (32). Evolving to recognize the O1 antigen as a receptor by all
366 three ICP phage is a natural example of such a concept as any *V. cholerae* that gain resistance
367 through loss of O1 have the tradeoff of loss of virulence and more sensitivity to host
368 antimicrobials like polymyxin B. Therefore, evolution to use the O1 antigen as a receptor by
369 ICP1, ICP2, and ICP3 has contributed to the evolutionary success of these lytic phage.

370

371 **ACKNOWLEDGEMENTS**

372 This research was supported and funded by NIH grants GM139537 and AI158433 to
373 C.M.W. D.A.B. was supported by a Professorial Assistant fellowship from Michigan State
374 University and the Norman E. & Hanna C. Kelker Research Scholarship and the Dietrich C.
375 Bauer Scholarship from the Department of Microbiology and Molecular Genetics. We thank
376 Andrew Camilli and Wai-Leung Ng at Tufts University for providing the ICP phage.

377

378 **MATERIALS AND METHODS**

379 **Bacterial culturing and molecular biology**

380 The WT culture is El Tor *V. cholerae* C6706:str2 (33). All cultures were grown in glass test tubes

381 (18 x 150 mm, Kimax) containing Luria-Bertani broth at 35°C with shaking, unless otherwise
382 stated. Selection of plasmids utilized kanamycin at 100 µg/mL. The *manB* (pDAB1) and *wbeU*
383 (pDAB2) complementation plasmids were generated using Gibson cloning. The gene inserts
384 were generated from C6706:str2 with GoTaq 2x MasterMix using the primers (overhang, RBS,
385 INSERT): 5'-acagcctcgacaggcctagggaggagctaaggaagctaaaGTGAAAGAGTTAACTTGTTTT-3'
386 and 5'-gcttgctcaatcaatcaccgTTAAATATCCAATTTCTTAATTAATTTAGTAAG-3' (*manB*) 5'-
387 acagcctcgacaggcctagggaggagctaaggaagctaaaATGCCATGGAAGACCTAC-3' and 5'-
388 gcttgctcaatcaatcaccgTCAACAGACATTTCCGAAG3' (*wbeU*). These inserts were Gibson (New
389 England Biolabs) cloned into an EcoRI/BamHI digested pEVS143 plasmid (34) using standard
390 methodologies followed by Sanger sequencing to confirm the proper clone was generated. The
391 empty vector and plasmids were conjugated from the *Escherichia coli* diaminopimelic acid
392 auxotrophic strain BW29427 (K. Datsenko and B.L. Wanner, unpublished) into *V. cholerae*
393 strains by mixing 25 µL of each culture overnight on an agar plate followed by selection on LB
394 plates with kanamycin.

395

396 **Liquid Infection Assays**

397 30 µL of overnight cultures were back diluted into 3 mL of LB and allowed to grow until
398 OD₆₀₀ (0.20-0.50). Each culture was then infected with ICP1, ICP2, ICP3, or SM buffer (50 mM
399 Tris-Cl + 100 mM NaCl + 8 mM MgSO₄). Phage stocks were maintained in filtered LB medium
400 with trace chloroform and the pfu/mL of each phage was: ICP1 9.2x10⁵ pfu/mL, ICP2 1.3x10⁸
401 pfu/mL, and ICP3 1.8x10⁹ pfu/mL. The amount of phage to add was determined by converting
402 the optical density (OD) of the culture into an estimated viable cell count in *cfu/mL* using a
403 quadratic regression equation generated from a previous growth curve: $cfu/mL = [1.4226 \times 10^9 \times$
404 $(OD)^2] + [4.2981 \times 10^8 \times OD] - 2.2521 \times 10^6$. Following infection, the OD₆₀₀ of cultures was
405 remeasured every 0.25 hr. For the recovery experiments, cultures were allowed to recover for
406 24 hours. The recovery pools were collected and frozen in 20% glycerol at -80°C. The recovery

407 pools were then struck onto LB agar plates and individual isolates were used for reinfection. The
408 same infection procedure was followed for the reinfection assays. For the liquid phage infection
409 assays in Fig. 4, three independent overnight cultures of the WT or seven phage resistant
410 isolates grown in 2 mL of LB were back diluted 1/100 in fresh LB and 100 μ L was added to four
411 wells into a clear 96-well plate (Costar). The plate was incubated at 35°C for 1.5 hours and to
412 one well 1 μ L of the ICP1, ICP2, or ICP3 phage stock was added. The OD₅₉₅ was monitored
413 every 0.25 hours in an Envision plate reader.

414

415 **Efficiency of Plaquing Assays**

416 Cultures of WT and resistant strains were grown overnight in LB and 250 μ L was
417 inoculated into 18 mL warm MMB (LB + 0.5% agar + 5 mM MgCl₂ + 5 mM CaCl₂ + 0.1 mM
418 MnCl₂) at 55°C. The mixture was poured onto an empty plate (150 x 15 mm, Fisher Scientific)
419 and allowed to solidify. Ten-fold serial dilutions of phage made in SM buffer were then spotted 5
420 μ L in succession onto each strain along with one SM buffer spot as a negative control. This was
421 performed in triplicate for each strain. Once plates were dried, they were allowed to incubate at
422 35°C overnight. Plaques were counted the next day to determine the efficiency of plaquing on
423 each strain for each phage (ICP1, ICP2, and ICP3).

424

425 **Whole Genome Sequencing and Analysis**

426 Genomic extraction for the WT C6706:str2 strain and each of the resistant mutants was
427 performed using Wizard Genomic DNA Purification Kit (Promega). The concentration of DNA
428 was quantified using Nanodrop spectrophotometer (ND-1000). Samples were then sequenced
429 at SeqCenter (Pittsburgh, PA). Results were processed by aligning each sample to the
430 reference chromosomes of *V. cholerae* (NZ_CP046844 and NZ_CP046845) using *breseq* (35).
431 Any mutations found in the WT strain as well as our resistant mutants were removed from
432 consideration.

433

434 **Polymyxin B minimum inhibitory concentration**

435 Three independent overnight cultures of the WT and all seven resistant mutants were prepared
436 in 2 mL LB and diluted 1/100 in fresh media. A final volume of 150 μ L of the diluted culture was
437 added to the well of a clear 96-well plate (Costar) and grown overnight at 35°C without shaking
438 in a humidity chamber with decreasing concentrations of polymyxin B starting at 555 mg/mL
439 followed by seven four-fold dilutions. The OD₅₉₅ was measured using an Envision plate reader.

440

441 **REFERENCES**

- 442 1. Dalusi L, Lyimo TJ, Lugomela C, Hosea KMM, Sjöling S. 2015. Toxigenic *Vibrio cholerae*
443 identified in estuaries of Tanzania using PCR techniques. *FEMS Microbiol Lett* 362:fnv009.
- 444 2. Ali M, Nelson AR, Lopez AL, Sack DA. 2015. Updated Global Burden of Cholera in Endemic
445 Countries. *PLoS Negl Trop Dis* 9:e0003832.
- 446 3. Yen M, Cairns LS, Camilli A. 2017. A cocktail of three virulent bacteriophages prevents
447 *Vibrio cholerae* infection in animal models. *Nat Commun* 8:14187.
- 448 4. Kanungo S, Azman AS, Ramamurthy T, Deen J, Dutta S. 2022. Cholera. *The Lancet*
449 399:1429–1440.
- 450 5. Ganesan D, Gupta SS, Legros D. 2020. Cholera surveillance and estimation of burden of
451 cholera. *Vaccine* 38:A13–A17.
- 452 6. Yen M, Camilli A. 2017. Mechanisms of the evolutionary arms race between *Vibrio cholerae*
453 and Vibriophage clinical isolates. *Int Microbiol* 20:116–120.
- 454 7. Severin GB, Ramliden MS, Hawver LA, Wang K, Pell ME, Kieninger A-K, Khataoakar A,
455 O’Hara BJ, Behrmann LV, Neiditch MB, Benning C, Waters CM, Ng W-L. 2018. Direct
456 activation of a phospholipase by cyclic GMP-AMP in El Tor *Vibrio cholerae*. *Proc Natl Acad*
457 *Sci USA* 115:E6048-E6055.

- 458 8. O'Hara BJ, Barth ZK, McKitterick AC, Seed KD. 2017. A highly specific phage defense
459 system is a conserved feature of the *Vibrio cholerae* mobilome. *PLoS Genet* 13:e1006838.
- 460 9. Hsueh BY, Severin GB, Elg CA, Waldron EJ, Kant A, Wessel AJ, Dover JA, Rhoades CR,
461 Ridenhour BJ, Parent KN, Neiditch MB, Ravi J, Top EM, Waters CM. 2022. Phage defence
462 by deaminase-mediated depletion of deoxynucleotides in bacteria. *Nat Microbiol* 7:1210–
463 1220.
- 464 10. Seed KD, Lazinski DW, Calderwood SB, Camilli A. 2013. A bacteriophage encodes its own
465 CRISPR/Cas adaptive response to evade host innate immunity. *Nature* 494:489–491.
- 466 11. Boyd CM, Angermeyer A, Hays SG, Barth ZK, Patel KM, Seed KD. 2021. Bacteriophage
467 ICP1: A Persistent Predator of *Vibrio cholerae*. *Annu Rev Virol* 8:285–304.
- 468 12. Seed KD, Bodi KL, Kropinski AM, Ackermann H-W, Calderwood SB, Qadri F, Camilli A.
469 2011. Evidence of a Dominant Lineage of *Vibrio cholerae*-Specific Lytic Bacteriophages
470 Shed by Cholera Patients over a 10-Year Period in Dhaka, Bangladesh. *mBio* 2:e00334-10.
- 471 13. Barman RK, Chakrabarti AK, Dutta S. 2022. Screening of Potential *Vibrio cholerae*
472 Bacteriophages for Cholera Therapy: A Comparative Genomic Approach. *Front Microbiol*
473 13:803933.
- 474 14. Seed KD, Faruque SM, Mekalanos JJ, Calderwood SB, Qadri F, Camilli A. 2012. Phase
475 Variable O Antigen Biosynthetic Genes Control Expression of the Major Protective Antigen
476 and Bacteriophage Receptor in *Vibrio cholerae* O1. *PLoS Pathog* 8:e1002917.
- 477 15. Xu J, Zhang J, Lu X, Liang W, Zhang L, Kan B. 2013. O Antigen Is the Receptor of *Vibrio*
478 *cholerae* Serogroup O1 El Tor Typing Phage VP4. *J Bacteriol* 195:798–806.
- 479 16. Seed KD, Yen M, Shapiro BJ, Hilaire IJ, Charles RC, Teng JE, Ivers LC, Boncy J, Harris JB,
480 Camilli A. 2014. Evolutionary consequences of intra-patient phage predation on microbial
481 populations. *eLife* 3:e03497.

- 482 17. Reyes-Robles T, Dillard RS, Cairns LS, Silva-Valenzuela CA, Housman M, Ali A, Wright ER,
483 Camilli A. 2018. *Vibrio cholerae* Outer Membrane Vesicles Inhibit Bacteriophage Infection. *J*
484 *Bacteriol* 200:e00792-17.
- 485 18. Chatterjee SN, Chaudhuri K. 2006. Lipopolysaccharides of *Vibrio cholerae*: III. Biological
486 functions. *Biochim Biophys Acta* 1762:1–16.
- 487 19. Mathur J, Waldor MK. 2004. The *Vibrio cholerae* ToxR-Regulated Porin OmpU Confers
488 Resistance to Antimicrobial Peptides. *Infect Immun* 72:3577–3583.
- 489 20. Gupta P, Modgil V, Kant V, Kaur H, Narayan C, Mahindroo J, Verma R, Mohan B, Taneja N.
490 2022. Phenotypic and genotypic characterization of antimicrobial resistance in clinical
491 isolates of *Vibrio cholerae* over a decade (2002-2016). *Indian J Med Microbiol* 40:24–29.
- 492 21. Miwanda B, Moore S, Muyembe J-J, Nguéfack-Tsague G, Kabangwa IK, Ndjakani DY,
493 Mutreja A, Thomson N, Thefenne H, Garnotel E, Tshapenda G, Kakongo DK, Kalambayi G,
494 Piarroux R. 2015. Antimicrobial Drug Resistance of *Vibrio cholerae*, Democratic Republic of
495 the Congo. *Emerg Infect Dis* 21:847–851.
- 496 22. Rodriguez-Gonzalez RA, Leung CY, Chan BK, Turner PE, Weitz JS. 2020. Quantitative
497 Models of Phage-Antibiotic Combination Therapy. *mSystems* 5:e00756-19.
- 498 23. Hsueh BY, Waters CM. 2019. Combating Cholera. *F1000research* 8:589.
- 499 24. Bhandare S, Colom J, Baig A, Ritchie JM, Bukhari H, Shah MA, Sarkar BL, Su J, Wren B,
500 Barrow P, Atterbury RJ. 2019. Reviving Phage Therapy for the Treatment of Cholera. *The J*
501 *Infect Dis* 219:786–794.
- 502 25. Chatterjee SN, Chaudhuri K. 2004. Lipopolysaccharides of *Vibrio cholerae*. *Biochim Biophys*
503 *Acta* 1690:93–109.
- 504 26. Silva-Valenzuela CA, Camilli A. 2019. Niche adaptation limits bacteriophage predation of
505 *Vibrio cholerae* in a nutrient-poor aquatic environment. *Proc Natl Acad Sci USA* 116:1627–
506 1632.

- 507 27. Stroehler UH, Karageorgos LE, Brown MH, Morona R, Manning PA. 1995. A putative
508 pathway for perosamine biosynthesis is the first function encoded within the *rfb* region of
509 *Vibrio cholerae* O1. *Gene* 166:33-42.
- 510 28. Lim ANW, Yen M, Seed KD, Lazinski DW, Camilli A. 2021. A Tail Fiber Protein and a
511 Receptor-Binding Protein Mediate ICP2 Bacteriophage Interactions with *Vibrio cholerae*
512 OmpU. *J Bacteriol* 203:e00141-21.
- 513 29. Silverman JA, Benson SA. 1987. Bacteriophage K20 requires both the OmpF porin and
514 lipopolysaccharide for receptor function. *J Bacteriol* 169:4830–4833.
- 515 30. Nesper J, Lauriano CM, Klose KE, Kapfhammer D, Kraiß A, Reidl J. 2001. Characterization
516 of *Vibrio cholerae* O1 El Tor *galU* and *galE* Mutants: Influence on Lipopolysaccharide
517 Structure, Colonization, and Biofilm Formation. *Infect Immun* 69:435–445.
- 518 31. Mutalik VK, Adler BA, Rishi HS, Piya D, Zhong C, Koskella B, Kutter EM, Calendar R,
519 Novichkov PS, Price MN, Deutschbauer AM, Arkin AP. 2020. High-throughput mapping of
520 the phage resistance landscape in *E. coli*. *PLoS Biol* 18:e3000877.
- 521 32. Burmeister AR, Fortier A, Roush C, Lessing AJ, Bender RG, Barahman R, Grant R, Chan
522 BK, Turner PE. 2020. Pleiotropy complicates a trade-off between phage resistance and
523 antibiotic resistance. *Proc Natl Acad Sci USA* 117:11207–11216.
- 524 33. Thelin KH, Taylor RK. 1996. Toxin-coregulated pilus, but not mannose-sensitive
525 hemagglutinin, is required for colonization by *Vibrio cholerae* O1 El Tor biotype and O139
526 strains. *Infect Immun* 64:2853–2856.
- 527 34. Dunn AK, Millikan DS, Adin DM, Bose JL, Stabb EV. 2006. New *rfp* - and pES213-Derived
528 Tools for Analyzing Symbiotic *Vibrio fischeri* Reveal Patterns of Infection and *lux* Expression
529 In Situ. *Appl Environ Microbiol* 72:802–810.
- 530 35. Deatherage DE, Barrick JE. 2014. Identification of Mutations in Laboratory-Evolved
531 Microbes from Next-Generation Sequencing Data Using *breseq*. *Methods Mol Biol*
532 1151:165-188.

- 533 36. Fallarino A, Mavrangelos C, Stroehler UH, Manning PA. 1997. Identification of additional
534 genes required for O-antigen biosynthesis in *Vibrio cholerae* O1. *J Bacteriol* 179:2147–
535 2153.
- 536 37. Pombo JP, Ebenberger SP, Müller AM, Wolinski H, Schild S. 2022. Impact of Gene
537 Repression on Biofilm Formation of *Vibrio cholerae*. *Front Microbiol* 13:912297.
538