

CONCISE REVIEW

Alternative splicing in mesenchymal stem cell differentiation

Jung Woo Park  | Siyi Fu | Borong Huang | Ren-He Xu

Center for Reproduction, Development, and Aging and Institute of Translational Medicine, Faculty of Health Sciences, University of Macau, Macau, People's Republic of China

Correspondence

Jung Woo Park, PhD, Center for Reproduction, Development, and Aging and Institute of Translational Medicine, Faculty of Health Sciences, University of Macau, Macau, People's Republic of China.
Email: jungwupark@um.edu.mo

Funding information

Macau Science and Technology Development Fund (FDCT), Grant/Award Numbers: #028/2015/A1, #095/2017/A1; University of Macau Research Committee funds (MYRG), Grant/Award Numbers: #2015-00169-FHS, #2016-00070-FHS, #2017-00124-FHS, #2017-00249-FHS

Abstract

The differentiation and maturation of mesenchymal stem cells (MSCs) to mesodermal and other lineages are known to be controlled by various extrinsic and intrinsic signals. The dysregulation of the MSC differentiation balance has been linked to several pathophysiological conditions, including obesity and osteoporosis. Previous research of the molecular mechanisms governing MSC differentiation has mostly focused on transcriptional regulation. However, recent findings are revealing the underrated role of alternative splicing (AS) in MSC differentiation and functions. In this review, we discuss recent progress in elucidating the regulatory roles of AS in MSC differentiation. We catalogue and highlight the key AS events that modulate MSC differentiation to major osteocytes, chondrocytes, and adipocytes, and discuss the regulatory mechanisms by which AS is regulated.

KEYWORDS

adipogenic, alternative splicing, chondrogenic, ESC differentiation, mesenchymal stem cells, MSC differentiation, neural differentiation, osteogenic, RNA-binding proteins

1 | INTRODUCTION

Mesenchymal stem or stromal cells (MSCs) are multipotent and multifunctional stem cells that have the potential to differentiate into mesodermal cell types, including osteoblasts, chondrocytes, and adipocytes, and they also play important roles as immune regulators and a critical niche in hematopoiesis.^{1,2} Due to the current debate surrounding their embryonic origin and in vivo functions, the term “MSC” has been suggested to be inaccurate as MSCs from different tissue depots clearly have different phenotypes. For example, not all MSCs support hematopoiesis. Thus, alternative terms such as “multipotent mesenchymal stromal cells” and “tissue specific stem cells” have been proposed. Given the broad therapeutic effects identified thus far for MSCs derived from various adult tissues and embryonic stem cells, it has even been suggested that MSCs be referred to as “medicinal signaling cells.”³⁻⁵ In this review, however, we use “MSC” throughout, for the simplicity of alluding to classically defined differentiation potential toward mesodermal derivatives in vivo and in vitro.^{6,7}

Dissecting the governing molecular mechanisms of MSC differentiation into mesenchymal cell types is critical for understanding MSC

function in vivo and evaluating and predicting the clinical outcome of MSC-based cell therapy. The balance of extrinsic signaling pathways through combinatorial or distinct sets of cytokines, growth factors, and hormones coordinates the activation or suppression of transcription factors that determine the fate of MSCs and the maturation of progenitors into terminally differentiated cell types.⁸⁻¹⁰ One of the best known examples is Wnt signaling, which simultaneously activates and represses the expression of Runt-related transcription factor 2 (*RUNX2*) and peroxisome proliferator-activated receptor gamma (*PPAR γ*), the master genes for osteogenesis and adipogenesis, respectively, and plays a critical role in lineage commitment during early MSC differentiation.¹¹

The importance of alternative splicing (AS), which will be discussed in more detail below, in controlling MSC differentiation has gained less attention than the mechanisms of transcriptional control. A recent study by Aprile et al identified a novel alternative isoform of *PPAR γ* , which modulates adipogenesis by blocking the function of wild type *PPAR γ* in two ways.¹² Interestingly, the expression level of the *PPAR γ Δ 5* isoform in subcutaneous adipose tissues was shown to be directly proportional to body mass index (BMI) in two independent

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

©2020 The Authors. *STEM CELLS* published by Wiley Periodicals LLC on behalf of AlphaMed Press 2020

cohorts of obese or type II diabetic patients, highlighting a potential link between AS and health risk of obesity.

Van de Peppel et al analyzed the gene expression dynamics for the early steps of human MSC (hMSC) differentiation into osteoblasts and adipocytes with a high-density temporal resolution.¹³ Upstream regulator analysis revealed 34 transcription factors (TFs) with a role in the early differentiation of MSC. Unexpectedly, the expression change of the differentially expressed genes (DEGs) that is thought to be regulated by the 34 TFs did not match with the expression change of the TFs themselves during the early stage of MSC differentiation.¹³ This suggests that additional gene expression regulatory mechanisms, including post-transcriptional regulation, may be needed to explain the aforementioned discrepancy in evidence for the control the expression of the MSC differentiation-associated genes.

Moreover, recent studies have demonstrated that pre-mRNAs encoding master transcription factors, the prerequisite for the early lineage specification of MSC differentiation, undergoes AS and encode multiple protein isoforms with functionally distinct activities. The pre-mRNAs of the osteogenic and adipogenic master regulators, *RUNX2* and *PPAR γ* , generate multiple mRNA isoforms by AS that are subsequently translated into the protein isoforms with the regulatory activities.^{12,14}

Girardot et al recently demonstrated that a master regulator for chondrogenesis, *Sox9*, regulates the splicing of hundreds of genes without affecting their transcription during chondrogenic differentiation.¹⁵ The study indicates that the regulation occurs through the direct association of *Sox9* with pre-mRNAs and other RNA-binding proteins (RBPs), rather than by altering the target RBP expression levels and indirectly modulating AS. Consistent with the notion, more than one-third of the regulators that control ES- and neural-specific AS were transcription factors, implying that some transcription factors have dual functions, regulating transcription and AS during ES and neural differentiation.¹⁶ One of the surprising aspects of the study was that the regulatory transcription factors control AS and shape the cell fate-specificity by directly binding to target pre-mRNA.¹⁶ Not surprisingly, the coupling between pre-mRNA splicing and transcription has been well-documented as transcription elongation rate controls AS, and AS controls the transcriptional rate.¹⁷

Yet, there is a lack of systematic review on the regulatory roles of AS in MSC differentiation despite the current research progress in the field. In this review, we will briefly discuss the regulatory mechanisms of AS, focusing on the dynamic interactions between cis-acting sequence elements within the pre-mRNA and trans-acting RBPs, and discuss the major AS of pre-mRNAs encoding the key transcription factors and critical factors that are involved in the cell fate determination, maturation, and metabolic status for MSC differentiation into osteocytes, chondrocytes, adipocytes, and neurons.

2 | AS AND ITS REGULATORY MECHANISMS

AS is one of the powerful regulatory mechanisms of gene expression by which a pre-mRNA generates multiple mRNA and protein isoforms

Significance statement

Mesenchymal stem cells (MSCs) are important cell sources for bone, fat tissue, cartilage, and other connective tissues for both in vivo and in vitro studies. A better understanding of the molecular mechanisms involved in the intrinsic regulatory network that governs the differentiation of MSCs is needed. In this review, the underrated role of alternative splicing in MSC differentiation, a process that produces multiple similar but functionally distinct proteins from one gene, is discussed.

to increase the functional capacity of cells. As high as 95% of all human genes undergo AS and the pattern of AS is highly variable among different human tissues, suggesting a correlation between the unique signature of the transcriptome, regulated by AS, and tissue-specific structure and function.¹⁸ Thanks to the recent development of high-throughput RNA sequencing (RNA-Seq), more widespread, evolutionarily conserved, overall biological roles of AS in the gene expression landscape have been recently recognized.¹⁹⁻²¹

In metazoans, introns are removed from pre-messenger RNAs (pre-mRNAs) to produce mature RNA. However, this seemingly simple chemical reaction is catalyzed by the largest known macromolecular complex inside the cell, called the spliceosome. This mega-sized machinery is made of more than 300 RNA and protein components.^{22,23} The regulation of AS is controlled by the interaction between cis-acting elements and trans-acting factors, primarily RBPs. Trans-acting factors interact with different components of the spliceosome during the assembly of the spliceosome on the nascent pre-mRNA. Here, we summarize the general mechanism of regulation by which several groups of RBPs (heteronuclear ribonucleoproteins [hnRNPs], serine-arginine rich proteins [SRs], and KH domain-containing protein such as NOVA) function by promoting or inhibiting the binding of components of the splicing machinery to the target splice sites (Figure 1). These auxiliary splicing factors recognize a variety of cis-acting sequence elements within the pre-mRNA and are known as exonic splicing enhancers, exonic splicing silencers, intronic splicing enhancers, and intronic splicing silencers (ISSs), by either recruiting or blocking the assembly of small nuclear ribonucleoprotein (snRNP) complexes (Figure 1). These RBPs are expressed in a tissue- and developmental-specific manner and the relative expression level of critical RBPs in a specific cell determines a ratio of the isoforms by regulating whether or not a given exon will be included in the mRNA.

3 | REGULATION OF STEM CELL FATE DETERMINATION BY AS

One of the earliest evidence that AS plays a critical role in the pluripotency-differentiation axis in ESCs came from a study in which a shorter FGF4 isoform, FGF4si, attenuated activity of FGF4 by a novel fibroblast growth factor 4 (FGF4), thereby inducing ESC




FIGURE 1 A schematic of the regulatory mechanisms by which interaction between cis-acting elements and trans-acting factors control the inclusion or exclusion of the cassette exon in three exons model. A, Serine-arginine (SR) proteins (green) bind to exonic splicing enhancer (ESE) and recruit U1 and U2 snRNP for the inclusion of exon 2. B, RNA binding protein, NOVA, binds to intronic splicing enhancer (ISE) and recruit U1 and U2 snRNP for the inclusion of exon 2. C, RNA binding protein, hnRNP, binds to exonic splicing silencer (ESS) to block the binding of U1 and U2 snRNP for the exclusion of exon 2. D, RNA binding protein, hnRNP, binds to intronic splicing silencer (ISS) to block the binding of U1 and U2 snRNP for the exclusion of exon 2

differentiation.²⁴ Later, Blencowe's group demonstrated that the muscleblind-like RNA binding proteins-mediated alternative splicing of a key transcription factor, FOXP1, makes a significant contribution to the switch between maintaining pluripotency and triggering differentiation of human ESCs.^{25,26} Moreover, Paired box 6 (PAX6), RNA binding motif protein 24 (RBM24)-mediated, Tcf3, and Sal4 AS have been associated with ectodermal, mesodermal, and endodermal lineage determination of human and mouse embryonic stem cells.²⁷⁻³¹

Su et al recently reported that RNA binding domain protein 4 (RBM4) regulates AS of PKM pre-mRNA, thereby promoting neural differentiation of hMSCs.³² RBM4 was originally shown to promote the differentiation of neuronal progenitor cells and the neurite outgrowth of cultured neurons via its role in splicing regulation. During neuronal differentiation, energy production shifts from glycolysis to oxidative phosphorylation. AS of Pyruvate kinase M (PKM) has been implicated as a fuel switch in determining the source of energy

production, glycolysis or oxidative phosphorylation. Inclusion of mutually exclusive exons, 9 and 10, results in two PKM isoforms, PKM1 and PKM2, differing in the formation of the tetramer, which is a limiting factor for the production of pyruvate. Tetramer formation is less efficient for PKM2, which results in the reduced production of pyruvate, substrates for oxidative phosphorylation, and in turn, promotes the production of energy through glycolysis. Indeed, a higher PKM2/PKM1 ratio is a hallmark of undifferentiated stem cells and cancer cells.³³ PKM pre-mRNA alternative splicing appears to be regulated by multiple RBPs including hnRNP A1 and A2, polypyrimidine-binding protein (PTB), RBM4, and SRSF3 by binding to an ISS to block the assembly or entry of U1 snRNP to the 5' splice site, thus preferentially selecting exon 10 for the production of the PKM2 isoform.³⁴⁻³⁶

Genome-wide RNA-seq analyses have revealed that a specific AS pattern is observed with ESCs and a global AS change in differentiated ESCs. Conversely, the global AS profile in differentiated cells reverts back to that of a pluripotent state during stem cell reprogramming. Although this finding does not provide the conclusion that AS program is the driving force of differentiation, it supports the notion that a signature pattern of AS for a particular cell type may contribute to the cell type-specific global gene expression landscape. More recently, it has been shown that the downregulation of U2AF, a core component of spliceosome, induces a switch in the AS of TFs involved in the differentiation of pluripotent stem cells and controlled cell-fate specific AS.³⁷ Besides, it has been shown that differential AS events were detected in MSCs derived from young and old donors, illustrating that AS is a widespread mechanism to shape the transcriptome landscape in various cell and tissue types, as well as during differentiation.^{38,39}

The regulation of MSC differentiation by AS has recently been recognized as an additional mechanism that controls the precise cell fate determination and maturation to a terminally differentiated cell type. One mechanism of AS in cell fate determination is the production of two functionally distinct or opposite isoforms by AS, forming a molecular switch during cell fate commitment.^{40,41} As mentioned above, pre-mRNAs of key transcription factors, which specify cell fate by controlling the expression of downstream target genes, undergo AS, resulting in alternative isoforms. These alternative isoforms play overlapping, distinct, and even opposite roles during MSC differentiation through various molecular mechanisms. Here, we describe the AS of key molecules that specify cell fate during early MSC differentiation and maturation steps (summarized in Table 1) and the major AS events associated with MSC differentiation (illustrated in Figure 2).

4 | REGULATION OF OSTEOGENIC DIFFERENTIATION BY AS

The activation of transcription factor RUNX2 is a prerequisite for the commitment to an early osteogenic lineage and is switched off later during osteocyte maturation.⁶⁵ The induction of RUNX2 in MSCs coincides with the commitment to osteogenic progenitors, and it also suppresses the cell fate specification toward an adipogenic lineage.¹⁴ The first hint that AS may be involved in regulating osteogenic

differentiation came from several studies that showed that a shorter RUNX2 protein isoform, produced by alternative splicing, exhibited distinct transcriptional activity.^{14,66} Makita et al demonstrated the detection of four RUNX2 mRNA isoforms; WT RUNX2, RUNX2 Δ 5, Δ 7, and Δ 5 Δ 7, through skipping one or both cassette exons (exon 5 and 7) in hMSCs.¹⁴ Two of these isoforms generated from the skipping of exon 5 encoded RUNX2 isoforms that were devoid of nuclear localization, thereby eliminating their transcriptional activities. On the other hand, a third mRNA isoform, RUNX2 Δ 7, encodes an isoform with a deletion in the region near the activation domain at the carboxy terminus of RUNX2. The RUNX2 Δ 7 protein isoform localizes in the nucleus and exhibits a similar binding affinity and transcriptional activity for the osteocalcin (OC) gene as WT RUNX2. However, RUNX2 Δ 7 is less dependent on coactivators and corepressors (CBP, p300, and HDAC3) in transactivating the OC gene. This raises the reasonable possibility that RUNX2 Δ 7 may transactivate both the same common target genes as the WT and distinct target genes in a CBP/p53- and HDAC3-independent manner.

Nonetheless, the expression pattern of two isoforms, WT and RUNX2 Δ 7, is distinct between chondrocytes and osteocytes. Roughly equal expression levels of the isoforms are present in chondrocyte differentiated from hMSCs, while exclusively the wild-type isoform is present in osteocyte differentiated from hMSCs, indicating that the WT/ RUNX2 Δ 7 ratio may be a key determinant for cell fate of hMSCs. Mechanistically, RUNX2 WT and Δ 7 possess distinct transcriptional activities for the target genes, including OC, and potentially provide the fine-tuning for osteogenic vs chondrogenic fate determination. It is also worth noting that the WT/ RUNX2 Δ 7 ratio varies among various osteoblast-like cells, including osteosarcomas, primary osteoblasts isolated from human donors, and differentiated osteocytes from hMSC, potentially the ratio of the two isoforms as a potential biomarker for cancers of osteocytic origin.

Although the regulation of RUNX2 alternative splicing has not been clearly elucidated, skipping RUNX2 alternative exons has been stimulated by knockdown (KD) of the components of U2 snRNP, namely, U2AF1, SF3A1, and SF3A3.⁶⁷ Interestingly, U2 snRNPs are components of the spliceosome and are considered essential for the catalysis of pre-mRNA splicing, rather than modulating AS. This finding is consistent with the previous studies that have shown that mutations in general splicing factors can indeed alter the global AS pattern in vitro and in vivo, rather than affecting global splicing defects in a cell- or tissue-specific manner.⁶⁸⁻⁷⁰ Furthermore, osteogenesis-specific RUNX2 target genes, including OSC, COL1A1, OPN, BSP, and OSX, undergo alternative splicing and produce multiple mRNA isoforms, although the exact biological implication is not known at present.⁶⁷

Of particular interest, OSX (Sp7) is an immediate downstream target gene of RUNX2, which drives the differentiation of pre-osteoblasts into mature osteoblasts and promotes bone formation, in part, by modulating the expression of many osteogenic genes including COL1A, SPARC, BSP-1, and BGLAP.⁷¹ Two mRNA isoforms, encoding 413 (Sp7S) and 431 (Sp7L) amino acids long peptides, respectively, through 3' alternative splicing of exon 3, are differentially expressed

TABLE 1 Summary of alternative splicing events for genes involved in the differentiation of embryonic stem cells (ESCs) and mesenchymal stem cells (MSCs)

Gene	Isoform	AS mode	Isoform function	Remarks	Regulation	Reference
ES cell differentiation						
FGF4	FGF4 WT	Full length	Promotes self-renew and pluripotency	Positive regulator of Erk1/2 phosphorylation	Unknown	24
	FGF4si	Exon skipping	Antagonizes self-renew and pluripotency	Negative regulator of Erk1/2 phosphorylation		
FOXP1	FOXP1	Mutually exclusive exon	Promotes hESC differentiation	Activates differentiation and suppresses pluripotency genes	MBNL1, RBFOX2	25
	FOXP1-ES	Mutually exclusive exon	Activates pluripotency genes	Activates pluripotency and suppresses differentiation genes		
PAX6	PAX6a	Exclusion of Cassette Exon	Neuroectoderm differentiation	Induces neuroectoderm	PNIN, ESRP1	31
	PAX6b	Inclusion of Cassette exon	Cell proliferation	No effect on neuroectoderm differentiation		
Tcf3 (TCF7L1)	Tcf3-E18a	Mutually exclusive exon	High in undifferentiated mES cells	Maintains pluripotency of ES cells	hnRNP H, hnRNP F	29
	Tcf3-E18b	Mutually exclusive exon	High in differentiated mES cells	Promotes differentiation		
Sall4	Sall4a	Alternative 5' splice site	Promotes pluripotency	Missing ZF domains, preferentially binds to ES-specific promoters	Unknown	30
	Sall4b	Alternative 5' splice site	Sufficient for pluripotency			
Mesenchymal stem cell differentiation						
RUNX2	RUNX2 WT	Full length	Promotes osteogenesis	Promotes early osteogenesis, activates osteocalcin	U2 snRNP, U2AF, SF3A1, SF3A3	14,44
	RUNX2Δ5	Cassette exon	Not known	Missing NLS		
	RUNX2Δ7	Cassette exon	Chondrocytes specific	RUNX2Δ7 antagonizes RUNX2, no activation of osteocalcin		
	RUNX2Δ5Δ7	Cassette exon	Not known	Missing NLS		
FosB	FosB WT	Full length	Osteogenesis	Truncation of transactivation domain	Unknown	45,46
	ΔFosB	Cassette exon	High in osteoblasts, promotes osteogenesis and inhibits adipogenesis			
OPN	Δ2ΔFosB	Alternative promoter, cassette exon	Promotes osteogenesis independent of AP-1 activity	Truncation of Fos-homology domain and transactivation domain		
	OPN-a	Cassette exon	Promotes angiogenesis and osteogenesis		Unknown	47,48
	OPN-b (ΔE5)	Cassette exon	Not known	Exon 5 containing serine phosphorylation site		
OSX (SP7)	OPN-c (ΔE4)	Cassette exon	Promotes angiogenesis and osteogenesis	Exon 4 containing tyrosine phosphorylation site		
	OSX-I	Alternative 5' splice site	Lower expression in osteoblasts		Unknown	49
	OSX-s	Alternative 5' splice site	Higher expression in osteoblasts and chondrocytes formation	Truncated N-terminal with distinct t/s activity		

(Continues)

TABLE 1 (Continued)

Gene	Isoform	AS mode	Isoform function	Remarks	Regulation	Reference
<i>Vegf-A</i>	<i>Vegf-A</i> WT	Full length	High expression in high-frequency mechanical loading	Matrix bound	Unknown	50
	<i>Vegf-A206</i>	Cassette exon	High expression in high-frequency mechanical loading	Matrix bound		
	<i>Vegf-A189</i>	Cassette exon	High expression in high-frequency mechanical loading	Matrix bound		
	<i>Vegf-A165</i>	Cassette exon	High expression in high-frequency mechanical loading	Partially matrix/partially secreted		
	<i>Vegf-A145</i>	Cassette exon	High expression in low-frequency mechanical loading	Matrix bound		
	<i>Vegf-A121</i>	Cassette exon	High expression in low-frequency mechanical loading	Secreted		
<i>Periostin (Postn)</i>	<i>Postn-001</i>	Full length	Osteoblast differentiation	Maximum number of RD domains	Unknown	51
	<i>Postn-001 - 008</i>	Cassette exons	Increased attachment of osteoblasts in periosteum	Various number of RD domains		
<i>NELL-1</i>	<i>NELL-1-810</i>	Full length	Promotes osteogenesis		Unknown	52
	<i>NELL-1-570</i>	Cassette exon	Enhanced MSC proliferation and osteogenesis	Truncation of LamG domain, required for heterodimer formation		
<i>PPARγ</i>	<i>PPARγ WT</i>	Full length	Promotes adipogenesis		SRSF1	12
	<i>PPARγΔ5</i>	Cassette exon	Disrupts adipogenesis	Dominant negative against the WT protein		
<i>PTHrP</i>	<i>PTHrP-L</i>	Alternative 3' splice site			Unknown	53,54
	<i>PTHrP-S</i>	Alternative 3' splice site	Promotes chondrocytes, suppresses MSC hypertrophy	EGF-dependent 3'-UTR enables RNA stability		
<i>mTOR</i>	<i>mTOR-int5</i>	Removal of intron	Promotes adipocyte differentiation	Increased mTOR signaling	Sam68	55
	<i>mTOR + Int5</i>	Retained intron	Suppresses adipocyte differentiation	Introduction of PTC and reduced mTOR signaling		
<i>LPIN1</i>	<i>LPIN1 WT</i>	Full length	Promotes adipocyte differentiation	Nuclear	SFRS10	56-58
	<i>LPIN1Δ7</i>	Cassette exon	Suppresses adipocyte maturation	Cytoplasmic		
<i>AMBN</i>	<i>AMBN WT</i>	Full length	Promotes MSC proliferation and enamel formation	Exon 5 peptide promotes MSC proliferation and osteogenesis	Unknown	59
	<i>AMBNΔ6,1-15</i>	Alternative 3' splice site	Promotes MSC proliferation and enamel formation	Exon 6 15 AA peptide suppresses MSC proliferation and osteogenesis		
<i>S6K1</i>	<i>S6K1 WT</i>	Cassette exon	Promotes adipogenesis		Sam68	60,61
	<i>S6K1Δ6a</i>	Cassette exon	Suppresses adipogenesis	Activates mTOR signaling		
<i>RUNX1T1</i>	<i>RUNX1T1-L</i>	Cassette exon	Suppresses adipogenesis	Increased m6A leads to SRSF2 binding and inclusion of exon 6	FTO, SRSF2	62

TABLE 1 (Continued)

Gene	Isoform	AS mode	Isoform function	Remarks	Regulation	Reference
Col2a1	RUNX1T1-S	Cassette exon	Promotes adipogenesis	Reduced m6A by FTO leads to lower SRSF2 binding to exon 6	Tia1, Sox9	15,63,64
	Col2a1 IIA	Alternative 5' splice site	High expression in chondroprogenitors			
TAF4	Col2a1 IIB	Cassette exon	High expression in mature chondrocytes			
	Col2a1 IIC	Alternative 5' splice site	Unknown	Alters IIA/IIB ratio		
PKM	Col2a1 IID	Alternative 5' splice site	Unknown	Alters IIA/IIB ratio		
	TAF4	Cassette exon	Promotes MSC proliferation and chondrogenic differentiation	Cell cycle exit via p53 activation and regulates Wnt signaling	Unknown	65
PKM2	TAF4ΔTAFH	Cassette exon	Promotes differentiation, including neural precursor cells	Tetramer formation, prefers oxidative phosphorylation	RBM4, SRSF3	32,66
	PKM1	Mutually exclusive exon	Promotes dedifferentiation	Dimer formation, prefers glycolysis		

between various cell and tissue types, especially fetal vs adult osteoblasts, and chondrocytes, suggesting a biological role of AS in cell or differentiation stage-specific AS.⁴⁷ Besides, multiple osteogenesis imperfecta patients harbor a splicing-sensitive mutation in the intron of COL1A1, implicating the mis-regulated intron retention as a causative mechanism for osteogenesis imperfecta.⁷²

Vascular endothelial growth factor A (VEGF-A), encoded by VEGFA, is a multifunctional growth factor that was shown to promote osteogenic differentiation of MSC by promoting mineralization in autocrine manner and angiogenesis in a paracrine manner.⁷³ Faure et al demonstrated a correlation between different VEGF-A splicing isoforms and the mechanical stress in osteoblastic cells, that is, the upregulation of the membrane bound VEGF-A isoforms under high mechanical stress and that of the secreted VEGF-A isoforms under low mechanical stress.⁴⁸ Generation of membrane-bound and secreted isoforms is carried out by a differential inclusion of three cassette exons (6A, 6B, and 7), within VEGF-A pre-mRNA. Nonetheless, regulated AS of VEGF-A by mechanical stress may provide an additional regulatory mechanism toward osteogenic differentiation, especially during osteogenesis upon a bone fracture.

Mechanical stress plays an important role during the formation of osteoblast's extracellular matrix (ECM). Physiologically, the lack of mechanical loading reduces the integrity of the ECM structure of osteocytes in the bone and eventually leads to bone loss. Roosa et al recently analyzed the global AS pattern of post-mechanical loading and subsequent bone formation using rats as a model system.⁷⁴ Their study identified numerous AS changes after the mechanical loading, potentially linking its AS pattern to the response of osteocytes to mechanical stress. A deeper understanding of the biological functions of AS isoforms during mechanical loading can further shed light on the regulation of bone maintenance and bone formation in general by AS in response to mechanical loading.

One of the known osteogenic growth factors, called neural epidermal growth-like (NEL)-like factor 1 (NELL-1), promotes osteogenesis in vitro and in vivo.⁷⁵ This gene generates two alternative isoforms, the full-length isoform (NELL-1810) and a shorter isoform (NELL-1570), by a combination of an alternative promoter and a 5' splice site for the inclusion of exon 2. Although both isoforms demonstrate an osteogenic differentiating effect, the shorter isoform had a stronger promoting effect on the proliferation of a murine primary MSC line and pericytes compared with the wild-type NELL-1.⁵⁰ However, the molecular mechanism by which NELL-1 splicing is regulated remains unknown.

Ameloblastin (AMBN) encodes a regulator of osteogenic differentiation, which is originally expressed in ameloblast for enamel formation and mineralization. AMBN pre-mRNA produces two isoforms, AMBN-WT and $\Delta 6_{1-15}$, by choosing alternative 3' splice sites of exon 6. It was recently shown that a synthetic peptide derived from the sequence of AMBN exon 5 promotes hMSC proliferation and activates expression of RUNX2 and OSX.⁵⁷ Intriguingly, 15 amino acid synthetic peptide corresponding to AMBN exon 6 abrogated the effect by exon 5 peptide. Although the regulatory mechanism by which AMBN exon 5/6 AS is regulated and the exact molecular function of the smaller




FIGURE 2 A schematic showing the important AS events involved in MSC differentiation to osteocytes, chondrocytes, adipocytes, and neurons. A schematic above each arrow represents the abbreviated splicing pattern with the resulting protein isoforms structure with indicated structural domains. A schematic below each arrow represents the relationship between expression level of each splicing isoform and functional consequence

AMBN isoform remain to be elucidated, AS of AMBN appears to be important for encoding the extracellular domain, the self-assembly of AMBN, and proliferation/differentiation of hMSC, particularly in osteogenesis.

5 | REGULATION OF CHONDROGENIC DIFFERENTIATION BY AS

SRY-related high-mobility group-box gene 9 (Sox9) is a master regulator of chondrogenesis.⁷⁶⁻⁷⁹ Hata et al demonstrated that Sox9 exerts its molecular function as an activator of chondrogenic genes, at least in part, through interacting with an RBP called p54nrb in a specialized region of the nucleus called the paraspeckle, as originally described in a mouse study.⁸⁰ The paraspeckle is a ribonucleoprotein body located in the interchromatin region and has been hypothesized to function as the retention center for a specific set of mRNAs as a controlling mechanism for gene expression.⁸¹ A recent report on a potential function of Sox9 in directly regulating alternative splicing raises an intriguing possibility that Sox9 may be a protein with dual functions—transcriptional and splicing regulation. Sox9 KD in colon tumor cancer cells induced a global change in AS without affecting their expression levels.^{15,82} Girardot et al identified several mutants in the HMG

domain, for example, W143R, Del400, and Del440, that lost transcriptional activity but not splicing.¹⁵ This indicates that Sox9 may play a dual role in transcription and splicing, and that Sox9 can regulate AS independently of its transcriptional activity. This idea that Sox9 is a splicing regulator, as well as a transcriptional factor, was validated by the fact that Sox9 affects the splicing of *Col2A1*, a key chondrogenic gene, via physical interaction with p54nrb in paraspeckles.⁸⁰

Another regulatory mechanism of chondrogenic differentiation via AS was described where a splicing variant of TATA binding protein-associate factor 4 (TAF4) repressed the proliferation of MSCs and promoted the chondrocyte differentiation of hMSCs.⁶³ TAF4 pre-mRNA was shown to produce four TAF4 mRNA isoforms, one of which produced an isoform lacking a TAFH domain (TAF4ΔTAFH), as a result of skipping two of the cassette exons (6 and 7). Kazantseva et al reported that the TAF4ΔTAFH expression level is directly correlated with the hMSC differentiation potential. Overexpression of the TAF4ΔTAFH isoform in hMSCs represses the proliferation of MSCs and preferentially promotes chondrogenic differentiation.⁶³ TAF4 is a subunit of the transcription factor IID (TFIID) complex of the transcriptional machinery, which typically controls a set of genes through the coordinated interaction of temporal- and spatial-specific transcriptional cofactors in response to external stimulation. Interestingly, Pijnappel et al recently demonstrated a higher expression level of

several components of the TFIID complex in mouse ES cells compared with its fibroblast counterpart and that KD of these TFIID components resulted in the differentiation of mouse ES cells.⁸³ Thus, this raises the interesting hypothesis where the increased expression of splicing variant TAF4ΔTAFH could perturb the stoichiometry and activity of TFIID, thereby modulating MSC differentiation and, in particular, the chondrogenic pathway.

Parathyroid hormone-related peptide (PTHrP) is a paracrine peptide hormone that is essential for osteogenesis and chondrogenesis. PTHrP has also previously been implicated in chondrogenesis, as it is secreted by immature chondrocytes, and chondrogenesis was shown to be negatively affected in PTHrP knockout mice.⁸⁴ The biological activity of PTHrP is exerted by the proteolytic cleavage of the translated peptide to produce three shorter peptides: PTH-like (amino acids 1-34), middle region (amino acids 38-94), and osteostatin (amino acids 107-139). PTHrP protein isoforms are generated from three different mRNA isoforms via distinct 3' terminal exons and a 3'-UTR through alternative 3' splice site selection.⁵² This results in three protein isoforms of 139, 141, and 173 amino acids with varying in the osteostatin domain, which have been implicated in differentially regulating chondrogenesis and hypertrophy in BM-MSCs.⁵¹ Since PTHrP pre-mRNA AS encodes three distinct osteostatin domain peptides, one can speculate that PTHrP exerts an important regulatory role in chondrogenesis and hypertrophy by AS.

The alternative splicing of *PTHrP* pre-mRNA can modulate MSC differentiation in two ways. First, the three mRNA isoforms with three distinct 3'-UTRs are subjected to the variable half-life, as the 3'-UTR serves as the binding site for RBPs and miRNA. By modulating alternative 3'-splice site selection, one can speculate that mRNA with a differing 3'-UTR can determine the subcellular localization and/or stability of the mature mRNA, leading to a control mechanism for the PTHrP level. Second, it has been suggested that the three PTHrP isoforms with variable C-terminal ends are implicated in the differential paracrine activity by producing three structurally distinct osteostatin peptides.⁵²

6 | REGULATION OF ADIPOGENIC DIFFERENTIATION BY AS

One genome-wide study examined the diversity of 3'-UTR structures of nascent transcripts bound to polysomes during adipogenesis.⁸⁵ The study not only found massive changes in the lengthening or shortening of the 3'-UTR after initiation of adipogenesis but also found a significant correlation between 3'-UTR length and the expression levels of the DEGs during adipogenesis.⁸⁵ The notion that there is a direct association between changes in gene expression and an increase in 3'-UTR diversity and length via alternative polyadenylation and splicing has been previously demonstrated in mouse embryogenesis.⁸⁶ The differentially expressed components of the polyadenylation machinery and pre-mRNA splicing are mainly responsible for generating different 3'-UTRs, as described above, leading to control of the expression levels of key regulators of adipogenesis through differential subcellular localization of mRNAs and translational repression.

The process of adipogenesis is activated by two key transcription factors: peroxisome proliferator-activated receptor γ (PPAR γ) and CCAAT/enhancer binding protein α (C/EBP α).⁸⁷ PPAR γ is a nuclear hormone receptor protein that binds to retinoic acid receptor alpha (RXR α) and activates downstream target genes associated with regulating fatty acid uptake and synthesis, thereby promoting adipogenesis.⁸⁷ Recently, a naturally occurring dominant negative isoform of PPAR γ has been discovered to regulate PPAR γ activity. Skipping cassette exon 5 generates a shorter isoform, PPAR γ Δ5, which is devoid of the entire ligand-binding domain, which may compete with WT PPAR γ by interacting with an unidentified coreceptor, thereby reducing the adipogenic differentiation of hMSCs.¹² Interestingly, the PPAR γ Δ5/PPAR γ ratio in human fat tissues was determined to be positively correlated to the BMI index, suggesting that adipose tissue dysfunction may be associated with the level of PPAR γ Δ5 isoform. The inclusion of exon 5 appears to be regulated by SRSF1 (SR domain-containing splicing regulator), also known as ASF/SF, which directly binds to PPAR γ pre-mRNA and promotes the exon's inclusion.¹² This represents a typical negative feedback loop in which the activation of adipogenic differentiation by PPAR γ expression, in response to external signaling cues, may be controlled by a counteractive balancing of the PPAR γ Δ5 expression via AS. Subsequently, an increased level of PPAR γ activation results in the reduction of SRSF1-mediated inclusion of exon 5, thereby producing more of PPAR γ Δ5 and negatively regulating the activity of WT PPAR γ .

LPIN1 was initially discovered as a protein associated with diseases of adipose tissue degeneration and was shown to be expressed predominantly in adipose tissues.⁵⁴ Peterfy et al demonstrated that *LPIN1* pre-mRNA undergoes alternative splicing to produce two isoforms, LPIN1 α and LPIN1 β , and observed that the isoform switch is accomplished during adipocyte differentiation from pre-adipocytes to mature adipocytes. Although the exact mechanism by which LPIN1 isoforms control adipocyte differentiation has not been completely elucidated, the differential subcellular localization patterns of the two isoforms may be involved in regulating adipocyte differentiation.⁵⁴

Src-associated substrate during mitosis of 68 kD (Sam68) is an RBP that modulates alternative splicing by binding to its consensus binding site, the U(U/A)AA motif. Sam68 has been shown to regulate the alternative splicing of various pre-mRNA substrates, such as *CD44*, *Bcl-x*, *Survival of Motor Neuron (SMN)*, *Srsf1*, and *Neurexin 1 (Nrxn1)*.⁸⁸⁻⁹² Sam68-dependent *mTOR* pre-mRNA splicing was shown to be correlated with adipogenesis and metabolism. Sam68 binds to the intron flanking exon 5 and 6 of *mTOR* pre-mRNA and recruits spliceosome to promote the splicing of the intron. Upon depletion of Sam68, the intron is retained and introduces a premature termination codon, resulting in the degradation of the *mTOR* mRNA and, subsequently, lower protein levels. Lower mTOR signaling, as a result, reduces the adipogenic differentiation. Song et al demonstrated that Sam68 expression in pre-adipocytes (3T3-L1 cells) suppresses the skipping of three cassette exons, exon 6a, b, and c, within *ribosomal S6 kinase (S6K1)* pre-mRNA, thereby inhibiting the expression of the dominant negative form, known as p31S6K1. The authors further demonstrated that the ectopic

expression of p31S6K1 causes the abrogation of adipogenesis by disrupting the mTOR pathway.⁵⁹

7 | CONCLUSION AND PERSPECTIVES

The biological functions of MSCs have many clinical implications, and it is imperative to understand their true cellular and molecular characteristics. In this review, we have summarized and emphasized the critical role of AS of key transcription factors and growth factors in MSC differentiation. The alternative splicing of key transcriptional regulators such as RUNX2, PPAR γ , and PTHrP during MSC differentiation into osteocytes, adipocytes, and chondrocytes plays an underappreciated role in modulating the proliferation and cell fate determination of MSCs. An expression of a specific isoform generated by AS contributes to determining the cell type specification by activating isoform-specific transcriptional targets or antagonizing the function of the WT protein. Furthermore, AS of many pre-mRNAs, which encode the critical determinants of MSC differentiation, ensures the MSC differentiation by fine-tuning the balance of isoform expressions, controlling the subcellular localization, and other intermolecular interactions such as RNA-protein and protein-protein interactions. AS of key transcription factors and growth factors is regulated by the differential expression of various RBPs (regulated both temporally and spatially) and modulates the binding of the spliceosome to the target splice site. One interesting regulatory mechanism that has been revealed in several studies suggests that key transcription factors regulate AS by directly binding to RNA and further recruiting splicing regulators to cis-acting elements. One potential of most work reported thus far in the role of AS in MSC differentiation are artifacts in the in vitro cultures, especially associated with origin, development, and endogenous functions of MSCs. Future studies with a focus on elucidating more detailed mechanisms and exact molecular functions in vivo will help to enhance the understanding of MSC biology and strengthen the therapeutic benefits of MSCs in the future.

ACKNOWLEDGMENTS

This work was supported by University of Macau Research Committee funds MYRG #2015-00169-FHS, #2016-00070-FHS, and #2017-00124-FHS, and Macau Science and Technology Development Fund (FDCT) #028/2015/A1, FDCT/095/2017/A2, and #095/2017/A1 to Ren-He Xu and 2017-00249-FHS to Jung Woo Park.

CONFLICT OF INTEREST

Ren-He Xu declared Intellectual property rights and consultancy role with ImStem Biotechnology, Inc. The other authors declared no potential conflicts of interest.

AUTHOR CONTRIBUTIONS

J.W.P., S.F., B.H., R.-H.X.: wrote the manuscript; J.W.P. and R.-H.X.: approved the manuscript.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Jung Woo Park  <https://orcid.org/0000-0003-0657-7401>

REFERENCES

1. Bianco P. "Mesenchymal" stem cells. *Annu Rev Cell Dev Biol.* 2014;30(1):677-704.
2. Spees JL, Lee RH, Gregory CA. Mechanisms of mesenchymal stem/stromal cell function. *Stem Cell Res Ther.* 2016;7(1):125.
3. Sipp D, Robey PG, Turner L. Clear up this stem-cell mess. *Nature.* 2018;561(7724):455-457.
4. Robey P. "Mesenchymal stem cells": fact or fiction, and implications in their therapeutic use. *F1000Res.* 2017;6(F1000 Faculty Rev):524-533.
5. Caplan AI. Medicinal signalling cells: they work, so use them. *Nature.* 2019;566(7742):39.
6. da Silva Meirelles L, Caplan AI, Nardi NB. In search of the in vivo identity of mesenchymal stem cells. *STEM CELLS.* 2008;26(9):2287-2299.
7. Gimble JM, Guilak F, Nuttall ME, Sathishkumar S, Vidal M, Bunnell BA. In vitro differentiation potential of mesenchymal stem cells. *Transfus Med Hemother.* 2008;35(3):228-238.
8. Sobacchi C, Palagano E, Villa A, Menale C. Soluble factors on stage to direct mesenchymal stem cells fate. *Front Bioeng Biotechnol.* 2017;5:32.
9. Almalki SG, Agrawal DK. Key transcription factors in the differentiation of mesenchymal stem cells. *Differentiation.* 2016;92(1-2):41-51.
10. James AW. Review of signaling pathways governing MSC osteogenic and adipogenic differentiation. *Scientifica (Cairo).* 2013;2013:684736.
11. Chen Q, Shou P, Zheng C, et al. Fate decision of mesenchymal stem cells: adipocytes or osteoblasts? *Cell Death Differ.* 2016;23(7):1128-1139.
12. Aprile M, Cataldi S, Ambrosio MR, et al. PPAR γ Delta5, a naturally occurring dominant-negative splice isoform, impairs PPAR γ function and adipocyte differentiation. *Cell Rep.* 2018;25(6):1577-1592.e1576.
13. van de Peppel J, Strini T, Tilburg J, Westerhoff H, van Wijnen AJ, van Leeuwen JP. Identification of three early phases of cell-fate determination during osteogenic and adipogenic differentiation by transcription factor dynamics. *Stem Cell Rep.* 2017;8(4):947-960.
14. Makita N, Suzuki M, Asami S, et al. Two of four alternatively spliced isoforms of RUNX2 control osteocalcin gene expression in human osteoblast cells. *Gene.* 2008;413(1-2):8-17.
15. Girardot M, Bayet E, Maurin J, Fort P, Roux P, Raynaud P. SOX9 has distinct regulatory roles in alternative splicing and transcription [in English]. *Nucleic Acids Res.* 2018;46(17):9106-9118.
16. Han H, Braunschweig U, Gonatopoulos-Pournatzis T, et al. Multilayered control of alternative splicing regulatory networks by transcription factors [in English]. *Mol Cell.* 2017;65(3):539.
17. Kornblihtt AR, De la Mata M, Fededa JP, et al. Multiple links between transcription and splicing [in English]. *RNA.* 2004;10(10):1489-1498.
18. Pan Q, Shai O, Lee LJ, Frey BJ, Blencowe BJ. Deep surveying of alternative splicing complexity in the human transcriptome by high-throughput sequencing. *Nat Genet.* 2008;40(12):1413-1415.
19. Barbosa-Morais NL, Irimia M, Pan Q, et al. The evolutionary landscape of alternative splicing in vertebrate species. *Science.* 2012;338(6114):1587-1593.
20. Merkin J, Russell C, Chen P, Burge CB. Evolutionary dynamics of gene and isoform regulation in mammalian tissues. *Science.* 2012;338(6114):1593-1599.

21. Nilsen TW, Graveley BR. Expansion of the eukaryotic proteome by alternative splicing. *Nature*. 2010;463(7280):457-463.
22. Nilsen TW. The spliceosome: the most complex macromolecular machine in the cell? [in English]. *Bioessays*. 2003;25(12):1147-1149.
23. Zhang XF, Yan CY, Hang J, Finci LI, Lei J, Shi Y. An atomic structure of the human spliceosome [in English]. *Cell*. 2017;169(5):918.
24. Maysnar Y, Rom E, Chumakov I, Kronman A, Yayon A, Benvenisty N. Fibroblast growth factor 4 and its novel splice isoform have opposing effects on the maintenance of human embryonic stem cell self-renewal [in English]. *STEM CELLS*. 2008;26(3):767-774.
25. Gabut M, Samavarchi-Tehrani P, Wang X, et al. An alternative splicing switch regulates embryonic stem cell pluripotency and reprogramming. *Cell*. 2011;147(1):132-146.
26. Han H, Irimia M, Ross PJ, et al. MBNL proteins repress ES-cell-specific alternative splicing and reprogramming. *Nature*. 2013;498(7453):241-245.
27. Zhang T, Lin Y, Liu J, et al. Rbm24 regulates alternative splicing switch in embryonic stem cell cardiac lineage differentiation [in English]. *STEM CELLS*. 2016;34(7):1776-1789.
28. Zhang XQ, Huang CT, Chen J, et al. Pax6 is a human neuroectoderm cell fate determinant [in English]. *Cell Stem Cell*. 2010;7(1):90-100.
29. Salomonis N, Schlieve CR, Pereira L, et al. Alternative splicing regulates mouse embryonic stem cell pluripotency and differentiation. *Proc Natl Acad Sci USA*. 2010;107(23):10514-10519.
30. Rao S, Zhen S, Roumiantsev S, McDonald LT, Yuan GC, Orkin SH. Differential roles of Sall4 isoforms in embryonic stem cell pluripotency. *Mol Cell Biol*. 2010;30(22):5364-5380.
31. Park JW, Yang J, Xu RH. PAX6 alternative splicing and corneal development. *Stem Cells Dev*. 2018;27(6):367-377.
32. Su CH, Hung KY, Hung SC, Tam WY. RBM4 regulates neuronal differentiation of mesenchymal stem cells by modulating alternative splicing of pyruvate kinase M. *Mol Cell Biol*. 2017;37(3):e00466-16.
33. Chen M, Zhang J, Manley JL. Turning on a fuel switch of cancer: hnRNP proteins regulate alternative splicing of pyruvate kinase mRNA. *Cancer Res*. 2010;70(22):8977-8980.
34. Calabretta S, Bielli P, Passacantilli I, et al. Modulation of PKM alternative splicing by PTBP1 promotes gemcitabine resistance in pancreatic cancer cells [in English]. *Oncogene*. 2016;35(16):2031-2039.
35. Kuranaga Y, Sugito N, Shinohara H, et al. SRSF3, a splicer of the PKM gene, regulates cell growth and maintenance of cancer-specific energy metabolism in colon cancer cells. *Int J Mol Sci*. 2018;19(10):3012-3028.
36. Clower CV, Chatterjee D, Wang ZX, Cantley LC, Vander Heiden MG, Krainer AR. The alternative splicing repressors hnRNP A1/A2 and PTB influence pyruvate kinase isoform expression and cell metabolism [in English]. *Proc Natl Acad Sci USA*. 2010;107(5):1894-1899.
37. Laaref AM, Manchon L, Bareche Y, et al. The core spliceosomal factor U2AF1 controls cell-fate determination via the modulation of transcriptional networks. *RNA Biol*. 2020;54(5):1-15.
38. Peffers MJ, Collins J, Fang Y, et al. Age-related changes in mesenchymal stem cells identified using a multi-omics approach. *Eur Cell Mater*. 2016;31:136-159.
39. Peffers MJ, Goljanek-Whysall K, Collins J, et al. Decoding the regulatory landscape of ageing in musculoskeletal engineered tissues using genome-wide DNA methylation and RNASeq. *PLoS One*. 2016;11(8):e0160517.
40. Fiszbein A, Kornblihtt AR. Alternative splicing switches: important players in cell differentiation [in English]. *Bioessays*. 2017;39(6):1-11.
41. Baralle FE, Giudice J. Alternative splicing as a regulator of development and tissue identity [in English]. *Nat Rev Mol Cell Bio*. 2017;18(7):437-451.
42. Jensen ED, Niu L, Caretti G, et al. p68 (Ddx5) interacts with Runx2 and regulates osteoblast differentiation. *J Cell Biochem*. 2008;103(5):1438-1451.
43. Sabatakos G, Rowe GC, Kveiborg M, et al. Doubly truncated FosB isoform (Delta2DeltaFosB) induces osteosclerosis in transgenic mice and modulates expression and phosphorylation of Smads in osteoblasts independent of intrinsic AP-1 activity. *J Bone Miner Res*. 2008;23(5):584-595.
44. Sabatakos G, Sims NA, Chen J, et al. Overexpression of DeltaFosB transcription factor(s) increases bone formation and inhibits adipogenesis. *Nat Med*. 2000;6(9):985-990.
45. Gimba ER, Tilli TM. Human osteopontin splicing isoforms: known roles, potential clinical applications and activated signaling pathways. *Cancer Lett*. 2013;331(1):11-17.
46. Lee GS, Salazar HF, Joseph G, et al. Osteopontin isoforms differentially promote arteriogenesis in response to ischemia via macrophage accumulation and survival. *Lab Invest*. 2019;99(3):331-345.
47. Milona MA, Gough JE, Edgar AJ. Expression of alternatively spliced isoforms of human Sp7 in osteoblast-like cells. *BMC Genomics*. 2003;4:43.
48. Faure C, Linossier MT, Malaval L, et al. Mechanical signals modulated vascular endothelial growth factor-A (VEGF-A) alternative splicing in osteoblastic cells through actin polymerisation. *Bone*. 2008;42(6):1092-1101.
49. Horiuchi K, Amizuka N, Takeshita S, et al. Identification and characterization of a novel protein, periostin, with restricted expression to periosteum and periodontal ligament and increased expression by transforming growth factor beta. *J Bone Miner Res*. 1999;14(7):1239-1249.
50. Pang S, Shen J, Liu Y, et al. Proliferation and osteogenic differentiation of mesenchymal stem cells induced by a short isoform of NELL-1. *STEM CELLS*. 2015;33(3):904-915.
51. Lee JM, Im GI. PTHrP isoforms have differing effect on chondrogenic differentiation and hypertrophy of mesenchymal stem cells [in English]. *Biochem Bioph Res Co*. 2012;421(4):819-824.
52. Sellers RS, Luchin AI, Richard V, Brena RM, Lima D, Rosol TJ. Alternative splicing of parathyroid hormone-related protein mRNA: expression and stability. *J Mol Endocrinol*. 2004;33(1):227-241.
53. Huot ME, Vogel G, Zabarauskas A, et al. The Sam68 STAR RNA-binding protein regulates mTOR alternative splicing during adipogenesis [in English]. *Mol Cell*. 2012;46(2):187-199.
54. Peterfy M, Phan J, Reue K. Alternatively spliced lipin isoforms exhibit distinct expression pattern, subcellular localization, and role in adipogenesis. *J Biol Chem*. 2005;280(38):32883-32889.
55. Li H, Cheng YM, Wu WW, et al. SRSF10 regulates alternative splicing and is required for adipocyte differentiation [in English]. *Mol Cell Biol*. 2014;34(12):2198-2207.
56. Pihlajamaki J, Lerin C, Ikonen P, et al. Expression of the splicing factor gene SFRS10 is reduced in human obesity and contributes to enhanced lipogenesis [in English]. *Cell Metab*. 2011;14(2):208-218.
57. Stakkestad O, Lyngstadaas SP, Vondrasek J, et al. Ameloblastin peptides modulates the osteogenic capacity of human mesenchymal stem cells. *Front Physiol*. 2017;8:58.
58. Ben-Hur V, Denichenko P, Siegfried Z, et al. S6K1 alternative splicing modulates its oncogenic activity and regulates mTORC1 [in English]. *Cell Rep*. 2013;3(1):103-115.
59. Song J, Richard S. Sam68 regulates S6K1 alternative splicing during adipogenesis. *Mol Cell Biol*. 2015;35(11):1926-1939.
60. Zhao X, Yang Y, Sun BF, et al. FTO-dependent demethylation of N6-methyladenosine regulates mRNA splicing and is required for adipogenesis [in English]. *Cell Res*. 2014;24(12):1403-1419.
61. McAlinden A, Liang L, Mukudai Y, Imamura T, Sandell LJ. Nuclear protein TIA-1 regulates COL2A1 alternative splicing and interacts with precursor mRNA and genomic DNA. *J Biol Chem*. 2007;282(33):24444-24454.
62. Hering TM, Wirthlin L, Ravindran S, McAlinden A. Changes in type II procollagen isoform expression during chondrogenesis by disruption

- of an alternative 5' splice site within Col2a1 exon 2 [in English]. *Matrix Biol.* 2014;36:51-63.
63. Kazantseva J, Kivil A, Tints K, Kazantseva A, Neuman T, Palm K. Alternative splicing targeting the hTAF4-TAFH domain of TAF4 represses proliferation and accelerates chondrogenic differentiation of human mesenchymal stem cells. *PLoS One.* 2013;8(10):e74799.
 64. Lin JC, Tam WY. RBM4 down-regulates PTB and antagonizes its activity in muscle cell-specific alternative splicing. *J Cell Biol.* 2011;193(3):509-520.
 65. Komori T. Regulation of osteoblast differentiation by Runx2 [in English]. *Osteoimmunology.* 2010;658:43-49.
 66. Sun L, Vitolo MI, Qiao M, Anglin IE, Passaniti A. Regulation of TGFbeta1-mediated growth inhibition and apoptosis by RUNX2 isoforms in endothelial cells. *Oncogene.* 2004;23(27):4722-4734.
 67. Venables JP, Lapasset L, Gadea G, et al. MBNL1 and RBFOX2 cooperate to establish a splicing programme involved in pluripotent stem cell differentiation [in English]. *Nat Commun.* 2013;4:2480.
 68. Joshi P, Halene S, Abdel-Wahab O. How do messenger RNA splicing alterations drive myelodysplasia? [in English]. *Blood.* 2017;129(18):2465-2470.
 69. Park JW, Parisky K, Celotto AM, Reenan RA, Graveley BR. Identification of alternative splicing regulators by RNA interference in drosophila. *Proc Natl Acad Sci USA.* 2004;101(45):15974-15979.
 70. Buskin A, Zhu L, Chichagova V, et al. Disrupted alternative splicing for genes implicated in splicing and ciliogenesis causes PRPF31 retinitis pigmentosa. *Nat Commun.* 2018;9(1):4234.
 71. Sinha KM, Zhou X. Genetic and molecular control of osterix in skeletal formation. *J Cell Biochem.* 2013;114(5):975-984.
 72. Johnson C, Primorac D, McKinstry M, McNeil J, Rowe D, Lawrence JB. Tracking COL1A1 RNA in osteogenesis imperfecta. Splice-defective transcripts initiate transport from the gene but are retained within the SC35 domain. *J Cell Biol.* 2000;150(3):417-432.
 73. Mayer H, Bertram H, Lindenmaier W, Korff T, Weber H, Weich H. Vascular endothelial growth factor (VEGF-A) expression in human mesenchymal stem cells: autocrine and paracrine role on osteoblastic and endothelial differentiation. *J Cell Biochem.* 2005;95(4):827-839.
 74. Mantila Roosa SM, Liu Y, Turner CH. Alternative splicing in bone following mechanical loading. *Bone.* 2011;48(3):543-551.
 75. Zhang X, Zara J, Siu RK, Ting K, Soo C. The role of NELL-1, a growth factor associated with craniosynostosis, in promoting bone regeneration. *J Dent Res.* 2010;89(9):865-878.
 76. Hino K, Saito A, Kido M, et al. Master regulator for chondrogenesis, Sox9, regulates transcriptional activation of the endoplasmic reticulum stress transducer BFBF2H7/CREB3L2 in chondrocytes [in English]. *J Biol Chem.* 2014;289(20):13810-13820.
 77. Lefebvre V, Dvir-Ginzberg M. SOX9 and the many facets of its regulation in the chondrocyte lineage [in English]. *Connect Tissue Res.* 2017;58(1):2-14.
 78. Oh CD, Lu Y, Liang SD, et al. SOX9 regulates multiple genes in chondrocytes, including genes encoding ECM proteins, ECM modification enzymes, receptors, and transporters (vol 9, e107577, 2014) [in English]. *PLoS One.* 2015;10(11):e0143156.
 79. Akiyama H, Chaboissier MC, Martin JF, et al. The transcription factor Sox9 has essential roles in successive steps of the chondrocyte differentiation pathway and is required for expression of Sox5 and Sox6 [in English]. *J Bone Miner Res.* 2002;17:S142-S142.
 80. Hata K, Nishimura R, Muramatsu S, et al. Paraspeckle protein p54nrB links Sox9-mediated transcription with RNA processing during chondrogenesis in mice. *J Clin Invest.* 2008;118(9):3098-3108.
 81. Fox AH, Lamond AI. Paraspeckles [in English]. *Csh Perspect Biol.* 2010;2(7):a000687.
 82. Guzzo RM, Scanlon V, Sanjay A, Xu RH, Drissi H. Establishment of human cell type-specific iPS cells with enhanced chondrogenic potential. *Stem Cell Rev Rep.* 2014;10(6):820-829.
 83. Pijnappel WWMP, Esch D, Baltissen MPA, et al. A central role for TFIIID in the pluripotent transcription circuitry [in English]. *Nature.* 2013;495(7442):516-519.
 84. Kronenberg HM. PTHrP and skeletal development [in English]. *Ann N Y Acad Sci.* 2006;1068:1-13.
 85. Spangenberg L, Shigunov P, Abud APR, et al. Polysome profiling shows extensive posttranscriptional regulation during human adipocyte stem cell differentiation into adipocytes [in English]. *Stem Cell Res.* 2013;11(2):902-912.
 86. Ji Z, Lee JY, Pan ZH, et al. Progressive lengthening of 3' untranslated regions of mRNAs by alternative polyadenylation during mouse embryonic development (vol 106, pg 7028, 2009) [in English]. *Proc Natl Acad Sci USA.* 2009;106(23):9535-9535.
 87. Darlington GJ, Ross SE, MacDougald OA. The role of C/EBP genes in adipocyte differentiation [in English]. *J Biol Chem.* 1998;273(46):30057-30060.
 88. Matter N, Herrlich P, Konig H. Signal-dependent regulation of splicing via phosphorylation of Sam68. *Nature.* 2002;420(6916):691-695.
 89. Paronetto MP, Achsel T, Massiello A, Chalfant CE, Sette C. The RNA-binding protein Sam68 modulates the alternative splicing of Bcl-x. *J Cell Biol.* 2007;176(7):929-939.
 90. Pedrotti S, Bielli P, Paronetto MP, et al. The splicing regulator Sam68 binds to a novel exonic splicing silencer and functions in SMN2 alternative splicing in spinal muscular atrophy. *EMBO J.* 2010;29(7):1235-1247.
 91. Valacca C, Bonomi S, Buratti E, et al. Sam68 regulates EMT through alternative splicing-activated nonsense-mediated mRNA decay of the SF2/ASF proto-oncogene. 2010;191(1):87-99.
 92. Iijima T, Wu K, Witte H, et al. SAM68 regulates neuronal activity-dependent alternative splicing of neuroligin-1. *Cell.* 2011;147(7):1601-1614.

How to cite this article: Park JW, Fu S, B Huang, Xu R-H. Alternative splicing in mesenchymal stem cell differentiation. *Stem Cells.* 2020;38:1229-1240. <https://doi.org/10.1002/stem.3248>