

1,2-Bis[2-(2-nitro-1H-imidazol-1-yl)-ethoxy]ethane

Shu-Xian Li,^{a‡} Lin Zhu,^a Hua-Min Li,^{a§} Hoong-Kun Fun^{b*} and Suchada Chantrapromma^{c¶}

^aDepartment of Chemistry, Beijing Normal University, Beijing 100875, People's Republic of China, ^bX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^cCrystal Materials Research Unit, Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat-Yai, Songkhla 90112, Thailand
Correspondence e-mail: hkfun@usm.my


Received 18 July 2008; accepted 27 July 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.037; wR factor = 0.106; data-to-parameter ratio = 15.2.

In the crystal structure, the title compound, $\text{C}_{12}\text{H}_{16}\text{N}_6\text{O}_6$, lies on an inversion centre. The molecule has an antiperiplanar conformation with respect to the C—C bond of the central ethane unit and the two imidazole rings are parallel to each other. The dihedral angle between the imidazole ring and the mean plane of the C and O atoms of the bis(ethoxy)ethane group is $76.04(6)^\circ$. The molecules are stacked along the c axis through a weak C—H \cdots O interaction and a $\pi\cdots\pi$ interaction between the imidazole rings with a centroid-centroid distance of $3.5162(6)$ Å. An intramolecular C—H \cdots O hydrogen bond is also present.

Related literature

For bond-length data, see: Allen *et al.* (1987). For the applications of nitroimidazoles, see, for example: Abdel-Jalil *et al.* (2006); Kennedy *et al.* (2006); Nagasawa *et al.* (2006); Nunn *et al.* (1995).



[‡] Department of Chemistry, Handan College, Handan, Hebei 056005, People's Republic of China.

[§] Additional correspondence author, e-mail: leehuamin@sina.com.

[¶] Additional correspondence author, e-mail: suchada.c@psu.ac.th.

Experimental

Crystal data

$\text{C}_{12}\text{H}_{16}\text{N}_6\text{O}_6$
 $M_r = 340.31$
Monoclinic, $P2_1/c$
 $a = 7.0534(1)$ Å
 $b = 15.5792(2)$ Å
 $c = 6.8069(1)$ Å
 $\beta = 99.560(1)^\circ$
 $V = 737.60(2)$ Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.13$ mm⁻¹
 $T = 100.0(1)$ K
 $0.40 \times 0.30 \times 0.15$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.952$, $T_{\max} = 0.982$
11084 measured reflections
2140 independent reflections
1864 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$
 $wR(F^2) = 0.106$
 $S = 1.05$
2140 reflections
141 parameters
All H-atom parameters refined
 $\Delta\rho_{\max} = 0.49$ e Å⁻³
 $\Delta\rho_{\min} = -0.32$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|---------------------------------|------------|-------------|-------------|---------------|
| C4—H4A \cdots O3 ⁱ | 0.977 (13) | 2.404 (13) | 3.3707 (11) | 170.1 (10) |
| C4—H4B \cdots O2 | 0.944 (14) | 2.408 (13) | 2.8169 (13) | 105.9 (9) |

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: APEX2 (Bruker, 2005); cell refinement: APEX2; data reduction: SAINT (Bruker, 2005); program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2003).

The authors gratefully acknowledge the financial assistance of Beijing Normal University. The authors also thank Universiti Sains Malaysia for the Research University Golden Goose (grant No. 1001/PFIZIK/811012).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2317).

References

- Abdel-Jalil, R. J., Uebele, M., Ehrlichmann, W., Voelter, W. & Machulla, H. J. (2006). *J. Radioanal. Nucl. Chem.* **267**, 557–560.
Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Kennedy, D. C., Wu, A., Patrick, B. O. & James, B. R. (2006). *J. Inorg. Biochem.* **100**, 1974–1982.
Nagasawa, H., Uto, Y., Kirk, K. L. & Hori, H. (2006). *Biol. Pharm. Bull.* **29**, 2335–2342.
Nunn, A., Linder, K. & Strauss, W. H. (1995). *Eur. J. Nucl. Med.* **22**, 265–280.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o1690 [doi:10.1107/S1600536808023726]

1,2-Bis[2-(2-nitro-1*H*-imidazol-1-yl)ethoxy]ethane

S.-X. Li, L. Zhu, H.-M. Li, H.-K. Fun and S. Chantrapromma

Comment

Depending on the availability of oxygen in tissue, nitroimidazoles can undergo different intracellular metabolism. In a normal cell, the molecule undergoes reduction to become a potentially reactive species and can be reoxidized in the presence of normal oxygen levels. In hypoxic tissue, however, the low oxygen concentration is not able to effectively reoxidize the molecule which results in more reactive intermediates that bind with components of hypoxic tissues (Nunn *et al.*, 1995). Thus these compounds can function as hypoxia markers for imaging of hypoxic cells and have received much attention in medicinal and clinic studies (Abdel-Jalil *et al.*, 2006; Kennedy *et al.*, 2006; Nagasawa *et al.*, 2006). In an attempt to develop new hypoxic cell radiosensitizers, we present herein the synthesis and crystal structure of the title nitroimidazole compound, (I).

The molecule of the title compound, C₁₂H₁₆N₆O₆, lies on an crystallographic inversion centre, so the asymmetric unit contains half of the molecule. The molecular structure has an antiperiplanar conformation with the two imidazole rings parallel to each other. The imidazole ring is planar, within a deviation of ± 0.003 Å. The nitro group is twisted from the mean plane of imidazole ring with torsion angles O1–N3–C1–N1 = -6.49 (16)° and O2–N3–C1–N1 = 173.61 (9)°. Atoms C5, O3, C6, C5ⁱ, O3ⁱ and C6ⁱ [symmetry code: (i) $1 - x, 1 - y, -z$] lie on the same plane. The interplanar angle between the C5/O3/C6/C5ⁱ/O3ⁱ/C6ⁱ plane and the imidazole ring (N1/N2/C1–C3) is 76.04 (6)°. The conformation of the ethoxyethane group is (-)-*syn*-clinal with respect to the imidazole ring, which is reflected by the torsion angle N2–C4–C5–O3 = -72.25 (9)°. Bond distances and angles have normal values (Allen *et al.*, 1987).

The crystal packing of (I) in Fig. 2 shows that the molecules are linked by weak C—H···O interactions (Table 1) and stacked into columns along the *c* axis. The molecules in the adjacent columns are in a face-to-face fashion (Fig. 3). The crystal is stabilized by a weak C—H···O interaction (Table 1). A π ··· π interaction was also observed in the crystal with the Cg₁···Cg₁ⁱⁱ [symmetry code: (ii) $x, 1/2 - y, -1/2 + z$] distance of 3.5162 (6) Å; Cg₁ is the centroid of the N1/N2/C1–C3 ring.

Experimental

To a solution of the triethyleneglycol ditosylate (0.458 g, 1.0 mmol) and triethylamine (244 mg, 2.4 mmol) in DMF (10 ml) was added a solution of 2-nitroimidazole (249 mg, 2.2 mmol) in DMF (10 ml) under argon. The mixture was stirred at 313 K for 4 days. After concentration on the rotary unit under reduced pressure, ethyl acetate (80 ml) was then added to the reaction residue, washed with water (20 ml \times 3), dried (Na₂SO₄) and the organic layer was evaporated to dryness and subjected to chromatography on silica with 50% EtOAc-hexane to afford the desired compound (I) (0.255 g, yield 75%). Analysis calcd for C₁₂H₁₆N₆O₆: C 42.35, H 4.74, N 24.70%; found: C 42.01, H 4.71, N 24.43%. Single crystals suitable for X-ray diffraction analysis were obtained by the slow diffusion of hexane into the dichloromethane solution of the title compound.

Refinement

All H atoms were located in a difference map and refined isotropically.

Figures




Fig. 1. The molecular structure of (I), showing 50% probability displacement ellipsoids and the atomic numbering.




Fig. 2. The crystal packing of (I), viewed down the *c* axis, showing stacking of molecules along the *c* axis. Hydrogen bonds are shown as dashed lines.




Fig. 3. The crystal packing of (I), viewed down the *a* axis. Hydrogen bonds are shown as dashed lines.

1,2-Bis(2-(2-nitro-1*H*-imidazol-1-yl)ethoxy)ethane

Crystal data

$C_{12}H_{16}N_6O_6$

$M_r = 340.31$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 7.0534\ (1)\ \text{\AA}$

$b = 15.5792\ (2)\ \text{\AA}$

$c = 6.8069\ (1)\ \text{\AA}$

$\beta = 99.560\ (1)^\circ$

$V = 737.597\ (18)\ \text{\AA}^3$

$Z = 2$

$F_{000} = 356$

$D_x = 1.532\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2140 reflections

$\theta = 2.6\text{--}30.0^\circ$

$\mu = 0.13\ \text{mm}^{-1}$

$T = 100.0\ (1)\ \text{K}$

Block, colorless

$0.40 \times 0.30 \times 0.15\ \text{mm}$

Data collection

Bruker SMART APEXII CCD area-detector
diffractometer

2140 independent reflections

| | |
|--|--|
| Radiation source: fine-focus sealed tube | 1864 reflections with $I > 2\sigma(I)$ |
| Monochromator: graphite | $R_{\text{int}} = 0.024$ |
| Detector resolution: 8.33 pixels mm^{-1} | $\theta_{\text{max}} = 30.0^\circ$ |
| $T = 100.0(1)$ K | $\theta_{\text{min}} = 2.6^\circ$ |
| ω scans | $h = -9 \rightarrow 9$ |
| Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005) | $k = -19 \rightarrow 21$ |
| $T_{\text{min}} = 0.952$, $T_{\text{max}} = 0.982$ | $l = -9 \rightarrow 9$ |
| 11084 measured reflections | |

Refinement

| | |
|--|--|
| Refinement on F^2 | Secondary atom site location: difference Fourier map |
| Least-squares matrix: full | Hydrogen site location: inferred from neighbouring sites |
| $R[F^2 > 2\sigma(F^2)] = 0.037$ | All H-atom parameters refined |
| $wR(F^2) = 0.106$ | $w = 1/[\sigma^2(F_o^2) + (0.0604P)^2 + 0.1626P]$ |
| $S = 1.05$ | where $P = (F_o^2 + 2F_c^2)/3$ |
| 2140 reflections | $(\Delta/\sigma)_{\text{max}} = 0.001$ |
| 141 parameters | $\Delta\rho_{\text{max}} = 0.50 \text{ e } \text{\AA}^{-3}$ |
| Primary atom site location: structure-invariant direct methods | $\Delta\rho_{\text{min}} = -0.32 \text{ e } \text{\AA}^{-3}$ |
| | Extinction correction: none |

Special details

Experimental. The low-temperature data was collected with the Oxford Cryosystem Cobra low-temperature attachment.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. The highest residual electron density peak is located at 0.76 Å from C6 and the deepest hole is located at 1.04 Å from C1.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|----|---------------|-------------|--------------|----------------------------------|
| O1 | -0.11911 (13) | 0.21335 (6) | 0.31449 (16) | 0.0405 (2) |
| O2 | -0.12171 (11) | 0.35244 (5) | 0.34055 (13) | 0.0319 (2) |
| O3 | 0.39505 (9) | 0.47777 (4) | 0.21737 (9) | 0.01816 (16) |
| N1 | 0.25143 (13) | 0.20738 (5) | 0.49696 (13) | 0.02131 (19) |
| N2 | 0.26946 (11) | 0.35116 (5) | 0.49565 (11) | 0.01551 (17) |
| N3 | -0.04022 (13) | 0.28202 (6) | 0.36307 (14) | 0.0243 (2) |

supplementary materials

| | | | | |
|-----|--------------|-------------|--------------|--------------|
| C1 | 0.15885 (13) | 0.27986 (6) | 0.45100 (14) | 0.0183 (2) |
| C2 | 0.43305 (14) | 0.23267 (6) | 0.57559 (14) | 0.0209 (2) |
| C3 | 0.44699 (13) | 0.32065 (6) | 0.57493 (13) | 0.01808 (19) |
| C4 | 0.22148 (14) | 0.44265 (6) | 0.47476 (13) | 0.01791 (19) |
| C5 | 0.20547 (13) | 0.47446 (6) | 0.26222 (13) | 0.01829 (19) |
| C6 | 0.39581 (14) | 0.49935 (6) | 0.01416 (13) | 0.0190 (2) |
| H2 | 0.535 (2) | 0.1921 (9) | 0.625 (2) | 0.024 (3)* |
| H3 | 0.5512 (19) | 0.3594 (8) | 0.6195 (19) | 0.020 (3)* |
| H4A | 0.3268 (19) | 0.4727 (8) | 0.5584 (19) | 0.020 (3)* |
| H4B | 0.1055 (19) | 0.4504 (8) | 0.5245 (19) | 0.022 (3)* |
| H5A | 0.1496 (18) | 0.5319 (8) | 0.2511 (18) | 0.020 (3)* |
| H5B | 0.1250 (17) | 0.4375 (8) | 0.1698 (18) | 0.017 (3)* |
| H6A | 0.3216 (18) | 0.4569 (8) | -0.0729 (19) | 0.021 (3)* |
| H6B | 0.3398 (17) | 0.5550 (8) | -0.0139 (18) | 0.017 (3)* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|----|------------|------------|------------|-------------|-------------|-------------|
| O1 | 0.0272 (4) | 0.0366 (5) | 0.0539 (6) | -0.0129 (3) | -0.0041 (4) | -0.0045 (4) |
| O2 | 0.0186 (4) | 0.0345 (4) | 0.0408 (5) | 0.0023 (3) | -0.0004 (3) | 0.0128 (3) |
| O3 | 0.0189 (3) | 0.0223 (3) | 0.0136 (3) | -0.0025 (2) | 0.0038 (2) | 0.0027 (2) |
| N1 | 0.0249 (4) | 0.0174 (4) | 0.0216 (4) | 0.0000 (3) | 0.0037 (3) | 0.0023 (3) |
| N2 | 0.0152 (3) | 0.0162 (4) | 0.0152 (3) | 0.0008 (3) | 0.0028 (3) | 0.0021 (2) |
| N3 | 0.0187 (4) | 0.0287 (5) | 0.0246 (4) | -0.0047 (3) | 0.0009 (3) | 0.0043 (3) |
| C1 | 0.0166 (4) | 0.0198 (4) | 0.0183 (4) | -0.0018 (3) | 0.0021 (3) | 0.0028 (3) |
| C2 | 0.0217 (5) | 0.0206 (4) | 0.0199 (4) | 0.0050 (3) | 0.0024 (3) | 0.0018 (3) |
| C3 | 0.0156 (4) | 0.0208 (4) | 0.0173 (4) | 0.0021 (3) | 0.0013 (3) | 0.0006 (3) |
| C4 | 0.0216 (4) | 0.0155 (4) | 0.0179 (4) | 0.0033 (3) | 0.0069 (3) | 0.0023 (3) |
| C5 | 0.0196 (4) | 0.0177 (4) | 0.0183 (4) | 0.0019 (3) | 0.0054 (3) | 0.0039 (3) |
| C6 | 0.0218 (5) | 0.0213 (4) | 0.0139 (4) | -0.0030 (3) | 0.0035 (3) | 0.0035 (3) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|----------|-------------|--------------------|-------------|
| O1—N3 | 1.2258 (12) | C2—H2 | 0.977 (14) |
| O2—N3 | 1.2361 (12) | C3—H3 | 0.961 (13) |
| O3—C5 | 1.4211 (11) | C4—C5 | 1.5156 (12) |
| O3—C6 | 1.4243 (10) | C4—H4A | 0.977 (13) |
| N1—C1 | 1.3160 (12) | C4—H4B | 0.943 (13) |
| N1—C2 | 1.3620 (13) | C5—H5A | 0.976 (12) |
| N2—C1 | 1.3623 (11) | C5—H5B | 0.964 (12) |
| N2—C3 | 1.3641 (11) | C6—C6 ⁱ | 1.5140 (19) |
| N2—C4 | 1.4664 (11) | C6—H6A | 0.980 (13) |
| N3—C1 | 1.4324 (13) | C6—H6B | 0.958 (13) |
| C2—C3 | 1.3741 (14) | | |
| C5—O3—C6 | 111.88 (7) | N2—C4—H4A | 105.6 (7) |
| C1—N1—C2 | 104.03 (8) | C5—C4—H4A | 109.1 (7) |
| C1—N2—C3 | 104.96 (8) | N2—C4—H4B | 106.8 (8) |
| C1—N2—C4 | 131.03 (8) | C5—C4—H4B | 111.7 (8) |

| | | | |
|-------------|-------------|--------------------------|-------------|
| C3—N2—C4 | 123.99 (8) | H4A—C4—H4B | 110.5 (11) |
| O1—N3—O2 | 124.07 (10) | O3—C5—C4 | 107.06 (7) |
| O1—N3—C1 | 117.50 (9) | O3—C5—H5A | 109.4 (7) |
| O2—N3—C1 | 118.43 (8) | C4—C5—H5A | 109.8 (7) |
| N1—C1—N2 | 113.79 (8) | O3—C5—H5B | 110.8 (7) |
| N1—C1—N3 | 122.19 (8) | C4—C5—H5B | 111.7 (7) |
| N2—C1—N3 | 124.01 (8) | H5A—C5—H5B | 108.1 (11) |
| N1—C2—C3 | 110.51 (8) | O3—C6—C6 ⁱ | 106.69 (9) |
| N1—C2—H2 | 122.7 (8) | O3—C6—H6A | 109.9 (7) |
| C3—C2—H2 | 126.7 (8) | C6 ⁱ —C6—H6A | 111.4 (7) |
| N2—C3—C2 | 106.70 (8) | O3—C6—H6B | 109.8 (7) |
| N2—C3—H3 | 120.7 (8) | C6 ⁱ —C6—H6B | 109.8 (7) |
| C2—C3—H3 | 132.6 (8) | H6A—C6—H6B | 109.2 (11) |
| N2—C4—C5 | 112.94 (7) | | |
| C2—N1—C1—N2 | -0.18 (11) | C1—N1—C2—C3 | -0.23 (11) |
| C2—N1—C1—N3 | -179.90 (9) | C1—N2—C3—C2 | -0.61 (10) |
| C3—N2—C1—N1 | 0.51 (11) | C4—N2—C3—C2 | 177.86 (8) |
| C4—N2—C1—N1 | -177.81 (8) | N1—C2—C3—N2 | 0.54 (10) |
| C3—N2—C1—N3 | -179.78 (9) | C1—N2—C4—C5 | -77.42 (12) |
| C4—N2—C1—N3 | 1.90 (15) | C3—N2—C4—C5 | 104.54 (10) |
| O1—N3—C1—N1 | -6.49 (16) | C6—O3—C5—C4 | 174.54 (7) |
| O2—N3—C1—N1 | 173.61 (9) | N2—C4—C5—O3 | -72.25 (9) |
| O1—N3—C1—N2 | 173.82 (9) | C5—O3—C6—C6 ⁱ | -179.94 (9) |
| O2—N3—C1—N2 | -6.08 (15) | | |

Symmetry codes: (i) $-x+1, -y+1, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|----------------------------------|------------|-------------|-------------|---------------|
| C4—H4A \cdots O3 ⁱⁱ | 0.977 (13) | 2.404 (13) | 3.3707 (11) | 170.1 (10) |
| C4—H4B \cdots O2 | 0.944 (14) | 2.408 (13) | 2.8169 (13) | 105.9 (9) |

Symmetry codes: (ii) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2




Fig. 3

