

Case Report**Elastofibroma of the Face: A Case Report**Sorena Fardisi ^a, Mohammad Javad Ashraf ^b, Mohammad Reza Zarei ^c, Negar Azarpira ^d, Maryam Raouf ^e, Sara Amanpour ^f^a Postgraduate Student, Dept. of Oral and Maxillofacial Surgery, School of Dentistry, Shiraz University of Medical Sciences, Shiraz, Iran.^b Dept. of Pathology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.^c Dept. of Oral Medicine, School of Dentistry, Kerman University of Medical Sciences, Kerman, Iran.^d Organ Transplant Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.^e Laboratory of Molecular Neuroscience, Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran, Dept. of Biology, Faculty of Sciences, Shahid Bahonar University, Kerman, Iran, Dept. of Endodontics, School of Dentistry, Kerman University of Medical Sciences, Kerman, Iran.^f Dept. of Oral and Maxillofacial Pathology, School of Dentistry, Kerman University of Medical Sciences, Kerman, Iran, Oral and Dental Disease Research Center, Kerman University of Medical Sciences, Kerman, Iran.**KEY WORDS**Elastofibroma;
Face;
Case Report

Received October 2014;

Received in revised form February 2015;

Accepted April 2015.

ABSTRACT

Elastofibroma is a rare neoplasm that characteristically occurs in subscapular area in response to microtrauma. There are some reports of this tumor in other sites of the body but, up till now, there has been no report of elastofibroma in the face. A 20-year-old man presented with a slow growing painless mass in the face without any history of trauma. Histopathologic examination revealed a soft tissue mass composed of eosinophilic fibers admixed with aggregation of fat cells, capillary blood vessels, and fibroblasts. Elastic stain and Masson's trichrome stain confirmed the nature of elastic and collagen fibers. It was a case of elastofibroma in the face.

Corresponding Author: Amanpour S., Dept. of Oral and Maxillofacial Pathology, School of Dentistry, Shafa street, Kerman, Iran. Tel: +98-9177048602 Fax: +98-3432118073
Email: Saraamanpour@gmail.com

Cite this article as: Fardisi S., Ashraf MJ., Zarei MR., Azarpira N., Raouf M., Amanpour S. Elastofibroma of the Face: A Case Report. *J Dent Shiraz Univ Med Sci., Spring 2015; 16 (Supplement): 73-75.*

Introduction

Elastofibroma is a rare benign connective tissue neoplasm that mostly occurs in subscapular area of elderly patients, deep in the serratus anterior muscle. [1] There are reports of this tumor arising in other sites such as hand, [2] foot, [3] thigh, [4] gastrointestinal tract, [5-6] neck, [7] and mouth. [8-10] The pathogenesis of elastofibroma is still unknown but it may result from a reactive response to repetitive microtrauma. [7] To the best of our knowledge, elastofibroma has not been reported in the face so far. In this paper, we describe a case of elastofibroma in the face of a 20-year-old man.

Case report

A 20-year-old man was referred for a painless mass in the left parotid area that has been growing slowly for about 3 years. Clinical examination revealed a firm non-tender mass of approximately 2×3 cm² which was easily

movable (Figure 1).



Figure 1: Clinical view; a non-tender, firm, and movable mass in left parotid area

The overlying skin was normal. There was no specific history of trauma to the area of the lesion and the patient did not report a family history of similar lesions. The differential diagnosis included a preauricular or sebaceous cyst and enlarged lymph nodes. Fine needle




Figure 2a: Microscopic picture of elastofibroma, mainly composed of fibers admixed with aggregation of fat cells and capillary blood vessels (Hematoxylin-Eosin, original magnification $\times 100$) **b:** Elastofibroma, (Hematoxylin-Eosin, original magnification $\times 400$)

aspiration of the lesion revealed few benign looking spindle shape cells in favor of a benign fibrous lesion. The mass was completely excised under local anesthesia and histopathologic examination revealed a soft tissue tumor mainly composed of irregular crinkled eosinophilic fibers with corrugated margins and variable shapes and sizes. Aggregation of fat cells, capillary-sized blood vessels and fibroblasts were also observed within the specimen (Figures 2).

A Verhoeff-Van Gieson stain and Masson's trichrome stain identified elastic fibers and abundant collagen fibers, respectively (Figures 3). Based on microscopic findings and special staining, the diagnosis of elastofibroma was made. Up to now, there has been no clinical sign of recurrence during the 3-year follow-up of the patient.

Discussion

Elastofibroma is a benign process involving the sub-

scapular region in most cases. However, isolated lesions have been reported in thigh, lip, deltoid muscle, and stomach. [1] It was first described by Jarvi and Saxen in 1961. [11] The lesion is typically seen in females over the age of 50. [12-13] Frictional trauma has been suggested as the etiopathogenesis of this lesion. [8] A neoplastic etiology is also possible in some cases based on unusual clinical presentations and molecular studies. [14] Surgical removal is the treatment of choice for this lesion and there has been no report of malignant transformation. [9]

Microscopically, collagen bundles alternate with numerous degenerative fibers in irregular shapes are seen which stain strongly with elastic stain that is allied to mucoid materials, fibroblasts, and small collections of mature lipocytes. [15-16] In the case of elastofibroma reported by Manchandu et al., the elastic fibers showed flower-like appearance with serrated borders. [14] Elastic fibers in this tumor seem to be the result of altered




Figure 3a: Elastin stain shows high density of elastic fibers (Verhoeff-Van Gieson stain, original magnification $\times 400$) **b:** Masson-trichrome stain identified collagen fibers within the lesion (Masson-trichrome stain, original magnification $\times 400$)

elastogenesis caused by frictional trauma. [8] However, several cases of elastofibroma have been reported in different locations without having trauma. [17-21] It may be better explained by a genetic predisposition. [8] The relationship between elastofibroma and genetic factors was first described by Nagamine et al. [5] They reported the influence of hereditary factors on one third of 170 patients. [5]

To the best of our knowledge, this is the first case of elastofibroma that has been presented as a facial mass. Thus, this benign soft tissue tumor should be considered in the differential diagnosis of painless soft tissue masses in the face.

Conflict of Interest

None to declare.

References

- [1] Rosai J. Ackermann's surgical pathology. 10th ed. St. Louis MO: CV Mosby Co.; 2011; p.2115-2116.
- [2] Kapff PD, Hocken DB, Simpson RH. Elastofibroma of the hand. *J Bone Joint Surg Br* 1987; 69: 468-469.
- [3] McPherson FC, Norman LS, Truitt CA, Morgan MB. Elastofibroma of the foot: uncommon presentation: a case report and review of the literature. *Foot Ankle Int* 2000; 21: 775-777.
- [4] Nishida A, Uetani M, Okimoto T, Hayashi K, Hirano T. Bilateral elastofibroma of the thighs with concomitant subscapular lesions. *Skeletal Radiol* 2003; 32: 116-118.
- [5] Nagamine N, Nohara Y, Ito E. Elastofibroma in Okinawa. A clinicopathologic study of 170 cases. *Cancer* 1982; 50: 1794-1805.
- [6] Hobbs CM, Burch DM, Sobin LH. Elastosis and elastofibromatous change in the gastrointestinal tract: a clinicopathologic study of 13 cases and a review of the literature. *Am J Clin Pathol* 2004; 122: 232-237.
- [7] Maldjian C, Adam RJ, Maldjian JA, Rudelli R, Bonakdarpour A. Elastofibroma of the neck. *Skeletal Radiol* 2000; 29: 109-111.
- [8] Potter TJ, Summerlin DJ, Rodgers SF. Elastofibroma: the initial report in the oral mucosa. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004; 97: 64-67.
- [9] Nonaka CF, Rêgo DM, Miguel MC, de Souza LB, Pinto LP. Elastofibromatous change of the oral mucosa: case report and literature review. *J Cutan Pathol* 2010; 37: 1067-1071.
- [10] Darling MR, Kutalowski M, MacPherson DG, Jackson-Boeters L, Wysocki GP. Oral elastofibromatous lesions: a review and case series. *Head Neck Pathol* 2011; 5: 254-258.
- [11] Jarvio O, Sasen E. Elastofibroma dorsae. *Acta Pathol Microbiol Scand Suppl* 1961; 51(Suppl 144): 83-84.
- [12] Montijano Huertes C, Chismol Abad J, Pons Soriano A, Seminario Eleta P, Fenollosa Gómez J. Elastofibroma dorsae. Report of five cases and review of the literature. *Acta Orthop Belg* 2002; 68: 417-420.
- [13] Pyne D, Mootoo R, Bhanji A, Amin S. Elastofibroma dorsae. *Ann Rheum Dis* 2002; 61: 278-279.
- [14] Manchandu R, Foote J, Alawi F. Elastofibroma presenting as an oral soft tissue mass. *J Oral Pathol Med* 2008; 37: 125-126.
- [15] Briccoli A, Casadei R, Di Renzo M, Favale L, Bacchini P, Bertoni F. Elastofibroma dorsae. *Surg Today* 2000; 30: 147-152.
- [16] Weiss SW, Goldblum JR. Enzinger and Weiss's soft tissue tumors, 5th ed. St. Louis: Mosby; 2008. p. 286-289.
- [17] De Nictolis M, Goteri G, Campanati G, Prat J. Elastofibrolipoma of the mediastinum. A previously undescribed benign tumor containing abnormal elastic fibers. *Am J Surg Pathol* 1995; 19: 364-367.
- [18] Enjoji M, Sumiyoshi K, Sueyoshi K. Elastofibromatous lesion of the stomach in a patient with elastofibroma dorsae. *Am J Surg Pathol* 1985; 9: 233-237.
- [19] Tsutsumi A, Kawabata K, Taguchi K, Doi K. Elastofibroma of the greater omentum. *Acta Pathol Jpn* 1985; 35: 233-241.
- [20] Sakatani T, Shomori K, Adachi H, Hosoda A, Ito H. Elastofibroma of the sigmoid colon. *Pathol Res Pract* 2000; 196: 205-207.
- [21] Hayashi K, Ohtsuki Y, Sonobe H, Iwata J, Furihata M, Hikita T, et al. Pre-elastofibroma-like colonic polyp: another cause of colonic polyp. *Acta Med Okayama* 1991; 45: 49-53.