



POSTER PRESENTATION

Open Access

# Role of CD4<sup>+</sup>CD25<sup>hi</sup>CD127<sup>lo/-</sup>FoxP3<sup>+</sup> regulatory T lymphocytes in the pathogenesis of Behçet's disease in children

TA Tran<sup>1\*</sup>, S Monteil<sup>2</sup>, A Letierce<sup>3</sup>, B Terrier<sup>2</sup>, G Geri<sup>2</sup>, D Saadoun<sup>4</sup>, I Kone-Paut<sup>1</sup>, B Salomon<sup>5</sup>, M Rosenzweig<sup>2</sup>

From 18th Pediatric Rheumatology European Society (PReS) Congress  
Bruges, Belgium. 14-18 September 2011

## Introduction

Behçet's disease (BD) is an idiopathic multisystem recurrent inflammatory disorder. Physiopathology of BD shows a role of neutrophils and cytotoxic T lymphocytes.

## Our aim

We were to assess the role of regulatory T lymphocytes (Tregs) in the pathogenesis of BD in children.

## Patients and methods

19 patients with active BD (group A) and 8 patients with inactive BD (group B) were compared with 25 healthy controls (group C). Percentages of blood CD4<sup>+</sup>CD127<sup>lo/-</sup>CD25<sup>hi</sup>FoxP3<sup>+</sup> Tregs and other T/B and NK cells subpopulations were analyzed by flow cytometry. The frequency of IL-17A and IFN- $\gamma$  producing T cells was analyzed by flow cytometer from PBMC after 4 hours stimulation with PMA-ionomycin. We measured serum cytokines by Luminex and ELISA. We compared the 3 groups by using the Wilcoxon-Rank-signed test. Values were expressed as mean and median.

## Results

Patients in the 3 groups (A, B, C respectively) were comparable in term of age and sex distribution (median age: 12.8, 9.9 and 9.7; F/M = 1/1). No differences were observed between the 3 groups concerning the absolute number of lymphocytes, CD4<sup>+</sup> T cells and the percentage of total Tregs (median: A: 1.9, B:1.1, C:2.8). Percentages of naïve Treg/memory Treg and markers of Treg function (GITR, LAP, CD152, DR) were also

similar in the 3 groups. However, there was increased CD8<sup>+</sup> T cells count in the BD patients groups compared to healthy controls (A: 552 $\pm$ 361, p=0.18; B: 627 $\pm$ 159, p=0.04, C: 479 $\pm$ 209). The NK cell (CD3-CD16+CD56+) were highest in group C compared to group A (p=0.4) or B (p=0.001). IL-17A secreting CD4<sup>+</sup> T cells were significantly higher in active BD patients (n=6) compared to controls (n=6) (5.3 $\pm$ 2 vs 2.5 $\pm$ 1.47, p=0.043). Serum IL-6 level was significantly higher in BD populations compared to controls subjects (A: 4.3 $\pm$ 1.22 vs C:3 $\pm$ 0.7 pg/ml, p=0.016).

## Conclusion

There is no deficit of Tregs number in BD patients. The high rate of peripheral IL-17 secreting CD4<sup>+</sup> T cells suggests a possible role of Th17 cells in the occurrence of BD attacks. The Tregs functional ability to regulate CD4 and CD8 T cells needs to be studied further.

## Author details

<sup>1</sup>Department of Paediatrics, Pediatric Rheumatology. CEREMAI Bicêtre Hospital, University of Paris Sud., France. <sup>2</sup>Service de Biothérapies/ UPMC CNRS 7211 INSERM 959. La Pitié Salpêtrière University Hospital. Paris, France. <sup>3</sup>Unité de Recherche Clinique Paris Sud. Bicêtre University Hospital. Le Kremlin Bicêtre. France. <sup>4</sup>Department of Internal Medicine. La Pitié Salpêtrière University Hospital, Paris, France. <sup>5</sup>Unité 2. UPMC-CNRS U7087. La Pitié Salpêtrière University Hospital, Paris, France.

Published: 14 September 2011

doi:10.1186/1546-0096-9-S1-P84

Cite this article as: Tran et al.: Role of CD4<sup>+</sup>CD25<sup>hi</sup>CD127<sup>lo/-</sup>FoxP3<sup>+</sup> regulatory T lymphocytes in the pathogenesis of Behçet's disease in children. *Pediatric Rheumatology* 2011 **9**(Suppl 1):P84.

<sup>1</sup>Department of Paediatrics, Pediatric Rheumatology. CEREMAI Bicêtre Hospital, University of Paris Sud., France  
Full list of author information is available at the end of the article