

POSTER PRESENTATION

Open Access

Contribution of Thomsen-Friedenreich antigens to bladder cancer malignancy: Characterization of cell line models

Paulo F Severino^{1,2*}, Mylène A Carrascal¹, Helena Gouveia¹, M Guadalupe Cabral¹, Manuela Correia¹, Nadia Malagolini², Mariella Chiricolo², Fabio Dall'Olio², Paula A Videira¹

From 16th International Charles Heidelberger Symposium on Cancer Research
Coimbra, Portugal. 26–28 September 2010

A particular type of glycans has been associated with cancer, the O-linked glycosidic Thomsen-Friedenreich (TF) antigens -T and Tn, and the sialylated forms sT and sTn. The expression of TF antigens in bladder cancer (BC) has also been reported, although the correlation with prognosis is controversial. Furthermore, the molecular basis underlying their expression and the role played by the enzymes sialyltransferases (STs), responsible for the expression of sialylated forms, are unknown [1].

For a better understanding, we chose to study the cell line models: MCR and HT1376, exhibiting a low cell surface expression of the sTn and sT respectively, that were transduced with “ST6GalNac.I” or “ST3Gal.I” STs cDNA. The resulted transduced cells MCR_{ST6GalNac1} and HT1376_{ST3Gal1} were purified by immune-magnetic sorting for the sTn, and selected clones for the expression of sT.

Our preliminary results show differences concerning to cell adhesion ability and immunogenicity. Namely, dendritic cells (DCs) and MCR_{ST6GalNac1} in co-cultures, results in an increased expression of TGF- β and IL-10, and decreased of TNF- α and IL-6. After adhesion, MCR negative control (MCR_{NC}) induces higher maturation on DCs (higher expression of HLA-DR, CD80, CD83 and CD86). Concerning to phagocytosis, it was seen that HT1376_{ST3Gal1} are apparently more resistant to phagocytosis by macrophages, in respect to negative control (HT1376_{NC}), and MCR_{ST6GalNac1} less resistant to phagocytosis by DCs, in respect to MCR_{NC}.

* Correspondence: paulo.severino@fcm.unl.pt

¹CEDOC, Departamento de Imunologia, Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Lisboa, Portugal

Full list of author information is available at the end of the article

Author details

¹CEDOC, Departamento de Imunologia, Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Lisboa, Portugal. ²Departamento di Patologia Sperimentale, Università di Bologna, Bologna, Italia.

Published: 24 September 2010

Reference

1. Videira PA, Correia M, Malagolini N, Crespo HJ, Ligeiro D, Calais FM, Trindade H, Dall'Olio F: **ST3Gal. I sialyltransferase relevance in bladder cancer tissues and cell line.** *BMC Cancer* 2009, **9**:357.

doi:10.1186/1471-2407-9-357

Cite this article as: Severino et al.: Contribution of Thomsen-Friedenreich antigens to bladder cancer malignancy: Characterization of cell line models. *BMC Proceedings* 2010 **4**(Suppl 2):P48.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

