

Industrialized foods in early infancy: a growing need of nutritional research

Célia Araújo, MSc^{a,b}, Karla Danielly Ribeiro, PhD^b, Patrícia Padrão, PhD^{a,c}, Pedro Moreira, PhD^{a,c,d,*}

A systematic analysis for the global burden of disease study assessing data from 195 countries, identified high intake of sodium, and low intake of whole grains, fruit, vegetables, nuts, and seeds as the main dietary risks factors for deaths and disability-adjusted life-years.¹ In this global overview, consumption of nearly all healthy foods such as fruit, vegetables, and legumes, was suboptimal, whereas the consumption of sugar-sweetened beverages was far higher than the optimal intake. High consumption of red meat, processed meat, and trans-fat were also toward the bottom when ranking the dietary risks for deaths and disability-adjusted life-years for most high-population countries.¹ These findings require an urgent need to implement healthy eating patterns² as soon as possible during the early windows of opportunity for preventing noncommunicable diseases (NCDs) such as the complementary feeding period and the early infancy.³ During these life stages, commercially produced infant foods may be particularly used, and many of them are also industrialized. At the present time, there is a complex and controversial debate about the role of food processing as the primary driver of diet quality.⁴ In general, food processing implies changing raw commodities into ingredients or food substances, after some changes that may be physical, chemical, or microbiological, for example, and the main goals of these transformations may include expanding shelf life, increasing nutritional quality, attending special nutritional requirements, or adding variety and convenience.⁵ Although many industrialized healthful foods are minimal processed, such as fruits, vegetables, nuts, legumes, or eggs, the specific category of ultraprocessed foods has been studied in relation to diet adequacy and to risk factors for NCDs. Monteiro et al^{4,6-8} proposed that the main shaping force of the global food system and the main determinant of poor diets and their related influences on obesity and diet related NCDs was “food processing.” The definitions of ultra-processed foods have been changing and showed variability particularly in the last 10 years.⁴ More recently, Monteiro et al^{9,10} defined ultraprocessed foods in NOVA classification as industrial formulations of several ingredients which, besides salt, sugar, oils and fats, include food substances not commonly used in

culinary preparations, such as hydrolysed protein, modified starches, and hydrogenated or interesterified oils, or additives whose purpose is to imitate sensorial qualities of unprocessed or minimally processed foods and their culinary preparations or to disguise undesirable qualities of the final product, such as colorants, flavorings, nonsugar sweeteners, emulsifiers, humectants, sequestrants, and firming, bulking, defoaming, anticaking, and glazing agents.

The contribution of ultraprocessed foods to energy intake of populations in the higher quintiles of intake may vary from 49% in Brazil, to 76% in Canada, 78% in the UK, and 81% in the USA; in Mediterranean countries highly processed foods may contribute with 70% of the energy, and 50% to 90% of the nutrients intake in Nordic/Central Europe (with the exceptions of vitamin C and beta-carotene).⁴ However, the contribution of processed and ultraprocessed foods in children is also referred as a factor that may decrease the quality of diet in children considering that many foods exhibit unhealthy nutrient profiles, with higher amounts of sugar, *trans* fat, and sodium, compared with unprocessed or minimally processed foods; furthermore, after adjustment for energy intake, they may provide lower amounts of zinc, calcium, and vitamins A, B12, C, and E.¹¹ In infants and young children from Canada, 63% of the food products sold in Canada were reported as having high levels of sodium or sugar contribution to energy content,¹² and in Brazil, the analysis of 2945 processed foods that are usually consumed as snacks by children showed that 21% presented high levels of sodium.¹³ Maalouf et al¹⁴ assessed the sodium, sugar, and fat content of complementary infant and toddler foods sold in USA, and most commercial toddler meals, cereal bars and breakfast pastries, and infant-toddler snacks and desserts have high sodium contents or contain added sugars, suggesting a need for continued public health efforts to support families implementing healthy eating strategies.

The described nutritional characteristics of these products are particularly important considering that (1) nutritional guidelines recommend to limit the intake of added sugars,¹⁵ particularly before 2 years of age,¹⁶ sodium,¹⁷ and saturated and *trans*-fat and (2) habitual diets of children aged 1 to 3 years across different parts of the world may exceed sodium¹⁸ or the energy contribution from sugar and saturated fat.¹⁹⁻²²

Bearing in mind this global health and dietary intake scenarios, it is important to assess across different nations, including those facing a nutrition transition and developed countries, information on the nutritional profiles, and food processing characteristics of industrialized foods and beverages that are specifically labeled for infants, toddlers, and children younger than 3 years, and to help parents and health professionals with the right information in choosing complementary and baby foods in this biological window of opportunity to prevent NCDs. Furthermore, data collection on industrialized baby foods will be useful to monitor nutritional composition, to assess the eventual need of product nutritional reformulation or other measures of quality control, such as consumer-friendly labeling or nutrient profiling tools, and to evaluate the impact of any future strategies aiming to improve industrialized foods for children.

^a Faculty of Nutrition and Food Sciences, University of Porto, Porto, Portugal,

^b Department of Nutrition, Federal University of Rio Grande do Norte, Natal-RN, Brazil, ^c EPIUnit - Epidemiology Research Unit, University of Porto, ^d Research Centre in Physical Activity, Health and Leisure, University of Porto, Porto, Portugal.

* Corresponding author. Faculdade de Ciências da Nutrição e Alimentação da Universidade do Porto, Porto, Portugal.

E-mail address: pedromoreira@fcna.up.pt (Pedro Moreira)

Copyright © 2019 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of PBJ-Associação Porto Biomedical/Porto Biomedical Society. All rights reserved.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Porto Biomed. J. (2019) 4:4(e47)

Received: 23 June 2019 / Accepted: 4 July 2019

<http://dx.doi.org/10.1097/j.pbj.0000000000000047>

References

- [1] Afshin A, Sur PJ, Fay KA, et al. GBD 2017. Diet Collaborators. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2019;393:1958–1972.
- [2] Schulze MB, Martinez-Gonzalez MA, Fung TT, et al. Food based dietary patterns and chronic disease prevention. *BMJ*. 2018;361:k2396.
- [3] Michaelsen KF, Grummer-Strawn L, Begin F. Emerging issues in complementary feeding: global aspects. *Matern Child Nutr*. 2017;13 (suppl):2.
- [4] Gibney MJ. Ultra-processed foods: definitions and policy issues. *Curr Develop Nutr*. 2019; 3:nzy077.
- [5] Jones JM. Food processing: criteria for dietary guidance and public health? *Proc Nutr Soc*. 2019;78:4–18.
- [6] Monteiro CA, Moubarac JC, Levy RB, et al. Household availability of ultra-processed foods and obesity in nineteen European countries. *Public Health Nutr*. 2018;21:18–26.
- [7] Monteiro CA, Cannon G, Moubarac J-C, et al. The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutr*. 2017:1–13.
- [8] Popkin BM, Reardon T. Obesity and the food system transformation in Latin America. *Obes Rev*. 2018;19:1028–1064.
- [9] Martinez Steele E, Raubenheimer D, Simpson SJ, et al. Ultra-processed foods, protein leverage and energy intake in the USA. *Public Health Nutr*. 2018;21:114–124.
- [10] Martinez Steele E, Baraldi LG, Louzada ML, et al. Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study. *BMJ Open*. 2016;6:e009892.
- [11] Cornwell B, Villamor E, Mora-Plazas M, et al. Processed and ultra-processed foods are associated with lower-quality nutrient profiles in children from Colombia. *Public Health Nutr*. 2018;21:142–147.
- [12] Elliott CD. Sweet and salty: nutritional content and analysis of baby and toddler foods. *J Public Health (Oxf)*. 2011;33:63–70.
- [13] Kraemer MV, Oliveira RC, Gonzalez-Chica DA, et al. Sodium content on processed foods for snacks. *Public Health Nutr*. 2016;19:967–975.
- [14] Maalouf J, Cogswell ME, Bates M, et al. Sodium, sugar, and fat content of complementary infant and toddler foods sold in the United States. *Am J Clin Nutr*. 2017;105:1443–1452.
- [15] WHO. Guideline: Sugars Intake for Adults and Children. Geneva: World Health Organization; 2015.
- [16] Vos MB, Kaar JL, Welsh JA, et al. Added sugars and cardiovascular disease risk in children: a scientific statement from the American Heart Association. *Circulation*. 2017;135:e1017–e1034.
- [17] WHO. Guideline: Sodium Intake for Adults and Children. Geneva: World Health Organization (WHO); 2012.
- [18] Pereira-da-Silva L, Rego C, Pietrobelli A. The diet of preschool children in the Mediterranean countries of the European Union: a systematic review. *Int J Environ Res Public Health*. 2016;13:6.
- [19] Syrad H, Llewellyn CH, Van Jaarsveld CH, et al. Energy and nutrient intakes of young children in the UK: findings from the Gemini twin cohort. *Br J Nutr*. 2016;115:1843–1850.
- [20] Bailey RL, Fulgoni VL, Cowan AE, et al. Sources of added sugars in young children, adolescents, and adults with low and high intakes of added sugars. *Nutrients*. 2018;10:1.
- [21] Fujiwara A, Murakami K, Asakura K, et al. Estimation of starch and sugar intake in a Japanese population based on a newly developed food composition database. *Nutrients*. 2018;10:10.
- [22] Carvalho CA, Fonseca PC, Priore SE, et al. Food consumption and nutritional adequacy in Brazilian children: a systematic review [in Portuguese]. *Rev Paul Pediatr*. 2015;33:211–221.