


## RESEARCH ARTICLE

# Clinical analysis of pregnant women with 2019 novel coronavirus pneumonia

Siyu Chen<sup>1</sup>  | E. Liao<sup>2</sup> | Dongmei Cao<sup>2</sup> | Ying Gao<sup>2</sup> | Guoqiang Sun<sup>2</sup> | Yong Shao<sup>1</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China

<sup>2</sup>Department of Obstetrics and Gynecology, Maternal and Child Hospital of Hubei Province, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China

**Correspondence**

Guoqiang Sun, MD, Department of Obstetrics and Gynecology, Maternal and Child Hospital of Hubei Province, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China.  
Email: sun821ch@126.com

Yong Shao, PhD, MD, Department of Obstetrics and Gynecology, The First Affiliated Hospital of Chongqing Medical University, 400016 Chongqing, China.  
Email: cqshaoyong@163.com

**Funding information**

Chongqing Municipal Public Health Bureau, Chongqing People's Municipal Government, Grant/Award Numbers: 2015XMSB0001298, 2019ZDXM055; National Natural Science Foundation of China, Grant/Award Number: 81471473

**Abstract**

The aim is to evaluate pregnant women infected with coronavirus disease 2019 (COVID-19) and provide help for clinical prevention and treatment. All five cases of pregnant women confirmed COVID-19 were collected among patients who admitted to the Maternal and Child Hospital of Hubei Province between January 20 and February 10, 2020. All patients, aging from 25 to 31 years old, had the gestational week from 38th weeks to 41st weeks. All pregnant women did not have an antepartum fever but developed a low-grade fever (37.5°C–38.5°C) within 24 hours after delivery. All patients had normal liver and renal function, two patients had elevated plasma levels of the myocardial enzyme. Unusual chest imaging manifestations, featured with ground-glass opacity, were frequently observed in bilateral (three cases) or unilateral lobe (two cases) by computed tomography (CT) scan. All labors smoothly processed, the Apgar scores were 10 points 1 and 5 minutes after delivery, no complications were observed in the newborn. Pregnancy and perinatal outcomes of patients with COVID-19 should receive more attention. It is probable that pregnant women diagnosed with COVID-19 have no fever before delivery. Their primary initial manifestations were merely low-grade postpartum fever or mild respiratory symptoms. Therefore, the protective measures are necessary on admission; the instant CT scan and real-time reverse-transcriptase polymerase-chain-reaction assay should be helpful in early diagnosis and avoid cross-infection on the occasion that patients have fever and other respiratory signs.

**KEYWORDS**

clinical manifestations, COVID-19, labor, pregnancy

## 1 | INTRODUCTION

In December 2019, a novel coronavirus (SARS-CoV-2)-induced pneumonia, started in Wuhan and rapidly spread in China and other countries, is causing a global health problem.<sup>1,2</sup> As February 8, 2020, the National Health Commission announced 37 198 confirmed cases and 28 942 suspected cases in China. Pregnant women were also reported being infected with coronavirus disease 2019 (COVID-19), which possesses similarity

with the severe acute respiratory syndrome (SARS). As reported by the previous SARS-related studies, pregnant women were more susceptible to coronavirus infection and at a high risk of poor perinatal outcomes.<sup>3,4</sup> Nevertheless, studies on COVID-19 are still quite limited, leading to a great challenge for pregnant women care. Our study aimed to describe the clinical characteristics of pregnant women infected with COVID-19, and we hope this study could provide a guidance for clinical prevention and treatment of COVID-19 in pregnant women.

## 2 | MATERIALS AND METHODS

### 2.1 | Data sources

Our study collected a total of five pregnant women with COVID-19, who were hospitalized for regular delivery in Maternal and Child Hospital of Hubei Province between January 20 and February 10, 2020 (Table 1). All patients were regularly given a prenatal routine examination that started from the first trimester. Ultrasound examination and fetal heart monitoring were offered regularly before delivery, no symptoms of viral pneumonia were observed. On latest admission, all patients received routine fetal heart monitoring and laboratory examinations including complete blood count, blood chemistry, coagulation test, liver and renal function, electrolytes, C-reactive protein, procalcitonin, lactate dehydrogenase, and creatine kinase. The delivery strategy is determined according to the pregnant women's general situation. Apgar scoring and nursing was routinely given for the newborn after delivery. Once the patient had suspected symptoms, chest computed tomography (CT) scan and other blood analysis were instantly performed, antibiotics were empirically given in time to prevent bacterium-related infection, patients were quarantined for an intensive treatment, and newborns were closely monitored with breastfeeding restriction. This study was approved by the ethical committee of Maternal and Child Hospital of Hubei Province, the work undertaken in this study conforms to the provisions of the Declaration of Helsinki (as revised in Tokyo 2004). Informed consent was obtained from all pregnant women.

### 2.2 | Diagnostic criteria

Cases were suspected based on the WHO interim guidance and the definition of COVID-19 issued by the National Health Commission of China.<sup>5</sup> (1) Severe acute respiratory infection that required hospitalized treatment, (2) with no other etiology that thoroughly explains the clinical presentation, and (3) a history of travel to or residence in the city of Wuhan, or contact with the environment of COVID-19 as a healthcare worker. Abnormal results included the aforementioned imaging characteristics of pneumonia, decreased white blood cell (WBC) or lymphocyte counts in the early stage of the disease.

The diagnostic criteria for COVID-19 are based on the criteria for the suspected cases. Positive results from real-time reverse transcription polymerase chain reaction (RT-PCR) assay with samples like sputum, throat swabs, lower respiratory tract secretions or other specimens, or sequencing-confirmed SARS-CoV-2 infection. All patients in this study were reported positive for SARS-CoV-2 infection through quantitative real-time RT-PCR tests.

## 3 | RESULTS

### 3.1 | Demographic characteristics

According to the demographic data of five patients (Table 1), they were between the ages of 25 and 31 years old. None was a healthcare worker. All of them were physically fit and conceived naturally. The gestational weeks on admission were the third trimester (38th-41st weeks). Patients 1 and 5 had gestational

**TABLE 1** Maternal characteristics and symptoms of pregnant women with coronavirus disease 2019 (COVID-19)

	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5
Age, y	29	30	25	31	29
Date of admission	23/01/2020	26/01/2020	25/01/2020	27/01/2020	26/01/2020
Date of delivery	24/01/2020 17:51	26/01/2020 22:55	25/01/2020 09:17	27/01/2020 21:30	27/01/2020 09:00
Date of diagnosis	27/01/2020 14:30	27/01/2020 09:50	25/01/2020 23:43	28/01/2020 02:36	28/01/2020 08:31
Place of residence	Wuhan, Hubei	Wuhan, Hubei	Wuhan, Hubei	Wuhan, Hubei	Wuhan, Hubei
Obstetrical complications	Gestational diabetes	Pre-eclampsia	None	None	Gestational diabetes
Post-partum fever, °C	38.5	37.5	37.4	37.4	37.8
Cough	–	–	+	–	–
Sputum	–	–	–	–	+
Coryza	–	–	–	–	+
Myalgia	–	–	–	–	–
Malaise	–	–	–	–	–
Dyspnea	–	–	–	–	–
Other symptoms	–	–	–	–	–
Systolic blood Pressure, mm Hg	125	153	104	95	119

diabetes and patient 2 had pre-eclampsia before delivery. All patients lived in Wuhan, Hubei province.

On admission, all patients had no fever or cough before delivery. However, all patients' low-grade fever (37.5°C-38.5°C), the most common onset symptom of COVID-19, occurred after delivery within 24 hours, and the highest body temperature of patient 1 was 38.5°C. Patient 3 had a cough 1 day before giving birth, and patient 5 had the symptoms of coughing and nasal runny 10 days before delivery. All patients had no symptoms such as hemoptysis, dyspnea, shortness of breath, nausea, and vomiting. All oxygen saturation tests were normal.

### 3.2 | Laboratory findings

Laboratory findings showed (Table 2) that in the early stage of the disease, the count of WBCs in all patients was normal, and two patients developed an increased WBC count 24 hours after the symptom onset. The WBC counts of patient 5 were always abnormally high during


hospital admission. C-reactive protein levels increased in all patients but except patient 4, particularly in patient 5 who had a substantial increase (40.98-125.90 mg/L). There was no obvious abnormality of the procalcitonin levels observed in all patients. The levels of albumin were decreased in all patients, whereas the levels of alanine aminotransferase, aspartate aminotransferase, and total bilirubin were normal. All patients except patient 1 presented with the elevated levels of alkaline phosphatase. CK was reduced in patient 1 (21 U/L) while increased in patient 3 (227 U/L), respectively.

All the serum samples of patients were tested for seven common respiratory pathogens (*Chlamydia pneumoniae*, *Mycoplasma pneumoniae*, adenovirus, parainfluenza virus, respiratory syncytial virus, influenza a virus, and influenza b virus) and N7 subtype avian influenza virus antigen. All the tests were negative except for the positive results of *Mycoplasma pneumoniae* in patients 3 and 5. Unusual chest imaging manifestations, featured with ground-glass opacity, were frequently observed in bilateral (three cases) or unilateral lobe (two cases) by CT scan (Figure 1). All COVID-19 were laboratory-confirmed by the RT-PCR method.

**TABLE 2** Laboratory characteristics of pregnant women with coronavirus disease 2019 (COVID-19)

Laboratory characteristics	Normal range	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5
White blood cell count (WBC), $\times 10^9/L$	3.5-9.5	6.77	10.78 $\uparrow$	9.31	16.27 $\uparrow$	12.97 $\uparrow$
Lymphocyte count (LYMPH), $\times 10^9/L$	1.1-3.2	1.05 $\downarrow$	0.77 $\downarrow$	0.91 $\downarrow$	0.97 $\downarrow$	2.48
Neutrophil cell count (NEUT), $\times 10^9/L$	1.8-6.3	5.28	9.44 $\uparrow$	7.91 $\uparrow$	15.52 $\uparrow$	9.54 $\uparrow$
Platelet count (PLT), $\times 10^9/L$	125-350	240	167	105	194	326
C-reactive protein (CRP), mg/L	<4	18.46 $\uparrow$	19.56 $\uparrow$	NA	8.96 $\uparrow$	82.14 $\uparrow$
Procalcitonin (PCT), ng/mL	<0.05	0.053	0.212	0.070	0.048	0.122
Hemoglobin (HGB), g/L	115-150	125	119	113	103 $\downarrow$	105 $\downarrow$
Prothrombin time (PT), s	9.4-12.5	10.3	9.9	11.1	11.4	12.1
Activated partial thromboplastin time (APTT), s	25.1-36.5	25.7	29.5	28.0	29.1	30.1
D-dimer (D-DI), $\mu g/mL$	<0.5	0.54 $\uparrow$	NA	NA	1.90 $\uparrow$	1.64 $\uparrow$
Albumin (ALB), g/L	40-55	35.5 $\downarrow$	31.3 $\downarrow$	31.8 $\downarrow$	29.4 $\downarrow$	28.3 $\downarrow$
Total bilirubin (TBIL), $\mu g/mL$	5-21	5.5	4.0	6.2	7.3	8.1
Alanine aminotransferase (ALT), U/L	0-35	14.7	15.5	7.6	5.9	7.6
Aspartate aminotransferase (AST), U/L	0-35	13.8	30.3	26.1	14.5	16.4
Alkaline phosphatase (ALP), U/L	35-100	100.0	145.0 $\uparrow$	234.0 $\uparrow$	266 $\uparrow$	146.0 $\uparrow$
Creatinine (CREA), $\mu mol/L$	41-73	62.0	58.0	38.4 $\downarrow$	56.0	51.0
Sodium (Na), mmol/L	137-147	143	138.0	133.0 $\downarrow$	140.0	143.0
Potassium (K), mmol/L	3.5-5.3	4.36	4.78	4.40	4.59	3.50
Calcium (Ca), mmol/L	2.11-2.52	2.13	1.91 $\downarrow$	1.97 $\downarrow$	2.25	2.13
Chloride (Cl), mmol/L	99-110	105.0	100.0	104.0	104.0	104.0
Lactose dehydrogenase (LDH), U/L	120-250	245.0	412.0 $\uparrow$	247.0	221.0	194.0
Creatinine kinase (CK), U/L	40-200	21.0 $\downarrow$	105.0	227.0 $\uparrow$	69.0	199.0
SARS-CoV-2 quantitative RT-PCR	—	+	+	+	+	+
Seven common respiratory pathogens	—	—	—	—	—	—

Note: Positive, —; negative, +; NA, not available.



**FIGURE 1** Chest computed tomography (CT) image of the pregnant women with coronavirus disease 2019 (COVID-19). Chest CT images of patient 1 showed ground-glass opacity in left lung on day 1 after symptom onset. Chest CT images of patient 2 showed ground-glass opacity in both lungs on day 1 after symptom onset. Chest CT images of patient 3 showed ground-glass opacity in the right lung on day 1 after symptom onset. Chest CT images of patient 4 showed ground-glass opacity in the right lung on day 1 after symptom onset. Chest CT images of patient 5 showed ground-glass opacity in both lungs on day 1 after symptom onset

### 3.3 | Perinatal outcomes

The perinatal outcomes and newborns are described in Table 3. Patient 2 underwent an emergent cesarean section due to the fetal tachycardia during pregnancy, patient 5 underwent elective cesarean section due to gestational diabetes. Patients 1, 3, and 4 had a natural delivery. All patients had a smooth labor process, and no partial complications occurred. The weights of newborns ranged from 3235 and 4050 g. Apgar scores were 10 points 1 and 5 minutes after delivery. No newborns showed the signs of perinatal COVID-19 infection, umbilical cord blood and amniotic fluid were not applied for virus detection due to the lack of reagent. No complications

of placenta infarction and chorionic amniotic inflammation were reported. All patients were advised to stop breastfeeding and empirically given oseltamivir and azithromycin for treatment.

## 4 | DISCUSSION

A study that analyzed 12 cases of pregnant women with SARS in 2004, Hong Kong, found that pregnant women with SARS were associated with high morbidity and mortality.<sup>6</sup> Alfaraj et al<sup>7</sup> reported that the neonatal mortality rate was up to 27% for pregnant women with Middle East respiratory syndrome (MERS). Pregnant women

**TABLE 3** Perinatal outcomes of pregnant women with COVID-19

	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5
Gestational age on delivery, wk	40 <sup>+4</sup>	39 <sup>+1</sup>	38 <sup>+6</sup>	39 <sup>+6</sup>	39
Gravidity (n) Parity (n)	G1P0	G1P0	G3P1	G3P0	G1P0
Fetal heart rate trace	Normal	Fetal tachycardia	Normal	Normal	Normal
Mode of delivery	Natural birth	C-section	Natural birth	Natural birth	C-section
Complications after delivery	No	No	No	No	No
Weights of newborns, g	3235	3800	3670	3700	4050
Apgar scores (5 min/10 min)	10/10	10/10	10/10	10/10	10/10
Neonatal symptoms infected with COVID-19	No	No	No	No	No
Neonatal SARS-CoV-2 quantitative RT-PCR	Negative	Negative	Negative	Negative	Negative

Abbreviations: COVID-19, coronavirus disease 2019; RT-PCR, reverse transcription polymerase chain reaction; SARS, severe acute respiratory syndrome.

infected with influenza showed a higher risk of hospitalizing than healthy pregnant women.<sup>8</sup> Therefore, during the outbreak of COVID-19, a disease that still lacks a specific and effective treatment, the prevention and management of pregnant women and newborns should be emphasized during the peripartur period.

The data in our study showed that all pregnant women had no fever before delivery, but after delivery two patients developed coughing and low-grade fever. Their CT scan results demonstrated the typical images of viral pneumonia. All five patients were confirmed with COVID-19 by the test of nucleic acid examination. The fetal tachycardia was observed in one pregnant woman who subsequently received an emergent cesarean section. Chen et al<sup>9</sup> reported that pregnancy with pneumonia would have a higher risk of cesarean section, premature delivery, decreased Apgar score of the newborn, and low birth weight of the newborn, etc. Even patients with COVID-19 may only have the mild respiratory symptoms during pregnancy, they are still at a high risk of severe pneumonia and adverse pregnancy outcomes, especially for patients who have pre-eclampsia or other complications, as pneumonia may aggravate pulmonary edema and oxygen saturation reduction.

Our study showed that the strategy of delivery was not affected by the coexistence of COVID-19, the choice of cesarean section was mainly determined by obstetric factors. According to the previous research on SARS, the peritoneal fluid<sup>10</sup> and faeces<sup>11</sup> could be positive for virus detection, and the virus might be transmitted during vaginal and cesarean delivery. As reported by a recent newspaper, a newborn was diagnosed as COVID-19 30 hours after delivery, which suggested that there may be a risk of virus intrauterine vertical transmission or infection through contact with the maternal secretion. However, another research including nine pregnant women showed no direct evidence of intrauterine infection,<sup>12</sup> which was consistent with the research on pregnant women with SARS,<sup>13</sup> therefore, we need to be alert to take preventive and treatment measures for newborns in such period. According to China experts' consensus, it is recommended to quarantine the infants whose mother is suspected or diagnosed with COVID-19 for 14 days. Currently, there is not enough evidence supporting the presence of SARS-CoV-2 in breast milk, however, breastfeeding is not recommended according to the experience from SARS.<sup>4</sup>

Owing to the absence of symptoms before delivery, all patients neglected CT scan and nucleic acid screening on admission, like most Chinese pregnant women usually do. It is probable to miss the diagnosis of COVID-19, which could delay the diagnosis and treatment of patients, and increase the risk of transmission for surrounding people, containing healthcare workers. The pregnant women's exposure to CT radiation barely reaches the dose that causes harm to the fetus,<sup>14</sup> so it is plausible to perform a COVID-19 screening during early pregnancy. In terms of treatment, pregnant women with MERS were not recommended to use ribavirin because of its risk to lead to fetal deformities.<sup>15</sup> New drugs for the treatment of COVID-19 are still being tested in clinical trials,<sup>16</sup> and the effects for pregnant women are still not definitive and require further study.<sup>12</sup>

## 5 | CONCLUSION

Pregnant women infected with COVID-19 need to receive more intensive attention. COVID-19 could asymptotically occur during gestation but get diagnosed after delivery. The manifestations include postpartum fever, mild respiratory symptoms, and typical CT images. Therefore, the protective measures for COVID-19 are necessary on admission; The CT scan and real-time RT-PCR assay could be helpful for the prevention of cross-transmission and early treatment of pregnant women with COVID-19.

## ACKNOWLEDGMENTS

This study is funded by the National Natural Sciences Foundation of China (No. 81471473) and Chongqing Municipal Public Health Bureau, Chongqing People's Municipal Government (Nos. 2015XMSB0001298 and 2019ZDXM055).

## CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

## AUTHOR CONTRIBUTIONS

The corresponding author GS and YS developed the study concept and design, critically reviewed the data and decided the content of the manuscript. All other authors acquired and/or analyzed data, performed the statistical analysis and/or provided technical or material support and/or wrote and/or critically reviewed the manuscript. Authors SC, EL and DC contributed equally to this study. All authors finally decided on the content of the manuscript.

## ORCID

Siyu Chen  <http://orcid.org/0000-0002-5734-2015>

## REFERENCES

1. The 2019-nCoV Outbreak Joint Field Epidemiology Investigation Team QL. An outbreak of NCIP (2019-nCoV) Infection in China—Wuhan, Hubei Province, 2019-2020. *China CDC Weekly*. 2020;2(5):79-80.
2. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507-513.
3. Tang P, Wang J, Song Y. Characteristics and pregnancy outcomes of patients with severe pneumonia complicating pregnancy: a retrospective study of 12 cases and a literature review. *BMC Pregnancy Childbirth*. 2018;18(1):434.
4. Maxwell C, McGeer A, Tai KFY, Sermer M Maternal Fetal Medicine Committee, Infectious Disease Committee. Management guidelines for obstetric patients and neonates born to mothers with suspected or probable severe acute respiratory syndrome (SARS). *J Obstet Gynaecol Can*. 2009;31(4):358-364.
5. China tNHCo. Interpretation of pneumonia diagnosis and treatment scheme for novel coronavirus infection (trial version 5). 2020; <http://www.nhc.gov.cn/yzygj/s7652m/202002/e84bd30142ab4d8982326326e4db22ea.shtml>
6. Wong SF, Chow KM, Leung TN, et al. Pregnancy and perinatal outcomes of women with severe acute respiratory syndrome. *Am J Obstet Gynecol*. 2004;191(1):292-297.
7. Alfaraj SH, Al-Tawfiq JA, Memish ZA. Middle East respiratory syndrome coronavirus (MERS-CoV) infection during pregnancy:

- report of two cases & review of the literature. *J Microbiol Immunol Infect.* 2019;52(3):501-503.
8. Mertz D, Geraci J, Winkup J, Gessner BD, Ortiz JR, Loeb M. Pregnancy as a risk factor for severe outcomes from influenza virus infection: A systematic review and meta-analysis of observational studies. *Vaccine.* 2017;35(4):521-528.
  9. Chen YH, Keller J, Wang IT, Lin CC, Lin HC. Pneumonia and pregnancy outcomes: a nationwide population-based study. *Am J Obstet Gynecol.* 2012;207(4):288 e281-287.
  10. Shek CC, Ng PC, Fung GPG, et al. Infants born to mothers with severe acute respiratory syndrome. *Pediatrics.* 2003;112(4):e254.
  11. Isakbaeva ET, Khetsuriani N, Beard RS, et al. SARS-associated coronavirus transmission, United States. *Emerg Infect Dis.* 2004;10(2):225-231.
  12. Chen H, Guo J, Wang C, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet.* 2020; 395(10226):809-815.
  13. Robertson CA, Lowther SA, Birch T, et al. SARS and pregnancy: a case report. *Emerg Infect Dis.* 2004;10(2):345-348.
  14. American College of Obstetricians and Gynecologists' Committee on Obstetric Practice. Committee Opinion No. 656: guidelines for diagnostic imaging during pregnancy and lactation. *Obstet Gynecol.* 2016; 127(2):e75-e80.
  15. Alserehi H, Wali G, Alshukairi A, Alraddadi B. Impact of Middle East respiratory syndrome coronavirus (MERS-CoV) on pregnancy and perinatal outcome. *BMC Infect Dis.* 2016;16:105.
  16. Guan W, Ni Z, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020:NEJMoa2002032.

**How to cite this article:** Chen S, Liao E, Cao D, Gao Y, Sun G, Shao Y. Clinical analysis of pregnant women with 2019 novel coronavirus pneumonia. *J Med Virol.* 2020;92:1556-1561.  
<https://doi.org/10.1002/jmv.25789>