

Ocrelizumab Treatment Modulates B-Cell Regulating Factors in Multiple Sclerosis

Samantha Ho, MSc, Eva Oswald, BSc, Hoi Kiu Wong, MSc, Atay Vural, MD, PhD, Vuslat Yilmaz, PhD, Erdem Tüzün, MD, Recai Türkoğlu, MD, Tobias Straub, MD, Ingrid Meinl, MD, Franziska Thaler, MD, Tania Kümpfel, MD, Edgar Meinl, MD,* and Simone Mader, PhD*

Neurol Neuroimmunol Neuroinflamm 2023;10:e200083. doi:10.1212/NXI.000000000200083

Correspondence

Dr. Meinl
Edgar.Meinl@med.uni-muenchen.de
or Dr. Mader
Simone.Mader@med.uni-muenchen.de

Abstract

Background and Objectives

Antibodies to CD20 efficiently reduce new relapses in multiple sclerosis (MS), and ocrelizumab has been shown to be effective also in primary progressive MS. Although anti-CD20 treatments efficiently deplete B cells in blood, some B cells and CD20⁻ plasma cells persist in lymphatic organs and the inflamed CNS; their survival is regulated by the B cell-activating factor (BAFF)/A proliferation-inducing ligand (APRIL) system. The administration of a soluble receptor for BAFF and APRIL, atacicept, unexpectedly worsened MS. Here, we explored the long-term effects of ocrelizumab on immune cell subsets as well as on cytokines and endogenous soluble receptors comprising the BAFF-APRIL system.

Methods

We analyzed immune cell subsets and B cell-regulating factors longitudinally for up to 2.5 years in patients with MS treated with ocrelizumab. In a second cohort, we determined B-cell regulatory factors in the CSF before and after ocrelizumab. We quantified the cytokines BAFF and APRIL along with their endogenous soluble receptors soluble B-cell maturation antigen (sBCMA) and soluble transmembrane activator and calcium-modulator and cyclophilin ligand (CAML) interactor (sTACI) using enzyme-linked immunosorbent assays (ELISAs). In addition, we established an in-house ELISA to measure sTACI-BAFF complexes.

Results

Ocrelizumab treatment of people with MS persistently depleted B cells and CD20⁺ T cells. This treatment enhanced BAFF and reduced the free endogenous soluble receptor and decoy sTACI in both serum and CSF. Levels of sTACI negatively correlated with BAFF levels. Reduction of sTACI was associated with formation of sTACI-BAFF complexes.

Discussion

We describe a novel effect of anti-CD20 therapy on the BAFF-APRIL system, namely reduction of sTACI. Because sTACI is a decoy for APRIL, its reduction may enhance local APRIL activity, thereby promoting regulatory IgA⁺ plasma cells and astrocytic interleukin (IL)-10 production. Thus, reducing sTACI might contribute to the beneficial effect of anti-CD20 as exogenous sTACI (atacicept) worsened MS.

Classification of Evidence

This study provides Class IV evidence that endogenous sTACI in blood and CSF is decreased after ocrelizumab treatment.

*These authors contributed equally to this work and are co-senior authors and shared corresponding authors.

From the Institute of Clinical Neuroimmunology (S.H., E.O., H.K.W., A.V., I.M., F.T., T.K., E.M., S.M.), Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München; Graduate School of Systemic Neurosciences (S.H.), Ludwig-Maximilians-Universität München, Germany; Department of Neurology (A.V.), Koc University School of Medicine; Department of Neuroscience (V.Y., E.T.), Aziz Sancar Institute of Experimental Medicine, Istanbul University; Department of Neurology (R.T.), Haydarpaşa Numune Education and Research Hospital, Istanbul, Türkiye; Core Facility Bioinformatics (T.S.), Biomedical Center, Ludwig-Maximilians-Universität München, Germany; Munich Cluster for Systems Neurology (SyNergy) (F.T.), Germany.

Go to [Neurology.org/NN](https://www.neurology.org/NN) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

MORE ONLINE

Class of Evidence

Criteria for rating therapeutic and diagnostic studies

[NPub.org/coe](https://www.npub.org/coe)

Glossary


APRIL = A proliferation-inducing ligand; **BAFF** = B cell-activating factor; **BAFF-R** = BAFF receptor; **BCMA** = B-cell maturation antigen; **BL** = baseline; **CAML** = calcium-modulator and cyclophilin ligand; **EAE** = experimental autoimmune encephalomyelitis; **ELISA** = enzyme-linked immunosorbent assay; **IL** = interleukin; **MS** = multiple sclerosis; **sTACI** = soluble transmembrane activator and CAML interactor; **sBCMA** = soluble B-cell maturation antigen; **TACI** = transmembrane activator and CAML interactor; **TP1** = time point 1.

Anti-CD20 mAbs are highly efficient in treating relapsing-remitting multiple sclerosis (MS), and the anti-CD20 mAb ocrelizumab is the first approved treatment that slows disability progression in primary progressive MS.^{1,2} Anti-CD20 treatment largely depletes circulating B cells and a subset of CD20⁺ T cells.¹⁻⁷

In contrast to the almost complete depletion of CD20⁺ cells in blood, there is evidence that CD20⁺ B cells in the lymphatic organs and the inflamed CNS are not eliminated to the same extent during anti-CD20 treatment.^{8,9} Furthermore, CD20⁻ plasma cells persist in their survival niches in the bone marrow and the inflamed CNS,¹ and the generation of mucosal IgA⁺ plasmablasts continues despite anti-CD20

treatment.¹⁰ The survival of B cells and plasma cells persisting during anti-CD20 is regulated by the B cell-activating factor (BAFF)/A proliferation-inducing ligand (APRIL) system, which comprises the ligands BAFF and APRIL and the receptors B-cell maturation antigen (BCMA), transmembrane activator and calcium-modulator and cyclophilin ligand (CAML) interactor (TACI), and BAFF receptor (BAFF-R).¹¹ We found previously that the membrane-bound receptors TACI and BCMA are shed from B cells and plasma cells yielding the endogenous soluble receptors, soluble B-cell maturation antigen (sBCMA)¹² and soluble transmembrane activator and CAML interactor (sTACI),¹³ reviewed in Ref. 14; these are elevated in the CSF of patients with MS and function as decoys.^{12,13} Figure 1 illustrates

Figure 1 Overview of the BAFF/APRIL System



sources of ligands, soluble receptors, and ligand binding. During anti-CD20 treatment, the interplay between BAFF and APRIL with their soluble receptors regulates the maintenance of remaining B cells and plasma cells. B-cell depletion with anti-CD20 results in elevated BAFF levels in serum,¹⁵ presumably due to reduced consumption,¹⁶ but the influence of anti-CD20 on the soluble receptors is unknown. The strong effect of soluble receptors of the BAFF-APRIL system on MS is evident from unexpected clinical observations: Application of exogenous soluble TACI, atacicept, which decreases B-cell numbers, unexpectedly increased MS activity¹⁷ and enhanced the conversion of optic neuritis to MS.¹⁸ The reasons for the apparently paradoxical effects of atacicept on MS are still not completely understood but may involve regulatory effects of B cells (reviewed in Ref. 19,20) and inhibitory effects mediated via receptors for BAFF and APRIL.²¹⁻²³ In the present study, we explored the long-term effects of ocrelizumab on immune cell subsets and on the BAFF-APRIL system focusing on the endogenous soluble receptors sBCMA and sTACI and complexes with their ligands.

Methods

Patients

First Cohort

From 36 patients with MS (summary in Table 1; individual details in eTable 1, links.lww.com/NXI/A799), blood was obtained before the first administration of ocrelizumab half-dose 300 mg infusion (baseline [BL]), before the second half-dose infusion 2 weeks apart (time point 1 [TP1]), and before every full-dose infusion 600 mg cycle with 6-month intervals (TP2–TP6) for up to 2.5 years. From this cohort, all patients were included in the analysis of immune cell subsets, and we randomly selected 17 of them for our enzyme-linked immunosorbent assay (ELISA) series.

Second Cohort

From 19 additional patients with MS (summary in Table 1; individual details in eTable 2, links.lww.com/NXI/A799), CSF and serum were obtained before and 12–19 months after ocrelizumab therapy. The patients received ocrelizumab at 6-month intervals, and CSF was obtained between 2 infusions.

Flow Cytometry

Immunophenotyping using anti-CD45 (clone ZD1), anti-CD20 (clone L27), anti-CD3 (clone SK7), and anti-CD19 (clone SJ25C1) was performed in cohort 1 as described.⁴

Measurement of Soluble Components of the BAFF-APRIL System

BAFF (DY124), APRIL (DY884), sBCMA (DY193), and sTACI (DY174) were analyzed in serum and CSF by ELISAs (R&D Systems, Minneapolis, MN). To determine whether these ELISAs detect bound or unbound forms of the proteins, we performed each of the ELISAs with a stable physiologic concentration of the protein of interest and added a titration of the interaction partner (including the physiologic concentration) to determine whether the complex formation hinders the detection.

Because we found that the detection of sTACI was blocked by addition of BAFF, we developed an ELISA to quantify the complexes sTACI-BAFF. To this end, we used the capture antibody from the BAFF ELISA kit (841899; DY124) for coating and the detection antibody from the sTACI ELISA kit (841862; DY174) for detection. The quantity of sTACI-BAFF complexes was determined by the quantity of detected sTACI present in the wells. In other words, the quantitative standard of the sTACI-BAFF complexes was conducted as in the sTACI ELISA (DY174). We calculated the total amount of sTACI as sTACI detected in the sTACI ELISA + sTACI detected in sTACI-BAFF complexes. We then calculated the percentage of sTACI in complex with BAFF with this

Table 1 Summary of Patients' Characteristics

	Patients included in the study ^a	Patients with ELISA analysis ^b	Age at baseline (y)	Sex	Diagnosis	Disease duration at baseline (mo)	Time between the last therapy and the start of ocrelizumab (wk)
Cohort 1	n = 36	n = 17	41 (23–62)	25 F 11 M	26 RRMS 8 PPMS 2 SPMS	139 (1–420)	29 (0–574)
Cohort 2	n = 19	n = 19	47 (27–68)	14 F 5 M	16 RRMS 0 PPMS 3 SPMS	149 (64–246)	11 (3–24)

Abbreviations: MS = multiple sclerosis; PPMS = primary progressive MS; RRMS = relapsing-remitting multiple sclerosis; SPMS = secondary progressive MS.
^a First cohort: 36 patients with MS without a previous depleting treatment were recruited from the Ludwig-Maximilians-Universität München Klinikum, Munich, Germany. All patients were included in immune cell phenotype analyses. Patients were analyzed longitudinally from baseline (before initiating ocrelizumab treatment) up to 2.5 years of Ocrelizumab treatment (8 patients were observed for 6–12 months, 5 patients up to 1.5 years, 10 patients up to 2 years, and 13 patients up to 2.5 years after baseline). Second cohort: 19 patients with MS were recruited at baseline and follow-up (12–19 months after the first ocrelizumab infusion from the Haydarpaşa Numune Education and Research Hospital, Department of Neurology, Istanbul, Türkiye). Patients designated as SPMS were in the transition phase from RRMS to SPMS.

^b Seventeen patients in cohort 1 (as indicated in the table) were included for measurement of B-cell regulatory cytokines with ELISA. All patients from cohort 2 were included for measurement of B-cell regulatory cytokines with ELISA. The patients from cohort 2 were not included for immune cell phenotype analyses.

formula: sTACI-BAFF complex detected (pg/mL)/Total sTACI detected (pg/mL) × 100.

Samples from all available time points of a patient were measured on the same plate. In addition, all 5 ELISAs for each patient were conducted on the same day simultaneously. Two day-to-day controls (sera from 2 healthy controls) were included on every ELISA plate to compare the consistency of ELISA measurements.

Statistics

All statistical analyses were performed using GraphPad Prism 8.0 (GraphPad Software, San Diego, CA). Levels of sBCMA, sTACI, BAFF, APRIL, and sTACI-BAFF complexes are presented as log₂ fold changes compared with BL, which is set as 100%, and were analyzed using a 1-sample *t* test and Bonferroni correction (*n* = 6). Pearson correlation analyses were applied to investigate log₂ fold change of BAFF to log₂ fold change of sTACI as well as log₂ fold change of BAFF to log₂ fold change of sTACI-BAFF complexes in the serum of patients. Differences were considered statistically significant for *p* values of **p* < 0.05, ***p* < 0.01, ****p* < 0.001, and *****p* < 0.0001 and marked accordingly in the figures.

Data Availability

After publication, anonymized data will be made available on reasonable request to the corresponding author.

Standard Protocol Approvals, Registrations, and Patient Consents

All patients provided written informed consent, and the study was approved by the ethical committees of the Medical Faculty of Ludwig-Maximilians-Universität Munich (protocol number 163-16) and Haydarpasa Numune Education and Research Hospital, Istanbul.

Results


Effect of Ocrelizumab Treatment on Immune Cell Subsets in Blood

We performed a longitudinal study of the effect of ocrelizumab on CD19⁺ B cells, CD3⁺ T cells, and CD20⁺ T cells in peripheral blood for up to 2.5 years. Our analysis showed that CD19⁺ B cells were persistently depleted. The percentage of CD3⁺ T cells among all lymphocytes increased, but the absolute number of CD3⁺ T cells per microliter blood remained unchanged. The relative increase of CD3⁺ T cells was due to the depletion of B cells. CD20⁺ T cells were also depleted by ocrelizumab treatment (eFigure 1, links.lww.com/NXI/A798; eTable 3, links.lww.com/NXI/A799).

Ocrelizumab Treatment Enhances BAFF and Decreases sTACI in Serum and CSF

Ocrelizumab treatment resulted in increased serum levels of BAFF and decreased sTACI, whereas APRIL and sBCMA did

Figure 2 Effect of Ocrelizumab Treatment on Components of the BAFF/APRIL System in the Serum of Patients



Serum levels of BAFF (A), sTACI (B), and APRIL (C) were evaluated from patient cohort 1 by ELISA from patients receiving ocrelizumab at baseline (*n* = 17), TP1 (mean 18 days; *n* = 17), TP2 (mean 6.7 months; *n* = 17), TP3 (mean 12.9 months; *n* = 17), TP4 (mean 19.1 months; *n* = 14), TP5 (mean 25.2 months; *n* = 10), and TP6 (mean 31.3 months; *n* = 7). Raw values are provided in eTable 3 (links.lww.com/NXI/A799). Data are presented as log₂ fold change compared with baseline. We performed a 1-sample *t* test to compare baseline and follow-up samples. *p* Values are shown if they were <0.05. Data are given as arithmetic mean ± SEM. Negative correlation between BAFF and sTACI levels is shown by the Pearson correlation analyses of log₂ fold change of BAFF to log₂ fold change of sTACI in the serum of patient cohort 1 (D). APRIL = A proliferation-inducing ligand; BAFF = B cell-activating factor; TACI = transmembrane activator and CAML interactor.

not change (Figure 2, A–D; eFigure 2A, links.lww.com/NXI/A798; eTable 3, links.lww.com/NXI/A799). The alterations of BAFF and sTACI appeared already at TP1 and were continuously observed up to TP6 (Figure 2, A and B). Because we obtained highly significant results for BAFF, sTACI, and the sTACI-BAFF complexes with cohort 1, we started the evaluation of a second cohort, from whom both serum and CSF were available.

Also in cohort 2, BAFF increased, sTACI decreased, and sBCMA remained unchanged, whereas we noted a slight reduction of APRIL (eFigure 2, B–E, links.lww.com/NXI/A798; eTable 3, links.lww.com/NXI/A799). sTACI correlated negatively with BAFF in both cohorts (Figure 2D; eFigure 2F, links.lww.com/NXI/A798). Also in CSF, BAFF increased, and sTACI decreased following ocrelizumab, whereas APRIL and sBCMA remained unaltered (Figure 3, A–D; eTable 3, links.lww.com/NXI/A799). Thus, we consistently observe that ocrelizumab decreases sTACI in serum and CSF.

Having observed that ocrelizumab strongly increased BAFF levels, we asked whether the elevated BAFF could interfere with the detection of sTACI or sBCMA. When we spiked BAFF in recombinant sTACI (Figure 4, A–C) and serum (data not shown), we observed reduced detection of sTACI; This was observed with 2 concentrations of sTACI reflecting sTACI levels in serum (Figure 4A). Although this shows that BAFF reduces the detection of sTACI in our ELISA, this does


not exclude that the sTACI ELISA partially also detects with low-efficiency sTACI-BAFF complexes. In contrast, the detection of sTACI was not reduced by spiking in APRIL (eFigure 3, links.lww.com/NXI/A798). Consequently, our sTACI ELISA detected free sTACI or sTACI-APRIL complexes, but rather not sTACI-BAFF complexes. The detection of sBCMA, in contrast to sTACI, was neither reduced by BAFF nor by APRIL (eFigure 3, links.lww.com/NXI/A798). Thus, the BCMA detected in our ELISA represents total sBCMA, free sBCMA, or BCMA bound by APRIL or BAFF.

We also noted that the spiking of sTACI reduced the detection of BAFF, but not of APRIL (eFigure 3, links.lww.com/NXI/A798). Thus, our APRIL ELISA detects total APRIL, both free APRIL and the complexes of APRIL and sTACI. The amount of sTACI is decreasing after ocrelizumab therapy in the CSF (Figure 3) and sTACI is a decoy for APRIL; this suggests that the amount of free APRIL measured in our ELISA is increasing.

Ocrelizumab Induces sTACI-BAFF Complexes


Having found that BAFF reduces the detection of sTACI (Figure 4A), we set up an ELISA to detect sTACI-BAFF complexes by combining well-established antibodies to sTACI and BAFF (Figure 4, B and C). As a result, we found that ocrelizumab induced the formation of sTACI-BAFF complexes in both cohorts (Figure 5, A and B; eTable 3, links.lww.com/NXI/A799). The BAFF levels in serum correlated

Figure 3 Effect of Ocrelizumab Treatment on Components of the BAFF/APRIL System in CSF



CSF levels of BAFF (A), sTACI (B), APRIL (C) and sBCMA (D) were evaluated from patient cohort 2 by ELISA from patients receiving ocrelizumab at baseline (n = 19) and follow-up (12–19 months after the first ocrelizumab infusion). Data are presented as \log_2 fold change compared with baseline. We performed a 1-sample *t* test to compare baseline and follow-up samples. *p* Values are shown if they were <0.05. Data are given as arithmetic mean \pm SEM. Raw values are provided in eTable 3 (links.lww.com/NXI/A799). APRIL = A proliferation-inducing ligand; BAFF = B-cell activating factor; sBCMA = soluble B-cell maturation antigen; CSF = cerebrospinal fluid; sTACI = soluble transmembrane activator and CAML interactor.

Figure 4 Establishment of ELISA Detecting sTACI-BAFF Complexes



(A) Measurement of sTACI (pg/mL) spiked with different amounts of BAFF using an ELISA approach. We added a 2-fold dilution series of BAFF (2,500 to 9.8 pg/mL) to 2 stable concentrations of sTACI (500 pg/ml left part) or 132 pg/mL of (right part) and compared with sTACI alone without BAFF. The left-most column represents no BAFF addition. (B) ELISA design used for the detection of sTACI-BAFF complexes. Anti-BAFF capture antibody and anti-sTACI detection antibody were used to detect sTACI-BAFF complexes. (C) Detection of sTACI-BAFF complexes (OD). Serial dilutions of 1,600, 800, and 400 pg/mL of BAFF were incubated with 500, 250, and 125 pg/mL of sTACI, respectively. BAFF = B cell-activating factor; TACI = transmembrane activator and CAML interactor.

with sTACI-BAFF complexes (Figure 5C). We calculated the percentage of sTACI in complexes with BAFF and observed a steady increase from 43.8% at BL to more than 80% in cohort 1 following ocrelizumab treatment (eTable 4, links.lww.com/NXI/A799). Similarly, also in cohort 2, the percentage of sTACI-BAFF complexes doubled after ocrelizumab therapy (eTable 4, links.lww.com/NXI/A799).

In our cohort 1, we followed the patients longitudinally for up to 2.5 years. Elevation of BAFF, reduction of sTACI, and formation of sTACI-BAFF complexes persist throughout this long observation period, which supports the view that this effect is a consequence of the anti-CD20 treatment and cannot be attributed to treatments before ocrelizumab therapy was started. This study provides Class IV evidence that


endogenous sTACI in blood and CSF is decreased after ocrelizumab treatment.

Discussion

Here, we report that endogenous sTACI in blood and CSF is decreased after ocrelizumab treatment. It is tempting to speculate that this contributes to the beneficial effect of ocrelizumab, as the application of exogenous soluble TACI (atacept) worsened MS.^{17,18}

By which mechanism could sTACI modulate MS? sTACI functions as a decoy for BAFF and APRIL^{13,24} (Figures 1 and 6). Thus, a reduction of sTACI enhances APRIL activity, and

Figure 5 Identification and Quantification of sTACI-BAFF Complexes



(A) sTACI-BAFF complexes were measured by ELISA in patient cohort 1 in serum samples of patients at baseline (n = 17), TP1 (mean 18 days; n = 17), TP2 (mean 6.7 months; n = 17), TP3 (mean 12.9 months; n = 17), TP4 (mean 19.1 months; n = 14), TP5 (mean 25.2 months; n = 10), and TP6 (mean 31.3 months; n = 7). (B) Serum levels of sTACI-BAFF complex were evaluated in 15 patients from cohort 2 at baseline and follow-up (12–19 months after the first ocrelizumab infusion). Data are presented as log₂ fold change compared with baseline. We performed a 1-sample *t* test to compare baseline and follow-up samples. Detailed values are given in eTable 3 (links.lww.com/NXI/A799). Data are given as arithmetic mean ± SEM. (C) The BAFF levels in serum correlated with sTACI-BAFF complexes (Pearson correlation analyses of log₂ fold change of BAFF to log₂ fold change of sTACI-BAFF complexes, *p* = 0.0032, *r* = 0.3252, cohort 1). BAFF = B cell-activating factor; TACI = transmembrane activator and CAML interactor; TP1 = time point 1.


intriguingly, recent work has indicated a beneficial effect of APRIL on CNS inflammation: APRIL, which is produced in MS lesions²³ and by meningeal fibroblasts,²⁵ stimulates astrocytes to secrete the inhibitory cytokine IL-10 (Ref. 23).

Although BAFF and APRIL share the receptor TACI, there are further aspects to be considered concerning their specific role in neuroinflammation. First, there is *in vitro* evidence that TACI, unlike BAFF-R, is solely activated by oligomeric BAFF and APRIL²⁶; BAFF in the serum and CSF of patients with MS is trimeric, not oligomeric.²⁷ APRIL can be oligomerized via heparan sulfate proteoglycan binding to activate TACI signaling.¹¹ Second, APRIL exerts functions beyond its binding to TACI, impressively shown by the ability of APRIL to limit atherosclerosis by binding to heparan sulfate proteoglycans.²⁸ APRIL induces IL-10 production by astrocytes²³ and astrocytes do not express TACI; these APRIL-specific effects are not understood in all details. Consistent with this, APRIL-deficient mice develop a more severe experimental autoimmune encephalomyelitis (EAE).²³

Furthermore, APRIL promotes IgA class switching²⁹ and induces regulatory B cells, some of which are IgA positive.^{30,31} Importantly, regulatory IgA⁺ plasma cells dampen CNS inflammation.^{32,33} It was already observed that the production of IgA⁺ plasmablasts continues despite anti-CD20 therapy.¹⁰ We propose that the reduced sTACI and enhanced APRIL activity are one explanation for this persistent production of IgA plasmablasts.

We addressed the mechanisms by which ocrelizumab treatment reduces sTACI. We found that ocrelizumab reduces sTACI but leaves sBCMA unaltered. This might be surprising because both sTACI and sBCMA are released from activated B cells and differentiated plasma cells.^{12,13} B-cell depletion reduces the amount of newly formed plasmablasts, but the long-lived CD20⁻ plasma cells in the bone marrow remain.¹ The unaltered sBCMA levels in blood during anti-CD20 treatment indicate that it is largely derived from long-lived plasma cells, consistent with the observation that IgG in blood derives mainly from long-lived plasma cells and shows little changes during anti-CD20 treatment at least in the first years of treatment.¹ We tested whether the sTACI reduction after ocrelizumab is due to interaction of sTACI with soluble ligands. B-cell depletion with anti-CD20 elevates BAFF, but not APRIL.¹⁵ We assume that this is due to reduced BAFF consumption because the depleted B cells express BAFF-R, which binds BAFF, but not APRIL.¹¹ We established an ELISA that specifically quantifies sTACI-BAFF complexes and found elevated sTACI-BAFF complexes during ocrelizumab treatment that correlated positively with BAFF and negatively with the sTACI levels. It might be surprising that elevated BAFF levels do not reduce sBCMA, although BCMA binds both BAFF and APRIL.¹¹ We observed that spiking in BAFF into our sBCMA ELISA had no effect on the measured sBCMA levels. This directly explains why the elevated BAFF does not reduce the detected sBCMA levels. Furthermore, we should consider the lower affinity of BCMA to BAFF

Figure 6 Illustration Showing Our Findings With the Proposed Consequences



The upper part illustrates our findings: ocrelizumab treatment results in an increase of BAFF and reduced free sTACI in both serum and CSF. In serum, this is associated with increased formation of sTACI-BAFF complexes. sTACI-BAFF complexes were not detected in CSF samples in our study. The lower part links our findings to published work: sTACI is a decoy for APRIL¹³ (left). In the absence of the decoy sTACI (right), APRIL might induce IL10 production from astrocytes²³ and promote the development of regulatory IgA plasma cells,²⁹⁻³¹ which can home to the brain and secrete IL-10.^{32,33} APRIL = A proliferation-inducing ligand; BAFF = B cell-activating factor; TACI = transmembrane activator and CAML interactor.

compared with APRIL; BAFF binding to BCMA requires a gain of avidity due to oligomerization of BCMA,³⁴ and therefore, membrane-bound BCMA also binds BAFF. Importantly, sBCMA is a monomer as we have observed before, which binds APRIL, but not BAFF,¹² whereas sTACI binds both BAFF and APRIL.¹³ All this explains why the elevation of BAFF has no effect on sBCMA but results in reduction of free sTACI.

Recently, it was observed that anti-CD20 elevated BAFF and this was inversely correlated with CNS inflammation, and it was proposed that anti-CD20 exerts a protective effect by providing a favorable niche for IL-10–producing B cells and suppressive IgA⁺ plasma cells.^{35,36} We now show that anti-CD20 reduces the decoy sTACI, which may result in enhanced APRIL activity; this could contribute to the proposed mechanism fostering suppressive plasma cells.¹⁸

We provide information about the concentrations of B-cell regulatory factors and soluble receptors in blood and CSF, but it is a limitation of our study that we cannot measure their concentrations directly in inflammatory lesions. The strong effects of the systemically given atacept, however, argue that the systemic levels of soluble TACI modulate the local microenvironment in the CNS. Although we selected only patients without prior depleting therapy, we cannot exclude the possibility that the therapy had an effect on our results. We show that anti-CD20 therapy reduces sTACI, a decoy for APRIL, but the linkage to local APRIL activity remains to be shown. To study the role of sTACI in neuroinflammation directly, a knock-in mouse expressing a mutated nonshed-dable TACI could be made; such an approach was applied to study the shedding of TNFR1 in EAE.³⁷

Altogether, we identify a novel effect of ocrelizumab treatment on the BAFF-APRIL system, a crucial regulatory element for B cells and plasma cells. We describe that the decoy sTACI is reduced over an observation time of up to 2.5 years of ocrelizumab therapy due to formation of sTACI-BAFF complexes. We propose that the reduced decoy sTACI leads to an enhanced local APRIL activity that could induce anti-inflammatory activity in the CNS and support the development and maintenance of regulatory plasma cells (Figure 6).

Acknowledgment

The authors thank Prof. Dr. Reinhard Hohlfeld and Prof. Dr. Pascal Schneider for continuous support and fruitful discussions, Damla Taskin and Yaren Canten for support with sample archiving, and Dipl.-Ing Benjamin Obholzer for graphical illustration. The authors thank the Institute of Laboratory Medicine, Ludwig-Maximilians-Universität München, for FACS measurements. The authors are grateful to PD Dr. Lisa Ann Gerdes and Dr. Anneli Peters for comments on the manuscript.

Study Funding

This work was supported by the DFG (SFB TR128) to E.M., the Verein zur Therapieforschung für MS Kranke to E.M., by

Roche to E.M. and the Else Kröner-Fresenius-Stiftung (EKFS) to S.M.

Disclosure

S. Ho, E. Oswald, H.K. Wong, A. Vural, V. Yilmaz, E. Tüzün, R. Türkoğlu, and T. Straub report no disclosures relevant to the manuscript. I. Meinel received payment from Roche. F. Thaler received grant support from Novartis Pharma GmbH. T. Kümpfel has served on advisory boards for Roche Pharma and has received personal compensations/speaker honoraria from Bayer Healthcare, Teva Pharma, Merck, Novartis Pharma, Sanofi-Aventis/Genzyme, Roche Pharma, and Biogen and grant support from Novartis and Chugai Pharma in the past. E. Meinel received funding for travel or speaker honoraria by (1) Roche, honorarium; (2) Novartis, honorarium; (3) Sanofi, honorarium; (4) Biogen, honorarium; (5) Bioeq, honorarium; and (6) Merck, honorarium. E. Meinel received research support from the following commercial entities: (1) Novartis, (2) Sanofi, (3) Merck, and (4) Roche. S. Mader received grant support from Novartis Pharma GmbH. Go to Neurology.org/NN for full disclosures.

Publication History

Received by *Neurology: Neuroimmunology & Neuroinflammation* June 10, 2022. Accepted in final form November 22, 2022. Submitted and externally peer reviewed. The handling editor was Deputy Editor Scott S. Zamvil, MD, PhD, FAAN.

Appendix Authors

Name	Location	Contribution
Samantha Ho, MSc	Institute of Clinical Neuroimmunology, Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München; Graduate School of Systemic Neurosciences, Ludwig Maximilians University Munich, Germany	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; and analysis or interpretation of data
Eva Oswald, BSc	Institute of Clinical Neuroimmunology, Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München, Germany	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; and analysis or interpretation of data
Hoi Kiu Wong, MSc	Institute of Clinical Neuroimmunology, Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München, Germany	Drafting/revision of the manuscript for content, including medical writing for content, and analysis or interpretation of data
Atay Vural, MD, PhD	Institute of Clinical Neuroimmunology, Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München, Germany; Department of Neurology, Koc University School of Medicine, Istanbul, Türkiye	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; and analysis or interpretation of data

Appendix (continued)

Name	Location	Contribution
Vuslat Yılmaz, PhD	Department of Neuroscience, Aziz Sancar Institute of Experimental Medicine, Istanbul University, Türkiye	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; and analysis or interpretation of data
Erdem Tüzün, MD	Department of Neuroscience, Aziz Sancar Institute of Experimental Medicine, Istanbul University, Türkiye	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; and analysis or interpretation of data
Recai Türkoğlu, MD	Department of Neurology, Haydarpaşa Numune Education and Research Hospital, Istanbul, Türkiye	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; and analysis or interpretation of data
Tobias Straub, MD	Core Facility Bioinformatics, Biomedical Center, Ludwig-Maximilians-Universität München, Germany	Drafting/revision of the manuscript for content, including medical writing for content, and analysis or interpretation of data
Ingrid Meinel, MD	Institute of Clinical Neuroimmunology, Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München, Germany	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; and analysis or interpretation of data
Franziska Thaler, MD	Institute of Clinical Neuroimmunology, Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München, Germany; Munich Cluster for Systems Neurology (SyNergy), Germany	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; and analysis or interpretation of data
Tania Kümpfel, MD	Institute of Clinical Neuroimmunology, Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München, Germany	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; and analysis or interpretation of data
Edgar Meinel, MD	Institute of Clinical Neuroimmunology, Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München, Germany	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; and analysis or interpretation of data
Simone Mader, PhD	Institute of Clinical Neuroimmunology, Biomedical Center and University Hospitals, Ludwig-Maximilians-Universität München, Germany	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; and analysis or interpretation of data

References

- Comi G, Bar-Or A, Lassmann H, et al. Role of B cells in multiple sclerosis and related disorders. *Ann Neurol*. 2021;89(1):13-23. doi:10.1002/ana.25927.
- Graf J, Mares J, Barnett M, et al. Targeting B cells to modify MS, NMOSD, and MOGAD: part 1. *Neurol Neuroimmunol Neuroinflamm*. 2021;8:e918. doi:10.1212/xxi.0000000000000918.
- Palanichamy A, Jahn S, Nickles D, et al. Rituximab efficiently depletes increased CD20-expressing T cells in multiple sclerosis patients. *J Immunol*. 2014;193(2):580-586. doi:10.4049/jimmunol.1400118.
- Schuh E, Berer K, Mulazzani M, et al. Features of human CD3+CD20+ T cells. *J Immunol*. 2016;197(4):1111-1117. doi:10.4049/jimmunol.1600089.
- Quendt C, Ochs J, Häusser-Kinzel S, Häusler D, Weber MS. Proinflammatory CD20(+) T cells are differentially affected by multiple sclerosis therapeutics. *Ann Neurol*. 2021;90(5):834-839. doi:10.1002/ana.26216.
- Sabatino JJ Jr, Wilson MR, Calabresi PA, Hauser SL, Schneke JP, Zamvil SS. Anti-CD20 therapy depletes activated myelin-specific CD8(+) T cells in multiple sclerosis. *Proc Natl Acad Sci*. 2019;116(51):25800-25807. doi:10.1073/pnas.1915309116.
- Meinel E, Hohlfeld R. CD20(+) T cells as pathogenic players and therapeutic targets in MS. *Ann Neurol*. 2021;90(5):722-724. doi:10.1002/ana.26232.
- Hausler D, Häusser-Kinzel S, Feldmann L, et al. Functional characterization of reappearing B cells after anti-CD20 treatment of CNS autoimmune disease. *Proc Natl Acad Sci USA*. 2018;115(39):9773-9778. doi:10.1073/pnas.1810470115.
- Theil D, Smith P, Huck C, et al. Imaging mass cytometry and single-cell genomics reveal differential depletion and repletion of B-cell populations following ofatumumab treatment in cynomolgus monkeys. *Front Immunol*. 2019;10:1340. doi:10.3389/fimmu.2019.01340.
- Mei HE, Frölich D, Giesecke C, et al. Steady-state generation of mucosal IgA+ plasmablasts is not abrogated by B-cell depletion therapy with rituximab. *Blood*. 2010;116(24):5181-5190. doi:10.1182/blood-2010-01-266536.
- Mackay F, Schneider P. Cracking the BAFF code. *Nat Rev Immunol*. 2009;9(7):491-502. doi:10.1038/nri2572.
- Laurent SA, Hoffmann FS, Kuhn PH, et al. γ -secretase directly sheds the survival receptor BCMA from plasma cells. *Nat Commun*. 2015;6(1):7333. doi:10.1038/ncomms8333.
- Hoffmann FS, Kuhn PH, Laurent SA, et al. The immunoregulator soluble TACI is released by ADAM10 and reflects B cell activation in autoimmunity. *J Immunol*. 2015;194(2):542-552. doi:10.4049/jimmunol.1402070.
- Meinel E, Thaler FS, Lichtenthaler SF. Shedding of BAFF/APRIL receptors controls B cells. *Trends Immunol*. 2018;39(9):673-676. doi:10.1016/j.it.2018.07.002.
- Pellkofer HL, Krumbholz M, Berthele A, et al. Long-term follow-up of patients with neuromyelitis optica after repeated therapy with rituximab. *Neurology*. 2011;76(15):1310-1315. doi:10.1212/WNL.0b013e3182152881.
- Kreuzaler M, Rauch M, Salzer U, et al. Soluble BAFF levels inversely correlate with peripheral B cell numbers and the expression of BAFF receptors. *J Immunol*. 2012;188(1):497-503. doi:10.4049/jimmunol.1102321.
- Kappos L, Hartung HP, Freedman MS, et al. Atacept in multiple sclerosis (ATAMS): a randomised, placebo-controlled, double-blind, phase 2 trial. *Lancet Neurol*. 2014;13(4):353-363. doi:10.1016/S1474-4422(14)70028-6.
- Sergott RC, Bennett JL, Rieckmann P, et al. ATON: results from a phase II randomised trial of the B-cell-targeting agent atacept in patients with optic neuritis. *J Neurol Sci*. 2015;351(1-2):174-178. doi:10.1016/j.jns.2015.02.019.
- Krumbholz M, Derfuss T, Hohlfeld R, Meinel E. B cells and antibodies in multiple sclerosis pathogenesis and therapy. *Nat Rev Neurol*. 2012;8(11):613-623. doi:10.1038/nrneuro.2012.203.
- Sabatino JJ Jr, Probstel AK, Zamvil SS. B cells in autoimmune and neurodegenerative central nervous system diseases. *Nat Rev Neurosci*. 2019;20(12):728-745. doi:10.1038/s41583-019-0233-2.
- Kumar G, Maria Z, Kohli U, et al. CNS autoimmune responses in BCMA-deficient mice provide insight for the failure of atacept in MS. *Neurol Neuroimmunol Neuroinflamm*. 2021;8(3):e973. doi:10.1212/xxi.0000000000000973.
- Kim SS, Richman DP, Zamvil SS, Agius MA. Accelerated central nervous system autoimmunity in BAFF-receptor-deficient mice. *J Neurol Sci*. 2011;306(1-2):9-15. doi:10.1016/j.jns.2011.04.008.
- Baert L, Benkhoucha M, Popa N, et al. A proliferation-inducing ligand-mediated anti-inflammatory response of astrocytes in multiple sclerosis. *Ann Neurol*. 2019;85(3):406-420. doi:10.1002/ana.25415.
- Hymowitz SG, Patel DR, Wallweber HJ, et al. Structures of APRIL-receptor complexes: like BCMA, TACI employs only a single cysteine-rich domain for high affinity ligand binding. *J Biol Chem*. 2005;280(8):7218-7227. doi:10.1074/jbc.m411714200.
- Brioschi S, Wang WL, Peng V, et al. Heterogeneity of meningeal B cells reveals a lymphopoietic niche at the CNS borders. *Science*. 2021;373(6553):eab9277. doi:10.1126/science.ab9277.
- Bossen C, Cachero TG, Tardivel A, et al. TACI, unlike BAFF-R, is solely activated by oligomeric BAFF and APRIL to support survival of activated B cells and plasmablasts. *Blood*. 2008;111(3):1004-1012. doi:10.1182/blood-2007-09-110874.
- Esiami M, Meinel E, Eibel H, et al. BAFF 60-mer, and differential BAFF 60-mer dissociating activities in human serum, cord blood and cerebrospinal fluid. *Front Cell Dev Biol*. 2020;8:577662. doi:10.3389/fcell.2020.577662.
- Tsiantoulas D, Esiami M, Obermayer G, et al. APRIL limits atherosclerosis by binding to heparan sulfate proteoglycans. *Nature*. 2021;597(7874):92-96. doi:10.1038/s41586-021-03818-3.
- Castigli E, Scott S, Dedeoglu F, et al. Impaired IgA class switching in APRIL-deficient mice. *Proc Natl Acad Sci USA*. 2004;101(11):3903-3908. doi:10.1073/pnas.0307348101.
- Hua C, Audo R, Yeremenko N, et al. A proliferation inducing ligand (APRIL) promotes IL-10 production and regulatory functions of human B cells. *J Autoimmun*. 2016;73:64-72. doi:10.1016/j.jaut.2016.06.002.
- Fehres CM, van Uden NO, Yeremenko NG, et al. APRIL induces a novel subset of IgA(+) regulatory B cells that suppress inflammation via expression of IL-10 and PD-L1. *Front Immunol*. 2019;10:1368. doi:10.3389/fimmu.2019.01368.

32. Pröbstel AK, Zhou X, Baumann R, et al. Gut microbiota-specific IgA(+) B cells traffic to the CNS in active multiple sclerosis. *Sci Immunol*. 2020;5(53):eabc7191. doi:10.1126/sciimmunol.abc7191.
33. Rojas OL, Pröbstel AK, Porfilio EA, et al. Recirculating intestinal IgA-producing cells regulate neuroinflammation via IL-10. *Cell*. 2019;177(2):492-493. doi:10.1016/j.cell.2019.03.037.
34. Day ES, Cachero TG, Qian F, et al. Selectivity of BAFF/BLyS and APRIL for binding to the TNF family receptors BAFR/BR3 and BCMA. *Biochemistry*. 2005;44(6):1919-1931. doi:10.1021/bi048227k.
35. Wang A, Rojas O, Lee D, Gomerman JL. Regulation of neuroinflammation by B cells and plasma cells. *Immunol Rev*. 2021;299(1):45-60. doi:10.1111/imr.12929.
36. Wang A, Ward L, Ramaglia V, Urich E, Gomerman JL. Impact of B cell-depletion on plasma cells in a pre-clinical model of meningeal inflammation and subpial demyelination. *J Immunol*. 2021;206(1 Suppl):21-06.
37. Xanthoulea S, Pasparakis M, Kousteni S, et al. Tumor necrosis factor (TNF) receptor shedding controls thresholds of innate immune activation that balance opposing TNF functions in infectious and inflammatory diseases. *J Exp Med*. 2004;200(3):367-376. doi:10.1084/jem.20040435.