DOI: 10.1111/irv.12915

ORIGINAL ARTICLE

WILEY

Early epidemiological investigations: World Health Organization UNITY protocols provide a standardized and timely international investigation framework during the COVID-19 pandemic

Isabel Bergeri ¹ 💿 Hannah C. Lewis ¹ 💿 Lorenzo Subissi ¹ 💿
Anthony Nardone ^{1,2} 💿 Marta Valenciano ^{1,2} 💿 Brianna Cheng ¹ 💿
Ketevan Glonti ¹ 💿 Bridget Williams ¹ 💿 Ibukun-Oluwa Omolade Abejirinde ¹ 💿
Alice Simniceanu ¹ 💿 Alessandro Cassini ¹ Rebecca Grant ¹ 💿
Angel Rodriguez ³ Andrea Vicari ³ Lubna Al Ariqi ⁴ Tasnim Azim ⁵
Pushpa Ranjan Wijesinghe ⁵ 💿 📔 Soatiana Cathycia Rajatonirina ⁶ 💿 🛛
Joseph Chukwudi Okeibunor ⁶ 💿 Linh-Vi Le ⁷ 💿 Mark Katz ⁸ 💿
Aisling Vaughan ⁸ 💿 Pernille Jorgensen ⁸ 💿 Gudrun Freidl ⁸ 💿
Richard Pebody ⁸ 💿 📔 Maria D. Van Kerkhove ¹ 💿
Alice Simniceanu ¹ Alessandro Cassini ¹ Rebecca Grant ¹ Tasnim Azim ⁵ Angel Rodriguez ³ Andrea Vicari ³ Lubna Al Ariqi ⁴ Tasnim Azim ⁵ Pushpa Ranjan Wijesinghe ⁵ Soatiana Cathycia Rajatonirina ⁶ Joseph Chukwudi Okeibunor ⁶ Linh-Vi Le ⁷ Mark Katz ⁸ Aisling Vaughan ⁸ Pernille Jorgensen ⁸ Gudrun Freidl ⁸ Richard Pebody ⁸ Maria D. Van Kerkhove ¹

¹WHO Health Emergencies Programme, World Health Organization Headquarters, Geneva, Switzerland

²Epidemiology Departement, Epiconcept, Paris, France

³Pan American Health Organization, Washington, D.C., USA

⁴World Health Organization for the Eastern Mediterranean, Cairo, Egypt

⁵Regional Office for South-East Asia, World Health Organization, New Delhi, India

⁶Regional Office for Africa, World Health Organization, Brazzaville, Republic of the Congo

⁷Regional Office for the Western Pacific, World Health Organization, Manila, Philippines

⁸Regional Office for Europe, World Health Organization, Copenhagen, Denmark

Correspondence

Isabel Bergeri, World Health Organization Headquarters, Geneva, Switzerland. Email: bergerii@who.int **Background:** The declaration of Coronavirus disease 2019 (COVID-19) as a Public Health Emergency of International Concern (PHEIC) on 30 January 2020 required rapid implementation of early investigations to inform appropriate national and global public health actions.

Methods: The suite of existing pandemic preparedness generic epidemiological early investigation protocols was rapidly adapted for COVID-19, branded the 'UNITY studies' and promoted globally for the implementation of standardized and quality studies. Ten protocols were developed investigating household (HH) transmission, the first few cases (FFX), population seroprevalence (SEROPREV), health facilities transmission (n = 2), vaccine effectiveness (n = 2), pregnancy outcomes and transmission, school transmission, and surface contamination. Implementation was supported by WHO and its partners globally, with emphasis to support building surveillance and research capacities in low- and middle-income countries (LMIC).

Results: WHO generic protocols were rapidly developed and published on the WHO website, 5/10 protocols within the first 3 months of the response. As of 30 June 2021, 172 investigations were implemented by 97 countries, of which 62 (64%) were LMIC. The majority of countries implemented population

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. Influenza and Other Respiratory Viruses published by John Wiley & Sons Ltd.

Funding information

8

German Federal Ministry of Health (BMG) COVID-19 Research and Development for WHO; COVID-19 Solidarity Response Fund for WHO

WILEY-

seroprevalence (71 countries) and first few cases/household transmission (37 countries) studies.

Conclusion: The widespread adoption of UNITY protocols across all WHO regions indicates that they addressed subnational and national needs to support local public health decision-making to prevent and control the pandemic.

KEYWORDS

COVID-19, infectious diseases, SARS-CoV-2, seroepidemiology, serology, seroprevalence

1 | BACKGROUND

In December 2019, a novel coronavirus, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), responsible for coronavirus disease 2019 (COVID-19), was first identified in Wuhan, China, from where it rapidly spread worldwide COVID-19 was declared as a public health emergency of international concern (PHEIC) by the World Health Organization (WHO) on 30 January 2020.

One of the key difficulties for decision makers in the early phase of any epidemic of a novel pathogen is to determine the appropriate public health measures in the absence of information on transmission characteristics, extent of infection, severity, and scale of the threat. Following a review of the global response to the last pandemic (2009 influenza pandemic H1N1),¹ the global Consortium for the Standardization of Influenza Seroepidemiology (CONSISE)² and WHO's Influenza Pandemic Special Investigations and Studies (IPSS) initiative were established to develop a suite of standardized early investigation protocols,³ supported by the Global Influenza Programme⁴ and the Pandemic Influenza Preparedness (PIP) Framework.⁵ Standardized protocols were also implemented following the emergence of Middle East Respiratory Syndrome coronavirus (MERS-CoV) in 2012⁶ and Zika virus in 2016⁷ with delineation of a comprehensive global research agenda.⁸

Building on these global pandemic preparedness efforts, we describe the rapid adaptation and implementation of WHO early investigation protocols for COVID-19, branded the 'UNITY studies',⁹ to generate local data to better inform appropriate national and global public health actions and guidelines. The on-going and future value of such studies for pandemic preparedness and response is discussed.

2 | METHODS

2.1 | Overall concept

UNITY studies enable all countries to rapidly and systematically collect standardized robust data on key epidemiological, virological, and clinical parameters to understand key characteristics of SARS-CoV-2.

2.2 | Embedment into global WHO COVID strategic plan, financing, and accountability

The UNITY initiative is embedded into both 2020 and 2021 WHO's COVID-19 Strategic Preparedness Response Plans (SPRP) and COVID-19 Monitoring and Evaluation framework.¹⁰ Financial support from May 2021 (see acknowledgements) was instrumental for LMIC.

2.3 | Generic protocol development

Ten generic protocols using eight study methodologies were developed (three adapted from available IPSS protocols). These investigate household (HH) transmission, the first few cases (FFX) and their contacts, population seroprevalence (SEROPREV), health facility transmission (two protocols), vaccine effectiveness (two protocols), pregnancy outcomes and transmission, school transmission, and surface contamination. More information on each protocol is described in Table 1.

Four protocols were designed from inception to be multi-country and multi-centre research studies with sharing of anonymised individual level data, a subsequent planned pooled analysis and related tools including joint data platforms (see Table 1). Remaining protocols were mostly considered to be enhanced surveillance protocols by implementing countries.

2.4 | Implementation strategy

WHO Headquarter (HQ), Regional Offices (RO) and Country Offices (CO) and its Special research programmes, supported the dissemination and implementation of protocols to countries, with translation in the six official languages of WHO. This was in interaction with Ministries of Health, local institutions, and through existing collaborations with relevant disease networks and national and international partners (e.g., academic institutions, US-CDC, and Institut Pasteur). Dissemination was also through United Nations and WHO information products and publications including the WHO website and a generic WHO email address. An insignia (Figure 1) was designed to foster belonging to a join global effort.

Number of countries Number of implementing study countries Date of first publication of LMIC (% HIC (% intending to Protocols name Description protocol on WHO website implement study Total of total) of total) 31/01/2020 24 3 First Few X (FFX) cases and A prospective case-17 (71%) 7 (29%) contacts transmission ascertained investigation of all identified close contacts of laboratoryconfirmed infections to characterize the transmission dynamics and clinical spectrum of COVID-19 infection in the general population Household (HH) A prospective investigation of 25/01/2020 13 5 (38%) 8 (62%) 1 household contacts of transmission laboratory-confirmed COVID-19 cases to reinforce the estimates of transmission dynamics and clinical spectrum of infection obtained from the FFX studies for the general population. Health facility transmission Two protocols available to 36 25 (69%) 11 (31%) 17 assess the transmissibility of and risk factors for COVID-19 infection among health workers: Case-control study 01/02/2020^a Prospective cohort study 26/05/2020 ٠ Surface contamination A practical "how to" guide on 20/02/2020 and transmission surface sampling for COVID-19 virus to estimate the persistence of contamination on different surfaces and their possible role in onward transmission. 71 Population-based age-Both cross-sectional and 19/03/2020 50 (70%) 21 (30%) 27 stratified seroprevalence longitudinal cohort designs (SEROPREV) to conduct serological surveys to estimate cumulative incidence in the population by age and sex. School transmission A prospective case-30/11/2020 ascertained investigation of school contacts of a laboratory-confirmed case of COVID-19 to understand the dynamics of infection among students and staff of schools and childcare institutions

TABLE 1 UNITY protocols description by publication date on the World Health Organization (WHO)'s website and number of countries intending to or having implemented UNITY protocols⁹ by investigation type, as of 30 June 2021

WILEY

(Continues)

TABLE 1 (Continued)

LEY-

	Description	Date of first publication of protocol on WHO website	Number of countries implementing study			Number of countries
Protocols name			Total	LMIC (% of total)	HIC (% of total)	intending to implement study
Maternal, pregnancy and neonatal outcomes and transmission	A prospective cohort study to determine if SARS-CoV-2 infection during pregnancy increases the risk of adverse pregnancy, post- partum or neonatal outcomes ^a	04/12/2020 ^a	10	8 (80%)	2 (20%)	9
Vaccine effectiveness	Two protocols available to measure product-specific COVID-19 vaccine effectiveness (VE):		17	2 (13%)	15 (87%)	1
	 Test negative case-control in hospitalized severe acute respiratory infections (SARI) patients 	18/03/2021				
	Prospective cohort study among health workers	04/05/2021				

Abbreviations: HIC, high-income country; LMIC, low- and middle-income country. ^aMulti-centre studies.

2.5 | Comparability of results through standardization and direct country support

WHO offered substantial technical, financial, and material support to national investigation teams to implement quality UNITY studies, particularly in LMIC. Technical support included scientific review of locally adapted protocols, provision of tools including the option of using Go.Data software¹¹), and guidance for field implementation and quality monitoring, advice on laboratory methods, and assistance with data analysis. Training in scientific writing was provided to national teams to facilitate public dissemination of results.

To enhance the comparability of different studies, WHO also advocated for the standardization of serology laboratory methods. Serology testing was standardized through various means: from March 2021 central WHO procurement of manual total antibody ELISAs targeting the receptor-binding domain (RCB) of SARS-CoV-2 spike (S) protein and that demonstrated sensitivity \geq 95% and specificity \geq 99% in multiple independent evaluations including also samples from mild and asymptomatic infections¹²; provision of research reagent and panel from April 2020¹³; comparative performance evaluation of commercially available assays from Dec 2020¹⁴; and availability of International Standard by December 2020.¹³

As collaborators undertaking UNITY investigations, national teams were asked to share aggregate results in standardized templates while being assured that ownership of primary data rested firmly with the implementing countries and institutions.



FIGURE 1 Insignia of WHO UNITY early investigations initiative

2.6 | Monitoring of country adoption of protocols

WHO ROs and COs provided weekly updates on the number of countries adopting UNITY studies, defined as either 'intending to undertake' or 'having implemented' a protocol. 'Implementation' was defined as a nationally adapted protocol scientifically validated by WHO, national ethical approval obtained (or waived), and initiation of the investigation (i.e., at least one participant enrolled or one sample taken).

3 | RESULTS

Five of the ten protocols were published on the WHO website within the first 3 months of response (before 1 April 2020). As of 30 June

10

2021, 129 countries registered intention to undertake investigations aligned with WHO UNITY studies, and 172 studies had been implemented by 97 countries of which 64% (62/97) were low- and middle-income countries (LMIC) (Table 1 and Figure 2). Studies have been implemented in all WHO regions and in half (53%, 34/64) of all Humanitarian Response Plan (HRP) countries.¹⁵

The most commonly implemented investigation was the SEROPREV investigation, which has been implemented by 71 countries (Table 1), followed by FFX and HH transmission investigations (n = 37), this last one particularly in the early stage of the pandemic. Health facility transmission investigations were implemented in 36 countries. The monitoring of countries which aligned with or adopted the school or surface contamination protocol was not possible with our human resource capacity at that time. Nevertheless, we retrospectively assess that their uptake was likely low for various reasons including the pandemic dynamic. The pregnancy outcomes and transmission protocol and the two vaccine effectiveness protocols were developed and published at later stage, in December 2020 and March 2021, respectively, to respond to emergent scientific priorities, including to measure the performance of newly implemented vaccine programmes. As of 30 June 2021, 39 countries across all six WHO Geographic Regions (most being LMIC) have been supplied by a centrally procured manual ELISA tests to promote standardization of results.

National teams are sharing standardized results in the templates provided. UNITY studies has enabled countries to more accurately estimate epidemiological indicators, such as transmission and severity parameters, risk factors, seroprevalence, cumulative incidence and secondary attack and infection rates, thereby enhancing routine surveillance.

4 | DISCUSSION

The widespread adoption of UNITY protocols across all WHO regions indicates that they have addressed subnational and national needs to collect critical data to support local and regional public health decision-making to prevent and control the pandemic. The quarterly monitoring of the implementation of UNITY investigations as a performance indicator of the COVID-19 SPRP¹⁰ further reinforces their significance and impact to also inform global normative work.

WHO rapidly adapted and established the suite of protocols as part of its pandemic preparedness and response planning, which were rapidly adopted by countries. FFX and HH investigations implementation was more relevant for country action early in the pandemic to robustly define key epidemiological and severity parameters of the infection. Nonetheless, the continued collection of such data can provide important insights into the transmission of variants viruses of interest or of concern. It is likely that SEROPREV investigations were of higher priority for countries for rapid policy decision making (and particularly in countries with low testing capacities or weak case-based reporting system) than for other less used protocols (e.g., school transmission and surface contamination). The assessment of uptake of the more recently developed pregnancy and Vaccine Effectiveness protocols will require more time.



FIGURE 2 Countries that have started implementing at least one sero-epidemiological investigation aligned with WHO UNITY generic protocols, from Jan 2020 to 30 June 2021

12 ⊥WILEY_

The large number of LMIC, including those affected by armed conflict and political instability (HRP countries), that adopted a UNITY protocol is of note and indicates the value of having readily available, generic protocols that can be adapted for various settings. The number of adopted UNITY studies in this analysis is likely to be under-reported as institutions may have employed the publicly available protocols independent of WHO support or knowledge, particularly in HIC.

The first set of significant funding support arrived 4 months after the first protocols publication which hampered timely implementation in LMIC. Tight deadlines for fund expenditure from effective reception (ex: less than 3.5 months) was an issue for such activities.

The provision of technical, financial, and laboratory assay support by WHO and international partners and using existing surveillance and research networks was instrumental in the implementation of timely, quality, and standardized studies to better understand COVID-19 transmission and extent of infection. The support provided enabled countries, regardless of economic status, to build on routine surveillance and develop research capacities, to further scientific learning and support evidence-based response decisions at local, national, regional and international level. Laboratory standardization has been key to enable inter-country comparisons. Main lessons learned are described in Box 1.

Box 1. Lessons learned from the WHO UNITY framework during the COVID-19 pandemic Lessons learned:

- Need for a standardized global framework for epidemiological and laboratory investigations, with ready-to-be used, easily adaptable tools, which are adaptable to any resource setting and which can allow inter-country comparisons
- Operationnal suite of tools supported country enhanced surveillance, the strengthening research capacities independently to resource setting, and in addressing national, regional and global knowledge gaps
- Such framework could be valuable and replicated for any emerging and re-emerging pathogen pandemic readiness and preparedness plans
- Timely and pooled fund availability to ensure equitable support is instrumental in LMIC
- Building on existing networks of partners, research groups, surveillance systems and public health services were essential for the feasibility of such initiatives

WHO is working with UNITY partners to summarise available study results and to conduct pooled analyses where relevant, by studying populations and at regional and global levels.

An after action review and evaluation of the UNITY initiatives and its protocols to assess their utility to countries in responding to this pandemic and whether they fulfilled key initial objectives started in September 2021 and will be valuable to inform future pandemic preparedness, readiness and response strategy to any respiratory pathogens.

ACKNOWLEDGEMENTS

The authors would like to thank people in all countries who are working in the global response effort to COVID-19, to thank all WHO regional and country offices and partners organizations such as Institut Pasteur and the Institut Pasteur International Network, US-CDC, the UNDP/UNFPA/UNICEF/WHO/World Bank Special Programme of Research, Development and Research Training in Human Reproduction (HRP), as well as all individuals who supported, conducted or participated in each of the studies supported.

Further acknowledgement to the following WHO staff: Ana Paula Coutinho Rehse, Giorgia Gon, Vicki Indenbaum, Amal Barakat, Eman Alv. Arash Rashidian, Bassim Zahed, Francis Inbanathan, Pushpa Wijesinghe, Belinda Herring, Pui Shan Chung, Juliana Leite, Joao Toledo, Daniel Feikin, Minal Patel, Sudhir Joshi, Jason Mathiu, Ambrose Talisuna, Ngyo Nsenga, Tom Grein, Anna Thorson, and Nathalie Broutet.

This work was supported by the COVID-19 Solidarity Response Fund for WHO and German Federal Ministry of Health (BMG) COVID-19 Research and Development grant for WHO.

AUTHOR CONTRIBUTIONS

Isabel Bergeri: Conceptualization; formal analysis; funding acquisition; investigation; methodology; project administration; resources; supervision; visualization. Hannah C. Lewis: Conceptualization; formal analysis; investigation; methodology; project administration; supervision. Lorenzo Subissi: Conceptualization; formal analysis; investigation; methodology; project administration; resources; supervision. Anthony Nardone: Conceptualization; formal analysis; investigation; methodology; supervision. Marta Valenciano: Conceptualization; formal analysis; investigation; methodology; supervision. Brianna Cheng: Investigation; project administration; resources; visualization. Ketevan Glonti: Investigation; project administration; resources; visualization. Bridget Williams: Conceptualization; investigation; project administration; resources; visualization. Ibukun-Oluwa Omolade Abejirinde: Formal analysis; funding acquisition; investigation; methodology; project administration; resources; supervision. Alice Simniceanu: Formal analysis; funding acquisition; investigation; methodology; project administration; resources; supervision. Alessandro Cassini: Formal analysis; methodology; project administration; resources; supervision. Rebecca Grant: Conceptualization; funding acquisition; methodology. Angel Rodriguez: Formal analysis; investigation; project administration; supervision. Andrea Vicari: Conceptualization; funding acquisition; investigation; methodology; project administration; resources; supervision. Lubna Al Arigi: Conceptualization; formal analysis; funding acquisition; investigation;

methodology; project administration; resources; supervision. Tasnim Azim: Conceptualization; funding acquisition; investigation; project administration; resources; supervision. Pushpa Ranjan Wijesinghe: Funding acquisition; investigation; project administration; supervision. Soatiana Cathycia Rajatonirina: Conceptualization; formal analysis; funding acquisition; investigation; project administration; resources; supervision. Joseph Chukwudi Okeibunor: Funding acquisition; investigation; project administration; resources; supervision. Linh-Vi Le: Formal analysis; funding acquisition; investigation; project administration; resources; supervision. Mark Katz: Formal analysis; investigation; methodology; project administration; supervision. Aisling Vaughan: Formal analysis; investigation; project administration; resources; supervision. Pernille Jorgensen: Formal analysis; investigation; project administration; supervision. Gudrun Freidl: Formal analysis; investigation; project administration; resources; supervision. Richard Pebody: Conceptualization; formal analysis; funding acquisition; investigation; methodology; project administration; supervision. Maria D van Kerkhove: Conceptualization; funding acquisition; methodology; project administration; supervision.

PEER REVIEW

The peer review history for this article is available at https://publons. com/publon/10.1111/irv.12915.

ORCID

Isabel Bergeri D https://orcid.org/0000-0002-1204-1753 Hannah C. Lewis D https://orcid.org/0000-0002-1925-3943 Lorenzo Subissi D https://orcid.org/0000-0001-5147-575X Anthony Nardone D https://orcid.org/0000-0003-1138-0937 Marta Valenciano D https://orcid.org/0000-0002-2056-1062 Brianna Cheng D https://orcid.org/0000-0003-2004-2251 Ketevan Glonti D https://orcid.org/0000-0001-9991-7991 Bridget Williams D https://orcid.org/0000-0002-9677-8305 Ibukun-Oluwa Omolade Abejirinde D https://orcid.org/0000-0003-0139-0541

Alice Simniceanu https://orcid.org/0000-0003-4068-6177 Rebecca Grant https://orcid.org/0000-0001-8412-8113 Angel Rodriguez https://orcid.org/0000-0002-0061-7567 Andrea Vicari https://orcid.org/0000-0001-5528-2517 Lubna Al Ariqi https://orcid.org/0000-0002-3343-6359 Tasnim Azim https://orcid.org/0000-0003-4127-5871 Pushpa Ranjan Wijesinghe https://orcid.org/0000-0002-7875-0165 Soatiana Cathycia Rajatonirina https://orcid.org/0000-0002-5869-5090

Joseph Chukwudi Okeibunor 🔟 https://orcid.org/0000-0002-6696-8503

Linh-Vi Le https://orcid.org/0000-0002-5816-7589 Mark Katz https://orcid.org/0000-0002-5532-7276 Aisling Vaughan https://orcid.org/0000-0002-5966-2303 Pernille Jorgensen https://orcid.org/0000-0001-5967-0998 Gudrun Freidl https://orcid.org/0000-0003-0702-0189 Richard Pebody https://orcid.org/0000-0002-9069-2885 Maria D. Van Kerkhove https://orcid.org/0000-0002-6135-0018

REFERENCES

- World Health Organisation. (2011). Final report of the IHR Review Committee (as presented to 64th WHA). Retrieved from https:// www.who.int/teams/ihr/ihr-review-committees/h1n1
- Consortium for the Standardization of Influenza Seroepidemiology (CONSISE). (n.d.). Protocols. Retrieved 2021, from CONSISE: https:// consise.tghn.org/protocols/
- World Health Organisation. (n.d.). Pandemic Influenza Special Investigations and Studies (IPSS). Retrieved from https://www.who.int/ influenza/surveillance_monitoring/influenza_pandemic_special_inves tigations/en/
- 4. World Health Organisation. (n.d.). Global influenza programme. Retrieved from https://www.who.int/influenza/about/en/
- World Health Organisation. (n.d.). Pandemic influenza preparedness (PIP) framework. Retrieved from https://www.who.int/influenza/ pip/en/
- World Health Organisation. (n.d.). Middle East respiratory syndrome coronavirus (MERS-CoV)/Technical guidance/Investigation tools. Retrieved 2021, from https://www.who.int/health-topics/middleeast-respiratory-syndrome-coronavirus-mers/technical-guidance/inv estigation-tools
- World Health Organisation. (2016). WHO Zika virus research agenda. Support and harmonization of research in countries: Standardized research protocols. Retrieved June 2021, from https://www.who. int/health-topics/zika-virus-disease/publications/who-response
- Van Kerkhove MD, Reveiz L, Souza JP, Jaenisch T, Carson G, Broutet N. Harmonisation of Zika virus research protocols to address key public health concerns. *Lancet Glob Health*. 2016 Dec;4(12): e911-e912.
- World Health Organisation. (n.d.). The unity studies: WHO seroepidemiological investigations protocols. Retrieved from https:// www.who.int/emergencies/diseases/novel-coronavirus-2019/ technical-guidance/early-investigations
- World Health Organisation. (2021, June). COVID-19 Strategy and planning. Retrieved from https://www.who.int/emergencies/ diseases/novel-coronavirus-2019/strategies-and-plans
- 11. World Health Organisation. (n.d.). Go.Data: managing complex data in outbreaks. Retrieved from https://www.who.int/godata
- Weidner L, Gänsdorfer S, Unterweger S, et al. Quantification of SARS-CoV-2 antibodies with eight commercially available immunoassays. J Clin Virol. 2020;129:104540.
- Kristiansen PA, Page M, Bernasconi V, et al. WHO International Standard for anti-SARS-CoV-2 immunoglobulin. *Lancet*. 2021, April; 397(10282):1347-1348. Retrieved from. https://www.nibsc.org/ products/brm_product_catalogue/detail_page.aspx?catid=20/136
- 14. World Health Organisation. (n.d.). SARS-CoV-2 antibody detecting assay performance evaluation. Retrieved from https://who.int/newsroom/articles-detail/who-sars-cov-2-antibody-detecting-assay-per formance-evaluation
- United Nations Office for the Coordination of Humanitarian Affairs (OCHA). (2020). Global humanitarian response plan for covid-19. Retrieved from https://www.unocha.org/sites/unocha/files/GHRP-COVID19_July_update.pdf

How to cite this article: Bergeri I, Lewis HC, Subissi L, et al. Early epidemiological investigations: World Health Organization UNITY protocols provide a standardized and timely international investigation framework during the COVID-19 pandemic. *Influenza Other Respi Viruses*. 2022;16 (1):7-13. doi:10.1111/irv.12915

WILEY