

Effects of *Pistacia atlantica subsp. kurdica* on Growth and Aflatoxin Production by *Aspergillus parasiticus*

Sadegh Khodavaisy,^{1,2} Sassan Rezaie,² Fatemeh Noorbakhsh,³ Elham Baghdadi,³ Somayah Sharifynia,² and Farzad Aala^{1*}

¹Department of Medical Mycology and Parasitology, Kurdistan University of Medical Sciences, Sanandaj, IR Iran

²Division of Molecular Biology, Department of Medical Mycology and Parasitology, Tehran University of Medical Sciences, Tehran, IR Iran

³Department of Microbiology, Faculty of Science, Islamic Azad University, Varamin-Pishva, IR Iran

*Corresponding author: Farzad Aala, Department of Medical Mycology and Parasitology, Kurdistan University of Medical Sciences, Sanandaj, IR Iran. Tel: +98-9197544944, E-mail: farzadala@yahoo.com

Received 2015 December 11; Revised 2016 February 14; Accepted 2016 April 18.

Abstract

Background: Aflatoxins are highly toxic secondary metabolites mainly produced by *Aspergillus parasiticus*. This species can contaminate a wide range of agricultural commodities, including cereals, peanuts, and crops in the field. In recent years, research on medicinal herbs, such as *Pistacia atlantica subsp. kurdica*, have led to reduced microbial growth, and these herbs also have a particular effect on the production of aflatoxins as carcinogenic compounds.

Objectives: In this study, we to examine *P. atlantica subsp. kurdica* as a natural compound used to inhibit the growth of *A. parasiticus* and to act as an anti-mycotoxin.

Materials and Methods: In vitro antifungal susceptibility testing of *P. atlantica subsp. kurdica* for *A. parasiticus* was performed according to CLSI document M38-A2. The rate of aflatoxin production was determined using the HPLC technique after exposure to different concentrations (62.5 - 125 mg/mL) of the gum. The changes in expression levels of the *aflR* gene were analyzed with a quantitative real-time PCR assay.

Results: The results showed that *P. atlantica subsp. kurdica* can inhibit *A. parasiticus* growth at a concentration of 125 mg/mL. HPLC results revealed a significant decrease in aflatoxin production with 125 mg/mL of *P. atlantica subsp. kurdica*, and AFL-B1 production was entirely inhibited. Based on quantitative real-time PCR results, the rate of *aflR* gene expression was significantly decreased after treatment with *P. atlantica subsp. kurdica*.

Conclusions: *Pistacia atlantica subsp. kurdica* has anti-toxic properties in addition to an inhibitory effect on *A. parasiticus* growth, and is able to decrease aflatoxin production effectively in a dose-dependent manner. Therefore, this herbal extract maybe considered a potential anti-mycotoxin agent in medicine or industrial agriculture.

Keywords: HPLC, Real-Time PCR, Aflatoxin, *Aspergillus parasiticus*, *Pistacia atlantica subsp. kurdica*

1. Background

Aflatoxins (AFs) are carcinogenic, mutagenic, and teratogenic metabolites mainly produced by *Aspergillus flavus* and the very closely related species *A. parasiticus* (1). These mycotoxins can grow on a wide range of food commodities, including cereals, peanuts, and crops in the field, but also during postharvest operations and storage (2). The group of AFs includes B1, B2, G1, and G2, while AFBs include AFB1 and AFB2, and AFGs include AFG1 and AFG2 (3). AFs are potent hepatotoxic, teratogenic, and mutagenic metabolites that are now well-recognized as a public health hazard (1, 4). Although some synthetic fungicides have been improved to protect crops against AF production, there are some proven indications that these are themselves hazardous to human health (5). Therefore, using natural sources is an affirmative approach to protecting from fun-

gal infections in agriculture (6).

A medicinal herb, *Pistacia atlantica subsp. kurdica*, has been traditionally used as a therapeutic agent for infections (7-9). The genus *Pistacia* belongs to the family Anacardiaceae, and among 15 known species of pistachios, only three grow in Iran: *P. vera*, *P. khinjuk*, and *P. atlantica* (10, 11). *P. atlantica* has three sub-species: *mutica*, *kurdica*, and *cabulica* (11). This plant is an evergreen bush that grows native in some southern and central American and eastern Mediterranean countries (10). *P. atlantica subsp. kurdica* thrives particularly around the Zagros mountains in Iran (10, 12). The gum of *P. atlantica subsp. kurdica* is obtained as an exudate after crushing the trunk and branches, and has numerous qualities. The gum of this plant is used in medicine, with many researchers reporting that it possesses considerable antimicrobial activity (12, 13).

2. Objectives

The aim of this study was to evaluate the effects of *P. atlantica subsp. kurdica*, as a natural compound, with regard to antifungal activity and inhibition of AF production in *A. parasiticus* infections.

3. Materials and Methods

3.1. Preparation of Gum

Industrial-grade gum of *P. atlantica* var. *kurdica* was obtained from Saghez Sazi (Van) (Kurdistan, Iran) in the summer of 2014. The gum was obtained as an exudate of the trunk and branches.

3.2. Fungal Strain and Susceptibility Test

The *A. parasiticus* strain (American type culture collection 15517) was incubated at 28°C for 72 hours on Sabouraud dextrose agar (Germany, Merck). In vitro antifungal susceptibility testing for the determination of minimum inhibitory concentration (MIC) was performed according to recommendations provided in the clinical and laboratory standards institute (CLSI) M38-A2 document, with some modification in gum concentration (14). RPMI 1640 medium (Sigma chemical Co.) was buffered to pH 7.0 with MOPS (Sigma), with L-glutamine and without bicarbonate. The gum of *P. atlantica* var. *kurdica* was dissolved in water to reach a concentration of 500 mg/mL. It was then diluted to achieve concentrations of 500 to 15.6 mg/mL (500, 250, 125, 62.5, 31.2, and 15.6 mg/mL). The concentration of fungal spores was calculated using the hemacytometry method, and finally adjusted to 2x concentrated suspensions in such a way that each test well contained 5×10^4 colony-forming units (CFUs)/mL. Next, 100 μ L of fungal suspension was added to the plates. Negative (only RPMI 1640 medium) and positive controls were also run alongside each experiment. The plates were incubated for 48 - 72 hours at 35°C. The MIC endpoints were determined as the lowest concentrations that prevented any recognizable growth, mostly based on the antifungal capacity of the gum (100% inhibition).

3.3. Detection of AFs

The amount of AFs produced by *A. parasiticus* alone and in combination with different gum concentrations (62.5 mg/mL and 125 mg/mL) was determined using the high-performance liquid chromatography (HPLC) technique (Waters 474 scanning fluorescence detector) (6). PDB plates containing 62.5 mg/mL or 125 mg/mL of gum were inoculated with 5×10^4 fungal spores. The medium plates

were incubated at 30°C for 7 days without agitation. Positive and negative controls were also run along with the test plates. Afterward, the obtained fungal biomasses were harvested and weighed. Then, 25 g of each fungal biomass with 2.5 g of sodium chloride was added into 100 mL of HPLC-grade methanol and blended for 3 minutes at 18,000 rpm. Cell-free filtrates were obtained using Whatman filter paper No.4 and subjected to analysis by HPLC.

3.4. *AflR* Gene Expression Assay

Quantitative changes in expression of the *aflR* gene were analyzed by use of a quantitative real-time PCR assay. *A. parasiticus* was cultured in the presence of 62.5 mg/mL or 125 mg/mL of gum of *P. atlantica* var. *kurdica* to be used for RNA extraction (6). RNA concentrations were determined spectrophotometrically (Biophotometer, Eppendorf, Germany). Equal amounts of RNA (1 μ g in 20 μ L) were subjected to cDNA synthesis by using random hexamer primers, according to the manual protocol (Fermentase). The primers Faflr (5'-CGGAACAGGGACTTCCGGCG-3') and Raflr (5'-GGGTGGCGGGGACTCTGAT-3') were designed on the basis of the published sequence of the *aflR* gene of *A. parasiticus* (NCBI, accession no: AF441438). The b-actin gene (ACT1) with Fact (5'-ACGGTATTGTTTCCAACACTGGGACG-3') and Ract (5'-TGGAGCTTCGGTCAACAAAAGTGG-3') primers was used as an endogenous reference gene. Quantitative real-time PCR was carried out using the SYBR green master mix (Applied Biosystems, Foster City, CA, USA) performed with a step one plus real-time PCR system (Applied Biosystems, Foster City, CA, USA). The PCR setup and program have been previously described (6). The *aflR* gene expression was analyzed with REST software (2008 V2.0.7, Corbett research, Sydney, Australia).

4. Results

Our results demonstrated that the gum of *P. atlantica subsp. kurdica* inhibited *A. parasiticus* growth at MIC values of 125 mg/mL. According to HPLC analysis results, AFL-B2 and AFL-G2 production was significantly decreased and even entirely inhibited in 125 mg/mL of gum ($P < 0.01$). However, the production of AFL-B1 was not fully inhibited at this concentration. The inhibitory effect of the gum on AF production is shown in Table 1. Based on quantitative real-time PCR results, the rate of *aflR* gene expression was significantly decreased after treating the *A. parasiticus* with 125 mg/mL of *P. atlantica subsp. kurdica* ($P < 0.05$). The results of an analysis of the relative quantification of the *aflR* gene after treatment with the gum of *P. atlantica subsp. kurdica* are shown in Table 2.

Table 1. Effect of Concentrations of *P. atlantica subsp. kurdica* Gum on AF Production

Sample	AFL-B1	AFL-B2	AFL-G1	AFL-G2	Total
Negative control	ND	ND	ND	ND	ND
Positive control	4254.10	1521.0	9561.0	1510.1	16846.2
PDA containing standard toxins	21.16	2.34	11.71	2.08	37.29
<i>A. parasiticus</i> exposed to 62.5 mg/mL of gum	110.5	4.55	4.27	ND	119.32
<i>A. parasiticus</i> exposed to 125 mg/mL of gum	6.5	ND	ND	ND	6.5

Table 2. Relative Expression of the *aflR* Gene Using Real-Time PCR Analysis

Gene	Type	Reaction Efficiency	Expression	Std. Error	95%CI	Result
Act beta	REF	1.0	1.000	-	-	-
<i>aflR</i> (62.5 mg/mL)	TRG	1.0	0.165	0.016 - 0.451	0.08 - 0.41	Down expression
<i>aflR</i> (125 mg/mL)	TRG	1.0	0.212	0.111 - 0.823	0.043 - 0.55	Down expression




Figure 1. Effects of *P. atlantica subsp. kurdica* Gum on *aflR* Gene Expression

5. Discussion

Aflatoxins are extremely toxic and carcinogenic metabolites that widely contaminate agricultural commodities and animal feed (15). In recent years, research has been ongoing into the development of new anti-fungal agents or through the search for natural sources to control growth and toxin production of *Aspergillus* species (8, 9, 16). In several studies, the antimicrobial activity of *P. atlantica* was identified. Ghalem et al. reported that the gum of *P. atlantica* has antimicrobial activity against gram-positive and -negative bacteria that are resistant to commonly used antimicrobial agents, and this was considerably dependent on concentration (17). Tassou et al. showed that the addition of *P. lentiscus* var. chia

gum inhibited the growth of organisms (13). In this study, the data show that the *P. atlantica subsp. kurdica* had an inhibitory effect on *A. parasiticus*. These results agree with the conclusion of previous works (7, 18). Hesami et al. suggested the potential substitution of antifungal chemicals by *P. atlantica subsp. kurdica* as a natural inhibitor to control the growth of *Botrytis cinerea* (19). Aflatoxin inhibition mainly occurs during the primary stages of cell growth, when the genes involved in aflatoxin biosynthesis are at maximum activity within the fungal cells (7).

Our study revealed that the rate of *aflR* gene expression was significantly decreased after treating the fungus with 125 mg/mL of *P. atlantica subsp. kurdica*. Mohseni et al. showed that licorice extract could efficiently inhibit *aflR* gene expression, and consequently, AF production, in *A. parasiticus* (6). In another study, Bagheri-Gavkosh et al. showed potent inhibition of *A. parasiticus* growth and AF production with 1 mg/mL of *Ephedra major* (20). Overall, we concluded that the gum of *P. atlantica subsp. kurdica* could efficiently inhibit *aflR* gene expression and the consequent AF production in *A. parasiticus*. This gum may therefore be considered a potential anti-mycotoxin agent in medicine or industrial agriculture. Further studies are recommended to determine the effective components of the gum of *P. atlantica subsp. kurdica* and to consider it as a potential candidate for controlling AF contamination of crops in the field.

Acknowledgments

This study was financially supported by a grant from the Kurdistan University of Medical Sciences, Iran. We are

also grateful to the Saghez Sazi (Van) company (Kurdistan, Iran).

Footnotes

Authors' Contribution: Study concept and design, Sassan Rezaie, Sadegh Khodavaisy; acquisition of data, Sassan Rezaie, Farzad Aala; analysis and interpretation of data, Sadegh Khodavaisy, Somayeh Sharifynia, Elham Baghdadi; drafting of the manuscript, Sassan Rezaie; critical revision of the manuscript for important intellectual content, Sassan Rezaie; statistical analysis, Sadegh Khodavaisy; administrative, technical, and material support, Sadegh Khodavaisy, Fatemeh Noorbakhsh, Elham Baghdadi; study supervision, Sassan Rezaie.

Conflict of Interest: The authors declare no potential conflicts of interest.

Funding/Support: The present work was financially supported by Kurdistan University of Medical Sciences, Iran.

References

- Kostarelou P, Kanapitsas A, Pyrri I, Kapsanaki-Gotsi E, Markaki P. Aflatoxin b1 production by aspergillus parasiticus and strains of aspergillus section nigri in currants of greek origin. *Food Control*. 2014;**43**:121-8. doi: [10.1016/j.foodcont.2014.02.011](https://doi.org/10.1016/j.foodcont.2014.02.011).
- Davari B, Khodavaisy S, Ala F. Isolation of fungi from housefly (*Musca domestica* L.) at Slaughter House and Hospital in Sanandaj, Iran. *Journal of preventive medicine and hygiene*. 2012;**53**(3):172-4. [PubMed: [23362625](https://pubmed.ncbi.nlm.nih.gov/23362625/)].
- Shephard G, Berthiller F, Burdaspal P, Crews C, Jonker M, Krska R, et al. Developments in mycotoxin analysis: an update for 2010-2011. *World Mycotoxin J*. 2012;**5**(1):3-30. doi: [10.3920/wmj2011.1338](https://doi.org/10.3920/wmj2011.1338).
- Khayoon W, Saad B, Yan C, Hashim NH, Ali ASM, Salleh MI, et al. Determination of aflatoxins in animal feeds by HPLC with multi-functional column clean-up. *Food Chemistry*. 2010;**118**(3):882-6. doi: [10.1016/j.foodchem.2009.05.082](https://doi.org/10.1016/j.foodchem.2009.05.082).
- Tripathi P, Dubey NK. Exploitation of natural products as an alternative strategy to control postharvest fungal rotting of fruit and vegetables. *Postharvest Biol Tec*. 2004;**32**(3):235-45. doi: [10.1016/j.postharvbio.2003.11.005](https://doi.org/10.1016/j.postharvbio.2003.11.005).
- Mohseni R, Noorbakhsh F, Moazeni M, Nasrollahi Omran A, Rezaie S. Antitoxin Characteristic of Licorice Extract: The Inhibitory Effect on Aflatoxin Production in *Aspergillus parasiticus*. *Food Safety*. 2014;**34**(2):119-25. doi: [10.1111/jfs.12104](https://doi.org/10.1111/jfs.12104).
- Rasooli I, Abyaneh MR. Inhibitory effects of Thyme oils on growth and aflatoxin production by *Aspergillus parasiticus*. *Food Control*. 2004;**15**(6):479-83. doi: [10.1016/j.foodcont.2003.07.002](https://doi.org/10.1016/j.foodcont.2003.07.002).
- Farag RS, Daw ZY, Abo-Raya SH. Influence of some spice essential oils on *aspergillus parasiticus* growth and production of aflatoxins in a synthetic medium. *Food Science*. 1989;**54**(1):74-6. doi: [10.1111/j.1365-2621.1989.tb08571.x](https://doi.org/10.1111/j.1365-2621.1989.tb08571.x).
- Karapinar M. Inhibitory effects of anethole and eugenol on the growth and toxin production of *Aspergillus parasiticus*. *Int J Food Microbiol*. 1990;**10**(3-4):193-9. [PubMed: [2397152](https://pubmed.ncbi.nlm.nih.gov/2397152/)].
- Sharifi M, Hazell SL. GC-MS Analysis and Antimicrobial activity of the essential oil of the trunk exudates from *Pistacia atlantica kurdica*. *Indian J Pharm Sci*. 2011;**3**(2):1364-7.
- Behboodi B. Ecological distribution study of wild pistachios for selection of rootstock. *Options Mediterr Ser A*. 2003;**63**:61-7.
- Taran M, Mohebbali M, Esmaeli J. In vivo efficacy of gum obtained *pistacia atlantica* in experimental treatment of cutaneous leishmaniasis. *Iran J Public Health*. 2010;**39**(1):36-41. [PubMed: [23112988](https://pubmed.ncbi.nlm.nih.gov/23112988/)].
- Tassou CC, Nychas GJE. Antimicrobial activity of the essential oil of mastic gum (*Pistacia lentiscus* var. *chia*) on Gram positive and Gram negative bacteria in broth and in Model Food System. *Int Biodeter Biodegr*. 1995;**36**(3-4):411-20. doi: [10.1016/0964-8305\(95\)00103-4](https://doi.org/10.1016/0964-8305(95)00103-4).
- CLSI. Reference method for broth dilution antifungal susceptibility testing of filamentous fungi: approved standard. 2008.
- Rasooli I, Owlia P. Chemoprevention by thyme oils of *Aspergillus parasiticus* growth and aflatoxin production. *Phytochemistry*. 2005;**66**(24):2851-6. doi: [10.1016/j.phytochem.2005.09.029](https://doi.org/10.1016/j.phytochem.2005.09.029). [PubMed: [16289146](https://pubmed.ncbi.nlm.nih.gov/16289146/)].
- Badii F, Moss MO. The effect of the fungicides tridemorph, fenpropimorph and fenarimol on growth and aflatoxin production by *Aspergillus parasiticus* Speare. *Lett Appl Microbiol*. 1988;**7**(2):37-9. doi: [10.1111/j.1472-765X.1988.tb01247.x](https://doi.org/10.1111/j.1472-765X.1988.tb01247.x).
- Ghalem BR, Mohamed B. Essential oil from gum of *Pistacia atlantica* Desf.: screening of antimicrobial activity. *J Pharm Pharmacol*. 2009;**3**(1):13-5.
- Mohseni R, Nasrollahi Omran A, Noorbakhsh F, Rezaie S, Hosseinjani H. A survey of the effect of licorice plant extract on aflr gene expression and aflatoxin production in *aspergillus parasiticus* via real-time pcr. *MJMS*. 2012;**15**(3):63-77.
- Hesami G, Bahramian S, Fatemi A, Hesamia S. Effect of *pistacia atlantica* subsp. *kurdica* essential oil and acetic acid on *botrytis cinerea* growth in culture media and strawberry fruits. *Bull Env Pharmacol. Life Sci*. 2014;**3**(2):100-6.
- Bagheri-Gavkosh S, Bigdeli M, Shams-Ghahfarokhi M, Razzaghi-Abyaneh M. Inhibitory effects of *Ephedra major* Host on *Aspergillus parasiticus* growth and aflatoxin production. *Mycopathologia*. 2009;**168**(5):249-55. doi: [10.1007/s11046-009-9220-x](https://doi.org/10.1007/s11046-009-9220-x). [PubMed: [19557546](https://pubmed.ncbi.nlm.nih.gov/19557546/)].