

©Suárez-Díaz S. Caminal-Montero I.

This work is licensed under a Creative Commons Attribution 4.0 International License.



LETTER TO THE EDITOR

Overlapping between IgG4-RD and Behçet's Disease

Silvia Suárez-Díaz1, Luis Caminal-Montero2

¹Internal Medicine Department, Hospital Universitario Central de Asturias, Oviedo, Spain, ²Systemic Autoinmune Disease Unit, Internal Medicine Department, Instituto de Investigación Sanitaria del Principado de Asturias (ISPA), Hospital Universitario Central de Asturias, Oviedo, Spain

Mediterr J Rheumatol 2020;31(1):92 https://doi.org/10.31138/mjr.31.1.92

Article Submitted: 19 Jan 2020; Revised Form: 30 Jan 2020; Article Accepted: 14 Feb 2020; Published: 31 Mar 2020

Keywords: IgG4 Related Disease, Behcet's Disease, Overlap syndrome

To the editor:

Alazani MB et al.¹ reported for the first time a case of IgG4-Related disease (IgG4-RD) retroperitoneal mass in a patient with Behçet's Disease (BD), and they found only one other similar patient with BD, and in that case, an IgG4-RD laryngeal mass.² Similarly, we have previously described another patient with recurrent aphthous stomatitis and IgG4-related laryngitis with the suspected diagnosis of overlapping with BD.³

Some conditions that were previously thought to be unique clinical syndrome are now recognized as a clinical manifestation of IgG4-RD. IgG4-RD physiopathological basis is the presentation of autoantigens by plasmablasts or B cells to CD4 cytotoxic T cells, which produce pro-fibrotic cytokines such as IFN-γ, IL1-β and TGF- β.⁴ Although BD shares some common features with autoimmune and autoinflammatory diseases, is ultimately caused by disturbance of T-cell homeostasis, especially Th1 and Th17 expansion, as well as Tregs response suppression. Neutrophil activity is increased at the earliest stage of inflammation in the affected organs, as well as the presence of HLA-B*5 and increased IL-17, which appears to play a major role in neutrophil activity.⁵ IgG4-RD is a rare, multiorgan condition characterized by histologic features, and ac-

Corresponding Author:

Silvia Suárez-Díaz Internal Medicine Department, Hospital Universitario Central de Asturias Oviedo 33011, Spain Tel.: +34 649 628 548 E-mail: silvia_porto@hotmail.es curate clinical diagnosis can be challenging due to the atypical manifestations or the presence of overlapping autoimmune diseases like BD or antineutrophil cytoplasmic antibody-associated vasculitides.⁶

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- Alanazi MB, Asiri YO, Al-Homood IA. IgG4-Related Disease coexisting with Behçet's Disease. Mediterr J Rheumatol 2019;30(4):228-30. [https://doi.org/10.31138/mjr.30.4.228]
- Shaib Y, Ton E, Goldschmeding R, Tekstra J. IgG4-related disease with atypical laryngeal presentation and Behcet/granulomatous polyangiitis mimicking features. BMJ Case Rep 2013;pii: bcr2013009158. [https://doi.org/10.1136/bcr-2013-009158] [PMID: 23813505] [PMCID: PMC3703034]
- Suárez-Díaz S, Núñez-Batalla F, Fernández-García MS, Fernández-Llana MB, Yllera-Gutiérrez C, Caminal-Montero C. Aphthous Stomatitis and Laryngitis, Another Form of Presentation of an IgG4-Related Disease? [Article in English, Spanish] Reumatol Clin 2018;pii: S1699-258X(18)30191-8. [https://doi.org/10.1016/j.reuma.2018.08.011] [PMID: 30297197]
- Zhang W, Stone JH. Management of IgG4-related disease. Lancet Rheumatol 2019;1(1):e55-65. [https://doi.org/10.1016/S2665-9913(19)30017-7]
- Leccese P, Alpsoy E. Behçet's Disease: An Overview of Etiopathogenesis. Front Immunol 2019;10:1067. [https://doi.org/10.3389/fimmu.2019.01067] [PMID: 31134098] [PMCID: PMC6523006]
- Danlos FX, Rossi GM, Blockmans D, Emmi G, Kronbichler A, Durupt S, Maynard C, et al. Antineutrophil cytoplasmic antibody-associated vasculitides and IgG4-related disease: A new overlap syndrome. Autoimmun Rev 2017;16(10):1036-43. [https://doi.org/10.1016/j.autrev.2017.07.020] [PMID: 28780079]