



Review

The Multifaceted Roles of Ku70/80

Sayma Zahid, Murielle Seif El Dahan, Florence Iehl, Paloma Fernandez-Varela , Marie-Helene Le Du, Virginie Ropars and Jean Baptiste Charbonnier *

Institute for Integrative Biology of the Cell (I2BC), Université Paris-Saclay, CEA, CNRS, 91198 Gif-sur-Yvette, France; sayma.zahid@i2bc.paris-saclay.fr (S.Z.); murielle.seif-el-dahan@i2bc.paris-saclay.fr (M.S.E.D.); florence.iehl@evotec.com (F.I.); paloma.fernandez-varela@i2bc.paris-saclay.fr (P.F.-V.); marie-helene.ledu@i2bc.paris-saclay.fr (M.-H.L.D.); virginie.ropars@i2bc.paris-saclay.fr (V.R.)

* Correspondence: jb.charbonnier@i2bc.paris-saclay.fr

Abstract: DNA double-strand breaks (DSBs) are accidental lesions generated by various endogenous or exogenous stresses. DSBs are also genetically programmed events during the V(D)J recombination process, meiosis, or other genome rearrangements, and they are intentionally generated to kill cancer during chemo- and radiotherapy. Most DSBs are processed in mammalian cells by the classical nonhomologous end-joining (c-NHEJ) pathway. Understanding the molecular basis of c-NHEJ has major outcomes in several fields, including radiobiology, cancer therapy, immune disease, and genome editing. The heterodimer Ku70/80 (Ku) is a central actor of the c-NHEJ as it rapidly recognizes broken DNA ends in the cell and protects them from nuclease activity. It subsequently recruits many c-NHEJ effectors, including nucleases, polymerases, and the DNA ligase 4 complex. Beyond its DNA repair function, Ku is also involved in several other DNA metabolism processes. Here, we review the structural and functional data on the DNA and RNA recognition properties of Ku implicated in DNA repair and in telomeres maintenance.



Citation: Zahid, S.; Seif El Dahan, M.; Iehl, F.; Fernandez-Varela, P.; Le Du, M.-H.; Ropars, V.; Charbonnier, J.B. The Multifaceted Roles of Ku70/80. *Int. J. Mol. Sci.* **2021**, *22*, 4134. <https://doi.org/10.3390/ijms22084134>

Academic Editors: Joanna Timmins and Jean-Luc Ravanat

Received: 4 March 2021
Accepted: 1 April 2021
Published: 16 April 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: double-strand break; c-NHEJ; telomeres; protein-DNA interactions; DNA repair machinery

1. Introduction

Living organisms are constantly exposed to genotoxic stress that can lead to double-strand breaks (DSBs). These DNA lesions can be of endogenous origin, such as cellular respiration and reactive oxygen. They can also be of exogenous origin such as ionizing radiation, pharmaceutical drugs, or chemical exposure [1] DSBs are repaired by two main DNA repair pathways: Homologous recombination (HR) and classical nonhomologous end-joining (c-NHEJ). The c-NHEJ is the prominent DSB repair pathway in multicellular eukaryotes, while HR is predominant in *S. cerevisiae*. The c-NHEJ pathway accounts for most DSB repair outside of the S and G2 phases [2,3].

The Ku70/80 complex was first identified in 1981 by Mimori et al. as a high-molecular-weight nuclear protein and as an antigen recognized by auto-antibodies in sera from patients with polymyositis-scleroderma diseases [4]. It was called Ku from the first two letters of the original patient's name whose serum was used as a prototype to identify and purify the heterodimer. The heterodimer Ku70/80 (called Ku hereafter) is a central actor of the c-NHEJ pathway. It is considered as one of the first factors to be recruited at DSB sites because of its high affinity for DNA ends (Kd in the nM range) and its high abundance in the nucleus [5]. Ku recognizes a large variety of DNA ends including blunt ends, hairpin DNA, or ends with protruding single-stranded overhangs (discussed in detail later on). Via its C-terminal region, Ku interacts with the DNA-dependent protein kinase catalytic subunit (DNA-PKcs) to form the DNA-PK holoenzyme, which controls the interaction between broken DNA ends during c-NHEJ [2–4]. Ku has a key role in c-NHEJ beyond the ends recognition step. It directly or indirectly recruits most c-NHEJ effectors, including ends processing by nucleases, kinases, and phosphatases; nucleotide addition by DNA

Pol X polymerases; and ligation by the XRCC4/Lig4 complex and the XLF and PAXX scaffolding proteins [5,6]. In this review, we mainly focus on the molecular basis of the interactions of Ku with various DNA ends observed during DSB formation and on its interactions with RNA and DNA substrates at telomeres.

The molecular basis of the protein interactions between Ku and the c-NHEJ factors are reviewed elsewhere [7–9]. Briefly, recent structural studies characterized interaction sites of Ku with c-NHEJ proteins. The structure of the DNA-PK, formed by Ku and DNA-PKcs, was determined by cryo-electron microscopy (cryoEM) by several groups (for example, [10,11]). Recently, Chaplin et al. identified particles corresponding to a dimer of DNA-PK holoenzymes that reconstitute a synapse of DNA ends. This dimer is mediated by a domain swap mechanism of the conserved C-terminal helix of Ku80. In this dimer, DNA ends present in each DNA-PK run almost parallel [11]. Ku can also recruit, at DSBs, several c-NHEJ factors containing motifs called Ku-binding motifs (KBMs) [12,13]. These KBMs were identified in the nuclease APLF, the ligation factors XLF (XRCC4-like factor) and PAXX (Paralog of XRCC4 and XLF), the regulation factor MRI/CYREN (MRI for Modulator of Retrovirus Infection, CYREN for Cell Cycle Regulator Of Nonhomologous End Joining), and the helicase/nuclease WRN protein [12,13]. Additional direct protein interactions were also reported with the Pol X polymerases, pol mu and TdT, and with the LIG4/XRCC4 complex [12]. Ku is thus a major hub of the c-NHEJ with key functions in DSB recognition and in coordination of the c-NHEJ enzymatic activities.

2. Ku70/80, an Abundant Nucleus Factor with High Affinity for DNA Ends

In human cells, Ku70 (69.9 kDa, 609 aa) and Ku80 (82.7 kDa, 732 aa) assemble to form a heterodimer. The role of Ku in DNA repair was first established experimentally when it was discovered that the Ku80 subunit was defective in X-ray-sensitive rodent cell mutants in the XRCC5 group (X-ray repair cross-complementing group 5) [7–9]. The same observation was made in embryonic stem cells XRCC6 defective for Ku70 [14]. The Ku70/80 heterodimer is one of the most abundant nuclear factors (about 500,000 copies by cell), and this corresponds to a cellular concentration of about 1.5 μM if we consider a nucleus with a radius of 5 μm . The Ku70 and Ku80 subunits are mainly addressed to the nucleus through the presence of a nuclear localization signal (NLS) present in the C-terminal region of each subunit (NLS in position (539–556) in Ku70 and in position (561–569) in Ku80) [15,16]. The crystal structures of the Importin α in a complex with the NLS of Ku70 or Ku80 were reported [17]. Both NLSs function as monopartite NLSs with only one basic cluster in the interaction with the importin. Mutagenesis of the NLS showed that each subunit can be imported via its own NLS or the NLS in the other subunit [18]. Ku is also present in the cytoplasm and at the membrane where it exerts several functions, including an anti-apoptotic activity by sequestering Bax (see [19] for a review).

In 1986, Mimori et al. unveiled in a seminal study the basic characteristics of the Ku protein regarding its DNA binding properties [20]. After the purification of Ku from HeLa extracts by immunoabsorbent column chromatography, they showed that Ku binds double-strand DNA (dsDNA) more efficiently than single-strand DNA (ssDNA), and they identified an interaction with dsDNA that is salt-dependent. They reported that Ku efficiently binds a linearized plasmid but not the circular form. Using restriction enzymes, they observed that the interaction is proportional to the number of cuts and is independent of the sequences at the cleavage sites. From all these data, they proposed that Ku interacts tightly with DNA ends in a sequence-independent manner [20]. The authors made the hypothesis that Ku may be “associated with DNA break points that might occur through the action of such agents as UV light or certain drugs” and thus have a role in DNA damage repair.

3. Affinities of Ku for DNA Measured by Biochemical and Biophysical Approaches

Several additional studies further documented the DNA binding properties of Ku. The interaction of Ku with short dsDNA (in the 20–200 bp range) was measured by different

approaches. Initially, most studies used EMSA (electrophoretic mobility shift assay), where the interaction between Ku and DNA results in characteristic “ladder” profiles. Different species with reduced mobilities form a ladder, with each band corresponding to a different number of Ku molecules bound on the DNA used. The EMSA assays indicate that one Ku molecule is loaded for every 30 bp of a DNA duplex and that Ku can translocate internally, leaving a place for new Ku molecules at the ends [21]. The first electron microscopy analyses of Ku on DNA confirmed the ability of Ku to thread on DNA and fully cover a DNA duplex [22].

The apparent K_d measured by EMSA with purified Ku and short dsDNA fragments (16–24 bp) ranged from 30 pM to 2 nM according to the experimental conditions [21,23]. The dissociation constants were further characterized using double-filter determination and anisotropy fluorescence [24]. Similar K_d values were measured in the 4–7 nM range. Recently, we used calorimetry to measure the interaction between Ku and different DNA substrates [25–27]. We obtained dissociation constants of 3.5 and 2.4 nM between Ku and two DNA duplexes of 18 and 42 bp, respectively, in good agreement with the previous values reported. The calorimetry gives access to the stoichiometry of the interaction. By titrating Ku with the 18 and 42 bp, we measured DNA:Ku stoichiometries of 0.97 and 0.47, respectively, which is in good agreement with the one and two Ku molecules bound on these substrates, respectively. Ku presents a slighter preference for DNA ends that are A/T-rich compared to G/C-rich, suggesting that some unwinding of the DNA ends may facilitate the interaction [23].

A helicase activity was initially proposed for Ku [28] with heterodimers purified from HeLa cells or *E. coli*. The absence of the motor domain and ATPase site in Ku suggests that it could correspond to a contaminant helicase co-purified with Ku. Indeed, Ku exhibits some affinity for ss oligos but in a much lower affinity range than for dsDNA [23,29,30]. Nevertheless, high concentrations of ssDNA can divert Ku from DSBs and favor Ku-independent end-joining at the expense of c-NHEJ [30]. Ku binding to single-stranded DNA (ssDNA) requires a free DNA end, a minimal length of 10 nucleotides with an optimal length of 25 nucleotides, and a preference for ssDNA with a free 5'-end [30]. Ku can also bind ssDNA once, forming a hairpin structure [31].

Ku70 and Ku80 are predominantly observed as a heterodimer when expressed in *E. coli* or in insect cells. Wang et al. showed that purified Ku70/80 and Ku70 expressed in rabbit kidney cells (RK13) bind dsDNA with a similar affinity and showed a similar sensitivity to high-ionic-strength buffers [32]. More recently, homodimers of Ku70 were isolated from the expression in *E. coli* and as a byproduct of the purification of Ku from insect cells [33]. The Ku70 homodimer stably binds 50 bp dsDNA substrate-forming complexes observed as two bands with an apparent lower affinity than the Ku heterodimer [33]. Ku70 and Ku80 subunits were separately expressed in vitro using rabbit reticulocytes lysate, and no DNA binding was observed with these isolated subunits [34].

4. Kinetic Analyses of Ku Binding on DNA

Kinetic analyses with DNA of increasing length (18 to 90 bp) were reported by Ma et al. using SPR (surface plasmon resonance) [35]. The authors showed that Ku binds a short DNA (18 bp) with a fast association rate (k_{on}) of $7.2 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$ and a dissociation (k_{off}) rate of $1.0 \times 10^{-2} \text{ s}^{-1}$, corresponding to a K_d of 1.4 nM. Similar kinetic constants were reported with Ku from *S. cerevisiae* using SPR [36]. A cooperative binding was observed for two sites (45 bp), suggesting that the first Ku molecule positioned at the DNA end favors the recruitment of the second Ku molecule. The use of DNA with 3 and 4 sites (68 and 90 bp) does not show additional cooperativity. Recently, some kinetic data were reported using SPR and the differences in Förster resonance energy transfer (FRET) with Ku70 and Ku80, respectively, tagged with an ECFP or EYFP protein in their N-terminus [37]. The SPR results were in the same range than previous measurements. Using FRET, a longer time release was observed with a k_{off} of $1.4 \times 10^{-4} \text{ s}^{-1}$, and the authors observed that a

deletion mutant of the SAP domain (SAF-A/B, Acinus, and PIAS) of the Ku70 Cter region rapidly dissociates, resulting in a 20-fold reduced affinity for DNA [37].

5. Ku Recognizes a Large Variety of DNA Ends

An important issue regarding the DNA ends recognition by Ku is the capacity of the heterodimer to recognize a wide spectrum of DSBs with different DNA ends structures. DSBs are generated by various endogenous and exogenous agents that result in chemically heterogeneous structures and chemical groups at DNA ends (reviewed in [38]) (Figure 1a). Using restriction enzymes generating blunt or 5'-or 3'-overhang DNA ends, Mimori et al. showed that the affinity of Ku is independent of the presence of 3–4 nucleotides overhangs [20]. This was confirmed by Falzon et al., who observed a tight interaction with a DNA containing an overhang of 21 nt on the four ends of a dsDNA [23]. Tadi et al. observed that Ku can accommodate DNA large single-strand overhangs in 3' or 5' [33] (Figure 1c,d). Ku can also accommodate hairpin structures (Figure 1e). Affinities similar to blunt DNA were measured with DNA containing hairpins of 4 to 20 nucleotides at both ends [23,24]. It is still unclear how the Ku ring structure can adapt to this variety of DNA ends.




Figure 1. Diversity of DNA ends that are recognized by Ku70/80. (a) Ku binds DNA ends efficiently with strands containing either hydroxyl or phosphate functions, as well as more heterogeneous ends with dRP or other chemical adducts. (b) Ku recognizes DSBs with a lesion present in the internal position close to the DNA ends. (c,d) Ku interacts efficiently with DNA having single-strand overhangs in 3' and/or in 5' up to 20 nt. (e) Ku can accommodate the presence of loops up to at least 8 nt at the ends. (f) Ku binds RNA hairpins at telomeres. (g,h) Ku interacts with the DNA–RNA duplex during replication fork restart (nascent RNA (green arrow)). (i) Ku binds single-ended DSB.

Ku also interacts with DNA ends, presenting DNA damage at proximity. Turchi et al. showed that Ku can bind a 75 mer DNA treated with Cisplatin (CisPt) (Figure 1b), although under this condition, it does not activate DNA-PKcs [39], contrary to what is observed in the absence of DNA damage close to the DSB ends [40]. Ku interacts with a DNA duplex containing an abasic site close to the DNA ends, and it was identified as the main protein trapped with the crosslinking reagent and DNA containing AP sites in the cellular extract [41]. In agreement with this observation, 5'dRP/AP lyase activity was reported for Ku. This activity is proposed to “clean” AP sites located close to DSBs and, thus, contribute to the DSB processing [42]. Ku lyase activity has a preference for the AP site located on a 5' overhang. Mutations of three Ku70 lysines (K31, K160, and K164) located on the inner part

of the ring lead to a major reduction of this activity [43]. The AP lyase activity was also characterized in two bacterial Ku proteins [44].

6. A Unique Pre-Formed Ring Structure among DNA Binding Proteins

A major advance in the understanding of the Ku mechanism was the crystal structure of the heterodimer in the complex with hairpin DNA in 2001 [45]. A key observation was the ring-shaped structure of the heterodimer surrounding the DNA (Figure 2). The crystal contained a DNA with a 14 bp duplex region followed by a 13 nt hairpin to limit Ku translocation during complex formation and crystallization. The protein was formed by three domains: An *N*-terminal α/β domain, a central-barrel domain, and a helical *C*-terminal arm (Figure 2). Ku70 and Ku80 have a *C*-terminal region (SAP domain (7 kDa) in Ku70 and a Cter of 17 kDa in Ku80). The Ku70 SAP domain is present in both crystals of Ku free (apo-Ku) and Ku complex, but it is only visible in the crystal structure of apo-Ku where it is in contact with the Ku80 α/β domain. Using electron microscopy, Rivera-Calzada mapped these *C*-terminal regions and their conformational changes after DNA and DNA-PKcs binding on maps at low resolution [46]. Recently, origami DNA were used to visualize, by EM, one or two Ku molecules bound at DNA ends specifically positioned on the origami [47].




Figure 2. Structure of the human Ku70/80 heterodimer. (a) Human Ku70 and Ku80 have similar organizations. They present some sequence variabilities in their *C*-terminal region. (b,c) Crystal structure of Ku70/80 complexed with a hairpin DNA and the Ku binding motif of APLF (magenta) [26]. Ku is colored in the same way as that in (a). The right view shows the top view of the complex. The DNA is a hairpin DNA used to limit Ku movement on the DNA for crystallization. (d) Structure of the SAP domain of Ku70 solved by NMR [48]. (e) *C*-terminal domain of Ku80 solved by NMR [49,50].

The Ku ring is formed by a large base that cradles DNA and a narrow bridge formed by β -strands from each subunit (Figure 2b,c). The N-terminal α/β domains are positioned at a remote distance from the ring structure. The central rounded β -barrel domains of Ku70 and Ku80 contribute to the dimer interface, and each contain a ~70 residues insertion that forms the two narrow bridges and the ring-shaped structure of Ku. The extremities of these insertions are buried in the Ku70 and Ku80 subunits so that their opening is highly unlikely and would need a large unfolding of the Ku subunits. Ku has an asymmetric orientation on DNA ends. Yoo et al. used cross-linking experiments and characterized a preferential orientation of the heterodimer. The Ku70 subunit is located proximal to the DNA end, and Ku80 is located toward internal DNA nucleotides [51]. The crystal structure thus reveals a preformed ring structure that fits well with the specific DNA ends recognition properties of Ku. Ku covers the 14 bp duplex in the crystal structure, but molecular modeling from this structure suggests that about 25 bp could bind to the large Ku cradle. This ring structure with two interdigitated insertions raised the question of the unloading of Ku after DNA ligation and was recently addressed (see below).

7. The Ku70/80 Has an Apparently Rigid Inner Face

The crystal structure of the Ku/DNA complex shows that several loops of Ku70 and Ku80 are in contact with the minor and major grooves of the DNA. The inner dimension of the Ku ring is thus smaller than the external radius of the DNA helix, similar to a screw fixed in a nut (Figure 3a,b). These loops superimpose well in the apo structure of Ku or in a complex with the DNA. This suggests a limited dynamic of these loops that are not engaged in crystal contacts (Figure 3a). In recent crystal and cryoEM structures of the Ku-DNA complex bound with DNA-PKcs or peptides [10,11,26], these loops also adopt the same conformation, suggesting a stable conformation of these loops with or without DNA. It remains to be documented if some rearrangements can happen during Ku translocation or if Ku threads in a helical manner like a screw on a nut.

Mutations of the Ku ring have been characterized with the aim to disrupt the interaction between Ku and DNA. Britton et al. thus designed a mutant called Mut6E with six residues of the Ku70 ring mutated in Glu to induce a repulsive charge (K282E K287E T300E K331E K338E R403E) [52] (Figure 3c). The Mut6E mutant does not bind DNA anymore and was used as a control in a super-resolution analysis of Ku localization.




Figure 3. Complementarity between Ku inner ring and the DNA (a) Structure of an inner ring loop of Ku (Ku80 Tyr397-Arg400) that interacts with the minor groove of the DNA. The figure shows a superimposition of this loop in the Ku/DNA complex and Ku free structures. (b) Surface representation of the Ku inner ring. Ku residues in contact with the DNA grooves are colored in blue. Due to these contacts, the radius of the inner ring of Ku is smaller than the DNA radius. (c) Surface representation of Ku. Residues in red are the six Ku70 positions mutated in Glu to disrupt the Ku-DNA interaction [52] (positions Lys282, Lys287, Thr300, Lys331, Lys338, and Arg403). Residues in magenta are the five positions mutated in Asp in [53] to mimic potential phosphorylation sites on the ring (Thr305, Ser306, Thr307, Ser314, and Thr316). Residues in purple are positions of Ku70 lysines (Lys31, Lys160, and Lys164) proposed to be involved in the Ku dRP/Lyase activity [43] (Lys31 is not observed, the first amino acid of Ku70 visible in the crystal is Arg35). Residues in light blue are a Cys of Ku80 that is a potential site of alkylation [54]. (d) CryoEM structure of (Ku70/80)/DNA-PKcs/DNA complex (PDB 6zha) [11]. DNA-PKcs is represented in the green cartoon with a semi-transparent surface. Ku70 and Ku80 are, respectively, shown in light and dark blue, while DNA is shown in purple. (e) Superimposition of Ku from the Ku/DNA-PKcs/DNA complex and Ku from the Ku/DNA complex (PDB 1jey) [45]. The loops of Ku in contact with DNA superimpose well, indicating that Ku has the same inner ring conformation with or without DNA-PKcs. (f) Inhibitors of Ku-DNA interactions reported by [55] (compound L) and by [56] (compound 245).

8. Insights from Recent 3D Structures of Ku-DNA Complexes Bound with C-NHEJ Factors

Recently, crystal structures of the Ku70/80/DNA complex were reported bound with peptides derived from Ku-binding motifs (KBM) of the nuclease APLF and of the ligation factor XLF [12,26]. The binding site of the KBM of APLF (A-KBM) is located at the periphery of the vWA (von Willebrand factor type A) domain of Ku80 in a position remote from the DNA binding site (Figure 2b). The amino acids involved in the interaction with

DNA superimpose well in the Ku-DNA-(A-KBM) and in the Ku-DNA structures [26]. In agreement with this observation, the affinity of Ku for dsDNA is the same with or without an A-KBM peptide. The KBM of XLF (X-KBM) is positioned in a remote site that is created after a large opening of the vWA domain that creates a binding pocket for this motif. The X-KBM is located closer to the DNA than the A-KBM is, but its binding has no impact on the affinity of Ku for a dsDNA. It is unknown if the other regions of the XLF and APLF can bind on Ku or DNA and if they have an impact on the Ku affinity for DNA or on its ability to translocate on DNA.

Several crystals and cryoEM structures of DNA-PKcs free and in a complex with Ku were reported [10,11,57–60]. Two recent cryoEM structures reported at a higher resolution used Y-shaped DNA with a 35 bp duplex region [10,11] (Figure 3d). In these structures, Ku interacts with DNA in a similar manner than in the crystal structures. The authors observed that the Ku70 and Ku80 subunits superimpose well with their counterparts in the absence of DNA-PKcs, except a slight movement of the α/β domain of Ku70 (Figure 3d,e). What is noteworthy is that the interactions of Ku with DNA-PKcs and the KBM peptides do not impact the position of the Ku loops that define the inner ring of the heterodimer. This observation favors the hypothesis of a conserved preformed state of the inner ring structure of Ku in different contexts.

9. Ku Can Recognize RNA Hairpins and RNA-DNA Hybrids

Ku was shown to bind about 200–500-fold less efficiently DNA–RNA hybrids such as poly(A)-poly(dT) or yeast tRNA, suggesting a marked specificity for dsDNA [20]. Another study by Kaczmarek et al. showed that Ku binds HIV-1 TAR RNA that forms a stable 50 nt hairpin structure with a 10-fold weaker affinity than dsDNA [61]. Using the SELEX approach (Systematic Evolution of Ligands by Exponential Enrichment), a series of RNA aptamers was characterized by Yoo et al. with tight affinities ($K_d \leq 2$ nM), similar to the K_d observed with dsDNA [62]. Another study by Anisenko et al. suggests that Ku70 without Ku80 can bind dsDNA and hairpin RNA with good affinities (K_d of 60 nM for dsDNA and 40 nM for hairpin RNA) [63]. A crystal structure of a complex, the Integrator complex with subunits 13 and 14 (INTS13/INTS14), with a structure reminiscent of the Ku heterodimer, was recently reported [64]. This complex processes 3'-ends of the spliceosome small nuclear RNAs (snRNAs). It presents structural similarities with Ku70/80, despite its low sequence similarities (identity 7–10%, similarity 17–19%). Ku70/80 and INTS13/INTS14 share highly similar domain architectures with N-terminal vWA domains, central β -barrels, and C-terminal α -helical domains. The INTS complex can bind single- and double-stranded DNA and RNA.

10. Ku Contributes to Synapse DNA Ends in a Complex with Other C-NHEJ Factors

An important issue in the c-NHEJ pathway is to maintain the DNA ends in close proximity and form a synapse between them. This is central to limit the ligation of one DNA extremity with the DNA end of another DSB and, thus, prevent chromosome translocation. Ku has a central role in the synapse formation all along the c-NHEJ pathway. It coordinates the alternative recruitment of the different c-NHEJ enzymatic activities on the DSB ends and passes on the baton from one to another without leaving DNA damage. This function is reminiscent of the role of factors of other DNA repair pathways like XRCC1 in BER (base excision repair) or MutS in MMR (mismatch repair) that stay at the site of damage, recruit downstream factors, and limit the release of DNA until the repair is complete [65]. In the first steps of c-NHEJ, the main synaptic activity is attributed to DNA-PKcs in a complex with Ku [66]. However, Zhao et al. showed with single-molecule FRET (smFRET) that both Ku and XRCC4:DNA ligase IV are necessary and sufficient to achieve a flexible synapsis of blunt DNA ends, whereas either alone is not [67]. Indeed, by itself, Ku has a weak synaptic activity between two DNA ends. Ramsden et al. also observed with an elegant assay using a biotin-DNA and a ^{32}P -labelled-DNA that in the presence of Ku, a fraction (8%) of labelled-DNA is captured on streptavidin beads [68]. Similarly, Wang

et al. showed with magnetic tweezers experiments that Ku by itself cannot form a synapse between two DNA ends [69]. They observed a transient synapse in the presence of Ku and DNA-PKcs with a half-life of 0.1 s. The addition of PAXX and Lig4/XRCC4/XLF reinforces the stability of the synapse up to 66 s in this system. In *S. cerevisiae*, no DNA-PKcs has been identified so far [70]. The MRX complex (Mre11/Rad50/Xrs2) has an important role in c-NHEJ and it is proposed to contribute to synapse formation as observed by AFM (atomic force microscopy) [71,72].

11. Number of Ku Molecules at the Ends: In Vitro versus Cellular Analyses

In vitro studies showed that several Ku molecules can thread on dsDNA with up to one Ku loaded every 25–30 bp. The first Ku molecules bind close to DNA ends and then translocate inward so that other Ku molecules can have access to the ends. An EMSA performed with Ku thus presented regular ladder profiles where the different bands correspond to an increasing number of Ku molecules on DNA. Biophysical measurements with calorimetry [26], anisotropy fluorescence [73] and SEC-MALS (size-exclusion chromatography with multiangle light scattering), SPR, or BLI (bio-layer interferometry) (unpublished data) are in agreement with this mechanism. The molecular mechanism of the translocation of Ku is not coupled to ATP hydrolysis as for translocases and helicases. From the structural studies, we can suggest two hypotheses. As described above, the inner part of the Ku ring makes tight contacts with the minor and major grooves of DNA. One possibility is that the ring presents some conformational changes on the loops in contact with the DNA and somehow enlarge to allow a passive one-dimensional diffusion of Ku on the DNA. A second hypothesis is that the inner ring has a rigid and stable conformation so that Ku translocate with a helical translation (Figure 3b).

The translocation of Ku on DNA has been proposed to be more limited in a cellular context by using high-resolution microscopy. A seminal study used ribonuclease- and detergent-based pre-extraction treatment to improve the visualization of Ku at DSB ends [52]. Ribonuclease treatment favors the release of a large proportion of Ku molecules bound to nuclear structures through interactions with RNA. This method combined with STED (stimulated-emission-depletion) microscopy allowed the visualization of Ku foci of small spherical shape (size of about 80 nm), corresponding to a maximal number of three to four molecules at each DNA end. Analyses by bleaching steps of fluorescent Ku molecules suggest that, predominantly, only one Ku molecule may be present on each side of the DSB [52]. These results indicate that despite a high concentration of Ku in the cell, the threading of Ku is reduced in the cell environment by some factors or processes.

12. Factors That May Limit Ku Threading in Cell

One hypothesis regarding the limited threading of Ku in the cell is the presence of nucleosomes at DSB ends that may block the Ku diffusion. In vitro studies with EMSA showed that Ku efficiently binds DNA ends close to the nucleosome, though with reduced apparent affinity (apparent K_d of 0.3 and 6 nM on free dsDNA and DNA ends, respectively, close to the nucleosome) [74]. In this study, the authors showed that Ku is able to displace the protein linker histone H1 from DNA, but it does not displace the nucleosome. They proposed that Ku “peels” some 50 bp DNA from the nucleosome [31,74]. The dynamic of nucleosomes following the formation of a DSB was analyzed using the I-PpoI homing endonuclease and the ChIP (chromatin immunoprecipitation) analysis of histone H3 around the I-PpoI break sites [75]. The authors observed a decrease in H3 over 1000 bp, corresponding to about eight nucleosomes (146 bp of DNA are wrapped around an octamer of histone H2A, H2B, H3, and H4). The disassembly is promoted in an ATM- and ATR-dependent manner by the nucleosome remodeler INO80 [75] (Figure 4). However, the extent of this nucleosome clearing would allow more Ku sliding than observed in the cells [52]. A second hypothesis is that other DNA binding proteins may limit Ku diffusion inward through interactions with a proximal Ku molecule and adjacent DNA. c-NHEJ factors interacting

with Ku with DNA binding properties may play such a role. These include DNA-PKcs, XLF, APLF, WRN, Lig4/X4, PAXX, and likely MRN (Mre11/Rad50/Nbs1) [12,13,76–78].

13. The Race for the DSB Ends

The two main DSB repair pathways are homologous recombination (HR) and non-homologous end-joining (NHEJ). HR requires a homologous template to guide the repair of the broken DNA and, therefore, functions in S and G2 phases due to the presence of a sister chromatid, whereas c-NHEJ is active during all cell cycles out of mitosis and mediates the direct re-ligation of the broken DNA. The complexity of the broken ends can determine the choice of the repair pathway [5]. Simple DSB ends are mainly repaired by c-NHEJ, and complex DSBs often require end resections. c-NHEJ requires limited sequence homologies (0–4 bp) between the overhangs of the DNA ends, while HR requires extensive end resection with a typically long homology tract (more than 100 bp). Several factors influence the choice of the pathway including the cell cycle stage, the DNA end resection, and specific post-translation modifications. At least four sensors detect DSB: Ku, PARP, MRN, and RPA after DSB processing. Using a pulsed near-infrared (NIR) laser, Mari et al. showed that Ku starts localizing at DSB within a few seconds and reaches a maximum after 3 min, and that 20% remains after 2 h [79]. Another laser-induced study showed that not only Ku but also MRE11 and PARP-1 are recruited immediately and independently to damage sites [80]. These factors are still detectable on DSB for about 2 h for PARP1, 4–6 h for Ku, and more than 8 h for MRE11. A meta-analysis of recruitment data of over about 80 DSB-related proteins was reported [81] (<http://www.dna-repair.live/cumulus/>, accessed on 17 March 2021). From this study, c-NHEJ factors are among the first to arrive at the DSB sites followed by the slower HR machinery. The median time for access on the DSB is about 10 s for Ku. It is among the fastest with c-NHEJ factors like Artemis, XLF, and PNKP (polynucleotide kinase 3'-phosphatase). PARP-1 (Poly (ADP-ribose) polymerase-1) belongs to the rare factors that are reported to be faster (about 3 s). This is in agreement with previous studies that report a fast access of PARP-1 at DSB [82]. In yeast, different studies showed that Ku and MRX/MRN are among the first proteins recruited to DSB [72,83–85].

Besides the competition to be first place, sensors can be actively removed. In budding yeast, meiotic DSB processing gave some clues to the removal of Ku by the combined activity of the Mre11-Rad50-Nbs1 complex (MRN) and CtIP (Ctp1/Sae2) required to initiate DSB resection [86,87]. It was shown that Sae2 (CtIP in humans) activates Mre11 endonuclease activity at a distance from the Ku-DNA ends and that this nick serves as an entry point for exonucleases [86]. This mechanism was confirmed by biochemical and single-molecules analyses [88–90], showing that Ku can be removed by the activity of these combined nucleases. It is well established that Ku prevents the exonucleolytic attack of free DNA ends [84,91–93]. In cells, Ku is also a major inhibitor of an alternative end-joining pathway involving an MRN-dependent ends resection step [94].

14. Ku Molecules Trapped after Ligation Are Actively Removed

The Ku ring has no clasp, and it is unlikely to open to dissociate. The crystal structure shows a large intertwining of both subunits (Figure 2b). An opening would require an unwinding of the C-terminal arms of Ku70 and Ku80 (constituting 95 and 113 amino acids, respectively). The ring structure of Ku is different from other ring shape factors with DNA binding properties like the PCNA trimer (proliferating cell nuclear antigen). The PCNA is a trimeric protein involved in the processivity of the replicative DNA polymerase. It can assemble and disassemble on dsDNA independently of the presence of DNA ends with the help of the replication factor C [95]. Paillard et al. showed in vitro that Ku is stable when trapped after ligation on mini-circles DNA. It remains bound in the presence of an increasing amount of NaCl from 0.1 to 2 M under conditions where it dissociates from linear DNA [31].

Different studies recently reported that Ku trapped after ligation would be removed after ubiquitination. Postow et al. showed that Ku80 is degraded in response to DSB in

a ubiquitin-dependent manner, but independent of the proteasome [96]. The extraction of Ku from repaired DSB can take place after ubiquitination mediated by three different E3 ligases (SCFFbx12, RNF8, or RNF138) [96–100]. After ubiquitination, Ku is removed by an AAA+ ATPase called Valosin-containing protein VCP/p97 with its co-factors Udf1-Npl4 [101]. This hexameric AAA+-type ATPase targets ubiquitinated proteins with the help of ubiquitin adaptor proteins. These ATPase use the energy of ATP hydrolysis to segregate their substrate proteins away from their cellular partners or structures (Figure 4e). In U2OS cells, Ku80 foci formed after ionizing radiation are largely resolved within 60 min, whereas in the presence of p97 inhibitor, the Ku80 foci persist later [101]. Recently, Ku70 was shown to be a direct target for the deubiquitinase USP14, leading to a reduced recruitment to DSB sites [102]. The conjugation of the ubiquitin-like protein NEDD8 to target proteins at breaks sites has been shown to be required for Ku ubiquitination and Ku and associated repair factors released from these sites following repair [103].




Figure 4. Association and removal of Ku in a chromatin context. (a) Upon DSB formation, Ku recognizes DNA ends. Super-resolution microscopy studies suggest that only one Ku may bind at the DSB and that no threading inward is observed with the loading of additional Ku molecules [52]. (b) Upon DSB formation, some nucleosomes are disassembled by the nucleosome remodeler INO80 [75]. Some factors not yet identified likely limit the Ku threading inward (light blue protein). c-NHEJ factors like XLF, APLF, or PAXX may be involved in this role. (c) Ku recruits DNA-PKcs, and the complex forms a synapse between the two DNA ends in a conformation that may correspond to the dimer observed by Chaplin et al. [11]. (d) Ku can recruit alternatively nucleases, polymerases, and the ligation complex at the DSB ends (here, the Ligase 4 is represented in a complex with an XLF-XRCC4 filament [104,105]). (e) Once the ligation is completed, Ku is trapped on the DNA. The AAA+ ATPase VCP/p97 will remove in place the trapped Ku molecules [101].

15. Ku at Single-Ended DSB and at Stalled Replication Fork

A frequent source of DSB in cells comes from the encounter of the replisome with a discontinuity in the DNA template that generates single-ended double-strand breaks (seDSB) [106] (Figure 1i). With the lack of a second DNA end, the damage cannot be repaired by c-NHEJ and is, therefore, processed by HR in the S/G2 phase of the cell cycle. Ku can bind at seDSB and, thus, needs to be removed to allow resection so that HR can proceed. MRN and EXO1 carry out an incision and excision upstream of the seDSB and remove Ku at the seDSB generated by CPT during replication [107,108]. Balestrini et al. performed a genetic screen in yeast to isolate yKU70 alleles that phenocopy the Ku deficiency with respect to single-ended DSB, but remain proficient for c-NHEJ [109]. They isolated six single mutants that present a reduced affinity for DNA ends and an enhanced capacity to translocate on DNA, as observed by AFM. MRN-Ctp1 also promotes Ku removal after an initial resection in the context of terminally arrested forks. Recent studies in fission yeast show that the presence of Ku at single DNA ends regulates the extension of resection, the level of recruitment of RPA and Rad51, and the efficiency of the fork-restart process (Figure 1h) [110].

16. Ku at Telomeres in Yeast

Telomeres have evolved to protect chromosome termini against degradation, checkpoint activation, and illicit repair [111]. In addition to telomere proteins, Ku70/80 is recruited constitutively at telomeres in budding yeast *S. cerevisiae* [112,113]. Ku directly binds the telomere dsDNA end [114] and protects telomeres against 5' resection [115–118]. To prevent Ku-dependent telomere fusions, the Ku function in NHEJ is blocked at telomeres by telomere factors (namely Rap1, Rif2, and Sir4) (Marcand et al., 2008, Lescasse et al., 2013, Roisé-Hamelin et al., 2021). Ku also participates in telomerase recruitment through its interaction with the TLC1 RNA subunit of telomerase and the telomere factor Sir4 [112,113,119–121]. The TLC1 RNA region that interacts with Ku is called the KBS for the Ku binding site. The TLC1-KBS specifically encompasses nucleotides 288 to 312 and forms a high-affinity interaction with Ku70/80 (Kd = 75 nM). The crystal structure of TLC1-KBS in the complex with full-length Ku70/80 was recently reported [122]. It reveals that TLC1 adopts a bulged stem-loop structure, and a bent conformation between the P1 and P2 helices of TLC1-KBS upon interaction with Ku [122] (Figure 5).

A direct interaction between the silencing factor Sir4 N-terminal KBM motif (100–115) and Ku80 recruits the Ku-telomerase complex to telomeres (Bertuch and Lundblad 2003 MCB) (Chen et al., 2018). Sir4 KBM forms a short alpha-helix that binds a hydrophobic groove inside the Ku80 vWA domain [122]. The authors proposed a model where, in the first step, Ku would interact with Sir4 at the telomeres and favor the telomerase recruitment in an inactive manner throughout the cell cycle. In the second step, the telomerase would interact with the ssDNA-bound cdc13 protein, in an active manner, a form competent for telomere elongation [122]. Thus, TLC1 and DNA binding to Ku is mutually exclusive.

As Sir4 is a core component of yeast heterochromatin (Faure et al. 2019 GBE), the KU-Sir4 interaction also favors heterochromatin interactions at telomeres (Ribes-Zamora et al. 2007 NSMB). This pathway is proposed to require a direct loading of KU to the chromosome DNA end (Lopez et al., 2011). Finally, Ku anchors telomeres at the nuclear periphery through Sir4-dependent and -independent pathways [119,120,123–125].




Figure 5. Ku contributes to the recruitment of the telomerase. (a,b) Crystal structure of *S. cerevisiae* Ku70/80 with the Ku binding site (KBS) of the TLC1 RNA of the yeast telomerase [122]. The KBS hairpin RNA is positioned in the ring channel of Ku in agreement with the competition observed between RNA and dsDNA binding. (c) Ku is proposed to be recruited at telomeres through its interaction between the Ku80 α/β domain and Sir4 KBM. Ku will contribute to the telomerase recruitment through its interaction with the KBS of TLC1. The interactions of Sir3 and TLC1 with the Mps3 factor at the nuclear envelope is indicated. (d) In a second step, the telomerase would interact with Cdc13 to adopt its active state conformation. It is not known whether Ku keeps interacting with TLC1 at this stage (indicated by a "?").

17. Ku at Telomeres in Mammalian Cells

The organization and functions associated with mammalian telomeres are different from budding yeast (no telomeric silencing or nuclear anchoring), but the Ku heterodimer is also present at the mammalian telomeres [126]. In human cells, Ku is essential to telomere stability and, therefore, cell viability [127]. It has been observed that Ku localizes to telomeres through its high affinity for the TRF1 shelterin protein ($K_d = 0.4$ nM) [128]. Ku can also bind with high affinities to DNA duplex telomeric sequences with TTAGGG repeats [129,130]. The presence of the G-quartet sequence in 3' and 5' of DNA does not prevent Ku from binding with high affinity. It has been shown that the paradoxical presence of the main end-joining initiator Ku at telomeres ensures protection against aberrant joining by an alternative pathway [131]. The use of knockdown mice for all shelterin

proteins also highlighted that the presence of Ku at mammalian telomeres is associated with the protection against alternative nonhomologous end-joining, and homologous-directed repair [132]. It was shown that Ku interacts with the RNA component of the human telomerase (hTR) *in vitro* and in a cellular context [133]. The interaction is mediated by 47 nt in 3' of hTR in a manner reminiscent of the interaction between yeast Ku and the stem loop of TLC1, though there is no sequence similarity between hTR and TLC1.

18. Ku Would also Be Able to Bind Internal DNA Sequences

Ku was also proposed to bind at DNA in the absence of DNA ends. Giffin et al. used micro-circles DNA containing a NRE1 (negative regulatory element 1) sequence, a 14 bp DNA sequence element. NRE1 is in the long terminal repeat (LTR) of mouse mammary tumor virus (MMTV) that is important for repressing inappropriate viral expression. Ku presents a strong affinity for the micro-circles that contain this NRE1 motif [134]. The micro-circles are resistant to several nucleases to exclude the possibility that Ku binds to nicks, to ends, or to structural features in the DNA. The molecular basis of the interaction of Ku on internal DNA motifs with a sequence specificity is not well understood. It is proposed that the C-terminal SAP domain of Ku70 may contribute with some DNA binding properties observed for this domain, though only a moderate affinity has been reported for this SAP domain [48,135].

19. Post-Translational Modifications of Ku Regulate Its Activities

Ku bears several post-translational modifications (PTM) that can potentially regulate its DNA recognition activities. Lees-Miller's laboratory showed that Ku is phosphorylated by DNA-PKcs *in vitro* at positions Ser6 of Ku70 and Ser 577, Ser 580, and Thr 715 of Ku80, though none of these positions are the SQ consensus motif for DNA-PKcs [4,136]. A form of Ku with mutations at these five positions in alanine proved to be fully able to complement the radiation sensitivity, suggesting that the DNA-PK-dependent phosphorylation of Ku70/80 is not required for c-NHEJ [137]. Another study proposed a role for the phosphorylation of Ku on its dissociation from DSBs in a cell [53]. Using laser-generated DNA damages and live cell microscopy, an increased retention of Ku at DSBs was observed in the presence of a broad-spectrum PI3K kinase inhibitor (wortmannin). A knowledge-based approach was used to design a Ku70 mutant with Asp mutations on five potential phosphorylation sites (mutant 5D). These positions are located in the junction of the pillar and bridge regions of Ku70 (Figure 3c). This mutant has a shorter time of retention than the wild-type Ku on a DSB in cells and a weaker DNA binding affinity. A second mutant with the five positions mutated in Ala (mutant 5A), resistant to phosphorylation, has a longer retention time than the wild-type Ku. Analyses of post-translational modifications (PTMs) on the Ku heterodimer by high-resolution 2D-gel electrophoresis and the mass spectrometry of chronic lymphocytic leukemia (CLL) proteins allowed the identification of the phosphorylation of Ser27 and Ser33 of Ku70 in the resistant form of CLL [138]. These positions are located in the N-terminal region of Ku70 that is predicted as disordered and not visible in the crystal. These phosphorylations promote fast but unfaithful DNA repair in cancer cell lines. Sumoylation has been characterized in the C-terminal region of budding yeast Ku70 and shown to strengthen the association with DNA [139–141]. Site-specific mapping of the human SUMO proteome identified sumoylation on both Ku70 and Ku80 subunits [142].

Other PTMs of Ku have been reported. In the presence of DNA and NAD⁺, PARP-1 can poly(ADP-ribose)ate itself and Ku70/80. EMSA showed that the poly-(ADP-ribose)ation of Ku70/80 decreases the DNA-binding affinity of Ku [143]. In line with this observation, Hohegger et al. reported that Parp-1 inhibits Ku from binding to DSB, thereby allowing access of the HR machinery [144]. The acetylation of Ku has been shown to regulate the interaction of Ku with Bax, which suppress apoptosis by sequestering Bax from mitochondria [145,146]. It is suspected that it is the less abundant cytoplasmic pool of Ku that is responsible for Bax sequestration. This role of Ku is mediated by acetylation on eight

lysines of Ku70 by either the CREB-binding protein (CBP) or P300/CBP-associated factor pCAF: Five in its C-terminus (Lys539, Lys542, Lys544, Lys553, and Lys) and three in its N-terminal DNA binding domain (Lys317, Lys331, and Lys338). The acetylation of Lys539 and Lys542 has been found to be critical for the regulation of Bax-mediated apoptosis [147]. Ku acetylation also disrupts Ku–p53 mRNA interactions, leading to a relief of Ku-mediated translational repression of the p53 mRNA after genotoxic stress [148].

20. Regulation of Ku Activities by Small Molecules

The interaction between Ku and DNA is dependent on reduced cysteines, and alkylation with *N*-ethylmaleimide prevents the interaction with DNA [54]. Similarly, the diamide treatment of Ku disrupts the Ku–DNA binding interaction, and this was abrogated by dithiothreitol treatment, demonstrating a reversible redox effect [149]. Hydroxyethyl disulfide also causes an enhanced radiation sensitivity and a direct inhibition of DNA end-binding activity of the Ku heterodimer by oxidation of its cysteine residues [150]. The analysis of the crystal structure showed that a Cys residue of Ku80 is located in the ring and could contribute to the impact of an alkylating agent (Figure 3c). An endogenous metabolite, inositol-6-phosphate (IP6), was shown to bind Ku heterodimer and stimulate the c-NHEJ pathway [151–153]. Molecular modeling and site-directed mutagenesis suggest an interaction site for this highly negatively charged molecule on a basic pocket of Ku at the interface between the two subunits [154]. Finally, some studies characterized interesting Ku inhibitors. Weterings et al. identified several potential inhibitors by computational screening and characterized a low micro-molar inhibitor that disrupts the binding of Ku70/80 to DNA substrates [55]. Recently, Gavande et al. described the discovery of highly potent and selective DNA-PK inhibitors that block the Ku70/80 heterodimer interaction with DNA [56] (Figure 3f).

21. A Well-Conserved Gene along Evolution

DSB repair factors equivalent to Ku70/80 have been identified and characterized in several bacteria [155,156], fungi [112,157–159], and bacteriophages [155,156,160]. Some NHEJ-like genes have also been identified in archaea, and a complete c-NHEJ system, comprising Lig, Pol, PE, and Ku, was characterized in the archaeon *Methanocella paludicola* (Mpa) [161]. Nenarokova reported frequent losses of Ku70 and Ku80 genes in several lineages of parasitic protists like human parasitic protists *Trypanosoma* spp., *Plasmodium* spp., and *Encephalitozoon cuniculi*. Multiple sequence alignments show that the core region of Ku is well conserved in most species. The similarity between the eukaryotic and bacterial Ku proteins indicates a likely common ancestral gene. The C-terminal region of Ku80 that is involved in the interaction with DNA-PKcs is conserved mainly in multicellular organisms that have a DNA-PKcs gene. It is absent, for example, in *S. cerevisiae* that has no identified DNA-PKcs.

In silico analyses of microbial genomes identified putative Ku genes in several prokaryote genomes [155,162] (Aravind and Koonin, 2001; Doherty et al., 2001). When bacterial NHEJ was studied in prokaryotes, a Ku homodimer was found to bind DNA ends and to recruit LigD, which carries nuclease, polymerase, and ligase activities [163]. Intriguingly, some species comprise up to four orthologues of Ku and Lig, which carry out different NHEJs under specific stress conditions [164]. The bacterial Ku homologs (~30–40 kDa) are smaller than their eukaryote counterparts. They comprise the central core of Ku70 and Ku80, and they lack the α/β N-terminal region and the C-terminal region of both subunits. No crystal structure of bacterial Ku has been available up until now but multiple sequence alignments suggest that the bacterial Ku core region shares the same fold as that of eukaryotic Ku. The mycobacterial Ku forms homodimers, binds DNA ends, and can thread on DNA, as observed in eukaryotes [165]. The c-NHEJ in many prokaryotes is performed by a two-component system constituted by the Ku homodimer and a multifunctional DNA ligase (LigD). The interaction between Ku and LigD involves the C-terminal region of Ku. Additional studies on Ku from *Bacillus subtilis* have revealed that the C-terminal region of

bacterial Ku is important for Ku threading on DNA and its synapse properties [166]. Using single-molecule DNA analysis, it was shown that the *Bacillus subtilis* Ku homodimer alone, contrary to human Ku, forms a ~2 s synapsis between blunt DNA ends that is increased to ~18 s upon addition of LigD. These synaptic properties are dependent on the C-terminal arms of Ku [167]. The bacterial Ku was also used as a tool to show that blocking DNA ends affects end resection and HR in mammalian cells [168].

Original properties of Ku were reported in ciliates like *Paramecium tetraurelia*. These organisms have massively programmed genome rearrangements (PGR) during the development of their somatic nucleus with the precise elimination of at least 45,000 germline sequences (internal eliminated sequences, IESs). IES excision proceeds through a cut-and-close mechanism mediated by a domesticated transposase, PiggyMac, essential for DNA cleavage, and the c-NHEJ pathway. The genome of *Paramecium tetraurelia* encodes for two Ku70 and three Ku80 paralogs. The Ku70a and Ku80c paralogs are produced during sexual processes. The Ku70a/80c heterodimer co-purifies with PiggyMac and is essential for the licensing of its endonuclease [169]. Recently, immunofluorescence microscopy and high-throughput DNA sequencing revealed that Ku80c stably anchors the PiggyMac (Pgm) endonuclease through an interaction on the α/β domain of Ku80c [170]. The ciliate model suggests a tight coupling between DSB introduction and repair with a central role of Ku [171].

22. Conclusions

In conclusion, several important issues regarding the molecular mechanism of Ku are still unresolved. It is still unclear to what extent Ku is sensitive to the diversity of structures and chemistry of the DNA ends that are generated during accidental or programmed DSB; The prominent role of Ku at telomeres are not clearly understood. It is not clear how Ku interacts with the other initial responders to DSB including MRN (Mre11-Rad50-Nbs1 complex) and PARP enzymes (Poly [ADP-ribose] polymerases). And additional studies are needed to characterize the functions of Ku that are conserved in the different organisms and those that are unique to some organisms.

Ku70/80 recognizes a large variety of DSBs with different chemistries and structures of the DNA ends. The heterodimer is one of the first players to be on the DSB site together with MRN and PARP-1, and it contributes in humans to orient DSB repair predominantly toward the c-NHEJ pathway. Despite important data on the molecular mechanism of Ku in vitro and in cells, additional studies are needed to further document how it interacts with various DNA and RNA ends and with DNA/RNA hybrids. It will be central in the future to understand how Ku interactions with its partners regulate its threading capacity in cells during the recognition of the ends and during the downstream steps of the c-NHEJ. The role of Ku beyond the c-NHEJ pathway in telomere maintenance, replication fork arrest/restart, and in the cytoplasm will also require further analyses. Finally, Ku is an important pharmaceutical target to modulate the c-NHEJ pathway [172]. The design of potent inhibitors of Ku-DNA or Ku-protein interactions will be central to characterize new molecules that may serve as a radiosensitizer or to improve genome editing.

Author Contributions: Writing—original draft preparation, S.Z., M.S.E.D., F.I., P.F.-V., M.-H.L.D., V.R., J.B.C.; writing—review and editing, S.Z., M.S.E.D., F.I., P.F.-V., M.-H.L.D., V.R., J.B.C.; supervision, J.B.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by INCA 2016-PLBIO-159, ANR-17-CE12-0015, ANR-17-CE12-0020, ANR-594 18-CE44-0008, ANR-20-CE11-0026 and from FRISBI ANR-10-INBS-0005.

Acknowledgments: We would like to thank Patrick Calsou (IPBS, Toulouse), Stephane Marcand and Florian Roisné-Hamelin (Institut Jacob, iRCM, Fontenay aux rose), and Mauro Modesti (CRCM, Marseille) for their fruitful discussions and critical reading of the manuscript. JBC would like to thank the support from INCA 2016-PLBIO-159, ANR-17-CE12-0015, ANR-17-CE12-0020, ANR-18-CE44-0008, ANR-20-CE11-0026, and FRISBI ANR-10-INBS-0005.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Mimori, T.; Akizuki, M.; Yamagata, H.; Inada, S.; Yoshida, S.; Homma, M. Characterization of a high molecular weight acidic nuclear protein recognized by autoantibodies in sera from patients with polymyositis-scleroderma overlap. *J. Clin. Investig.* **1981**, *68*, 611–620. [[CrossRef](#)] [[PubMed](#)]
2. Gell, D.; Jackson, S.P. Mapping of protein-protein interactions within the DNA-dependent protein kinase complex. *Nucleic Acids Res.* **1999**, *27*, 3494–3502. [[CrossRef](#)] [[PubMed](#)]
3. Woods, D.S.; Sears, C.R.; Turchi, J.J. Recognition of DNA Termini by the C-Terminal Region of the Ku80 and the DNA-Dependent Protein Kinase Catalytic Subunit. *PLoS ONE* **2015**, *10*, e0127321. [[CrossRef](#)] [[PubMed](#)]
4. Jette, N.; Lees-Miller, S.P. The DNA-dependent protein kinase: A multifunctional protein kinase with roles in DNA double strand break repair and mitosis. *Prog Biophys Mol. Biol.* **2015**, *117*, 194–205. [[CrossRef](#)]
5. Zhao, B.; Rothenberg, E.; Ramsden, D.A.; Lieber, M.R. The molecular basis and disease relevance of non-homologous DNA end joining. *Nat. Rev. Mol. Cell Biol.* **2020**, *21*, 765–781. [[CrossRef](#)]
6. Lieber, M.R.; Lu, H.; Gu, J.; Schwarz, K. Flexibility in the order of action and in the enzymology of the nuclease, polymerases, and ligase of vertebrate non-homologous DNA end joining: Relevance to cancer, aging, and the immune system. *Cell Res.* **2008**, *18*, 125–133. [[CrossRef](#)]
7. Smider, V.; Rathmell, W.K.; Lieber, M.R.; Chu, G. Restoration of X-ray resistance and V(D)J recombination in mutant cells by Ku cDNA. *Science* **1994**, *266*, 288–291. [[CrossRef](#)]
8. Rathmell, W.K.; Chu, G. Involvement of the Ku autoantigen in the cellular response to DNA double-strand breaks. *Proc. Natl Acad. Sci. USA* **1994**, *91*, 7623–7627. [[CrossRef](#)]
9. Taccioli, G.E.; Gottlieb, T.M.; Blunt, T.; Priestley, A.; Demengeot, J.; Mizuta, R.; Lehmann, A.R.; Alt, F.W.; Jackson, S.P.; Jeggo, P.A. Ku80: Product of the XRCC5 gene and its role in DNA repair and V(D)J recombination. *Science* **1994**, *265*, 1442–1445. [[CrossRef](#)]
10. Yin, X.; Liu, M.; Tian, Y.; Wang, J.; Xu, Y. Cryo-EM structure of human DNA-PK holoenzyme. *Cell Res.* **2017**, *27*, 1341–1350. [[CrossRef](#)]
11. Chaplin, A.K.; Hardwick, S.W.; Liang, S.; Kefala Stavridi, A.; Hnizda, A.; Cooper, L.R.; De Oliveira, T.M.; Chirgadze, D.Y.; Blundell, T.L. Dimers of DNA-PK create a stage for DNA double-strand break repair. *Nat. Struct. Mol. Biol.* **2020**. [[CrossRef](#)]
12. Frit, P.; Ropars, V.; Modesti, M.; Charbonnier, J.B.; Calsou, P. Plugged into the Ku-DNA hub: The NHEJ network. *Prog. Biophys. Mol. Biol.* **2019**. [[CrossRef](#)]
13. Grundy, G.J.; Moulding, H.A.; Caldecott, K.W.; Rulten, S.L. One ring to bring them all—the role of Ku in mammalian non-homologous end joining. *DNA Repair* **2014**, *17*, 30–38. [[CrossRef](#)]
14. Gu, Y.; Jin, S.; Gao, Y.; Weaver, D.T.; Alt, F.W. Ku70-deficient embryonic stem cells have increased ionizing radiosensitivity, defective DNA end-binding activity, and inability to support V(D)J recombination. *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 8076–8081. [[CrossRef](#)]
15. Koike, M.; Ikuta, T.; Miyasaka, T.; Shiomi, T. Ku80 can translocate to the nucleus independent of the translocation of Ku70 using its own nuclear localization signal. *Oncogene* **1999**, *18*, 7495–7505. [[CrossRef](#)] [[PubMed](#)]
16. Koike, M.; Ikuta, T.; Miyasaka, T.; Shiomi, T. The nuclear localization signal of the human Ku70 is a variant bipartite type recognized by the two components of nuclear pore-targeting complex. *Exp. Cell Res.* **1999**, *250*, 401–413. [[CrossRef](#)]
17. Takeda, A.A.; de Barros, A.C.; Chang, C.W.; Kobe, B.; Fontes, M.R. Structural basis of importin- α -mediated nuclear transport for Ku70 and Ku80. *J. Mol. Biol.* **2011**, *412*, 226–234. [[CrossRef](#)]
18. Bertinato, J.; Schild-Poulter, C.; Hache, R.J. Nuclear localization of Ku antigen is promoted independently by basic motifs in the Ku70 and Ku80 subunits. *J. Cell Sci.* **2001**, *114*, 89–99.
19. Fell, V.L.; Schild-Poulter, C. The Ku heterodimer: Function in DNA repair and beyond. *Mutat. Res. Rev. Mutat. Res.* **2015**, *763*, 15–29. [[CrossRef](#)]
20. Mimori, T.; Hardin, J.A. Mechanism of interaction between Ku protein and DNA. *J. Biol. Chem.* **1986**, *261*, 10375–10379. [[CrossRef](#)]
21. Blier, P.R.; Griffith, A.J.; Craft, J.; Hardin, J.A. Binding of Ku protein to DNA. Measurement of affinity for ends and demonstration of binding to nicks. *J. Biol. Chem.* **1993**, *268*, 7594–7601. [[CrossRef](#)]
22. De Vries, E.; van Driel, W.; Bergsma, W.G.; Arnberg, A.C.; van der Vliet, P.C. HeLa nuclear protein recognizing DNA termini and translocating on DNA forming a regular DNA-multimeric protein complex. *J. Mol. Biol.* **1989**, *208*, 65–78. [[CrossRef](#)]
23. Falzon, M.; Fewell, J.W.; Kuff, E.L. EBP-80, a transcription factor closely resembling the human autoantigen Ku, recognizes single-to double-strand transitions in DNA. *J. Biol. Chem.* **1993**, *268*, 10546–10552. [[CrossRef](#)]
24. Arosio, D.; Cui, S.; Ortega, C.; Chovanec, M.; Di Marco, S.; Baldini, G.; Falaschi, A.; Vindigni, A. Studies on the mode of Ku interaction with DNA. *J. Biol. Chem.* **2002**, *277*, 9741–9748. [[CrossRef](#)]
25. Gontier, A.; Varela, P.F.; Nemoz, C.; Ropars, V.; Aumont-Nicaise, M.; Desmadril, M.; Charbonnier, J.B. Measurements of Protein-DNA Complexes Interactions by Isothermal Titration Calorimetry (ITC) and Microscale Thermophoresis (MST). *Methods Mol. Biol.* **2021**, *2247*, 125–143. [[CrossRef](#)]
26. Nemoz, C.; Ropars, V.; Frit, P.; Gontier, A.; Drevet, P.; Yu, J.; Guerois, R.; Pitois, A.; Comte, A.; Delteil, C.; et al. XLF and APLF bind Ku80 at two remote sites to ensure DNA repair by non-homologous end joining. *Nat. Struct. Mol. Biol.* **2018**, *25*, 971–980. [[CrossRef](#)]

27. Velours, C.; Aumont-Nicaise, M.; Uebel, S.; England, P.; Velazquez-Campoy, A.; Stroebel, D.; Bec, G.; Soule, P.; Quétard, C.; Ebel, C.; et al. Macromolecular interactions in vitro, comparing classical and novel approaches. *Eur. J. Biophys.* **2021**, in press. [[CrossRef](#)]
28. Tuteja, N.; Tuteja, R.; Ochem, A.; Taneja, P.; Huang, N.W.; Simoncsits, A.; Susic, S.; Rahman, K.; Marusic, L.; Chen, J.; et al. Human DNA helicase II: A novel DNA unwinding enzyme identified as the Ku autoantigen. *EMBO J.* **1994**, *13*, 4991–5001. [[CrossRef](#)]
29. Krasner, D.S.; Daley, J.M.; Sung, P.; Niu, H. Interplay between Ku and Replication Protein A in the Restriction of Exo1-mediated DNA Break End Resection. *J. Biol. Chem.* **2015**, *290*, 18806–18816. [[CrossRef](#)]
30. Yuan, Y.; Britton, S.; Delteil, C.; Coates, J.; Jackson, S.P.; Barboule, N.; Frit, P.; Calsou, P. Single-stranded DNA oligomers stimulate error-prone alternative repair of DNA double-strand breaks through hijacking Ku protein. *Nucleic Acids Res.* **2015**, *43*, 10264–10276. [[CrossRef](#)]
31. Paillard, S.; Strauss, F. Analysis of the mechanism of interaction of simian Ku protein with DNA. *Nucleic Acids Res.* **1991**, *19*, 5619–5624. [[CrossRef](#)]
32. Wang, J.; Satoh, M.; Chou, C.H.; Reeves, W.H. Similar DNA binding properties of free P70 (KU) subunit and P70/P80 heterodimer. *FEBS Lett.* **1994**, *351*, 219–224. [[CrossRef](#)]
33. Tadi, S.K.; Tellier-Lebegue, C.; Nemoz, C.; Drevet, P.; Audebert, S.; Roy, S.; Meek, K.; Charbonnier, J.B.; Modesti, M. PAXX Is an Accessory c-NHEJ Factor that Associates with Ku70 and Has Overlapping Functions with XLF. *Cell Rep.* **2016**, *17*, 541–555. [[CrossRef](#)]
34. Griffith, A.J.; Blier, P.R.; Mimori, T.; Hardin, J.A. Ku polypeptides synthesized in vitro assemble into complexes which recognize ends of double-stranded DNA. *J. Biol. Chem.* **1992**, *267*, 331–338. [[CrossRef](#)]
35. Ma, Y.; Lieber, M.R. DNA length-dependent cooperative interactions in the binding of Ku to DNA. *Biochemistry* **2001**, *40*, 9638–9646. [[CrossRef](#)]
36. Chen, X.; Tomkinson, A.E. Yeast Nej1 is a key participant in the initial end binding and final ligation steps of nonhomologous end joining. *J. Biol. Chem.* **2011**, *286*, 4931–4940. [[CrossRef](#)]
37. Inagawa, T.; Wennink, T.; Lebbink, J.H.G.; Keijzers, G.; Florea, B.I.; Verkaik, N.S.; van Gent, D.C. C-Terminal Extensions of Ku70 and Ku80 Differentially Influence DNA End Binding Properties. *Int. J. Mol. Sci.* **2020**, *21*. [[CrossRef](#)]
38. Andres, S.N.; Schellenberg, M.J.; Wallace, B.D.; Tumbale, P.; Williams, R.S. Recognition and repair of chemically heterogeneous structures at DNA ends. *Environ. Mol. Mutagen.* **2015**, *56*, 1–21. [[CrossRef](#)]
39. Turchi, J.J.; Henkels, K. Human Ku autoantigen binds cisplatin-damaged DNA but fails to stimulate human DNA-activated protein kinase. *J. Biol. Chem.* **1996**, *271*, 13861–13867. [[CrossRef](#)]
40. Yaneva, M.; Kowalewski, T.; Lieber, M.R. Interaction of DNA-dependent protein kinase with DNA and with Ku: Biochemical and atomic-force microscopy studies. *EMBO J.* **1997**, *16*, 5098–5112. [[CrossRef](#)]
41. Ilina, E.S.; Lavrik, O.I.; Khodyreva, S.N. Ku antigen interacts with abasic sites. *Biochim. Biophys. Acta* **2008**, *1784*, 1777–1785. [[CrossRef](#)]
42. Strande, N.; Roberts, S.A.; Oh, S.; Hendrickson, E.A.; Ramsden, D.A. Specificity of the dRP/AP lyase of Ku promotes nonhomologous end joining (NHEJ) fidelity at damaged ends. *J. Biol. Chem.* **2012**, *287*, 13686–13693. [[CrossRef](#)]
43. Roberts, S.A.; Strande, N.; Burkhalter, M.D.; Strom, C.; Havener, J.M.; Hasty, P.; Ramsden, D.A. Ku is a 5'-dRP/AP lyase that excises nucleotide damage near broken ends. *Nature* **2010**, *464*, 1214–1217. [[CrossRef](#)]
44. De Ory, A.; Zafra, O.; de Vega, M. Efficient processing of abasic sites by bacterial nonhomologous end-joining Ku proteins. *Nucleic Acids Res.* **2014**, *42*, 13082–13095. [[CrossRef](#)]
45. Walker, J.R.; Corpina, R.A.; Goldberg, J. Structure of the Ku heterodimer bound to DNA and its implications for double-strand break repair. *Nature* **2001**, *412*, 607–614. [[CrossRef](#)]
46. Rivera-Calzada, A.; Spagnolo, L.; Pearl, L.H.; Llorca, O. Structural model of full-length human Ku70-Ku80 heterodimer and its recognition of DNA and DNA-PKcs. *EMBO Rep.* **2007**, *8*, 56–62. [[CrossRef](#)]
47. Aissaoui, N.; Lai-Kee-Him, J.; Mills, A.; Declerck, N.; Morichaud, Z.; Brodolin, K.; Baconnais, S.; Le Cam, E.; Charbonnier, J.B.; Sounier, R.; et al. Modular Imaging Scaffold for Single-Particle Electron Microscopy. *ACS Nano* **2021**. [[CrossRef](#)] [[PubMed](#)]
48. Zhang, Z.; Zhu, L.; Lin, D.; Chen, F.; Chen, D.J.; Chen, Y. The three-dimensional structure of the C-terminal DNA-binding domain of human Ku70. *J. Biol. Chem.* **2001**, *276*, 38231–38236. [[CrossRef](#)] [[PubMed](#)]
49. Harris, R.; Esposito, D.; Sankar, A.; Maman, J.D.; Hinks, J.A.; Pearl, L.H.; Driscoll, P.C. The 3D solution structure of the C-terminal region of Ku86 (Ku86CTR). *J. Mol. Biol.* **2004**, *335*, 573–582. [[CrossRef](#)] [[PubMed](#)]
50. Zhang, Z.; Hu, W.; Cano, L.; Lee, T.D.; Chen, D.J.; Chen, Y. Solution structure of the C-terminal domain of Ku80 suggests important sites for protein-protein interactions. *Structure* **2004**, *12*, 495–502. [[CrossRef](#)]
51. Yoo, S.; Dynan, W.S. Geometry of a complex formed by double strand break repair proteins at a single DNA end: Recruitment of DNA-PKcs induces inward translocation of Ku protein. *Nucleic Acids Res.* **1999**, *27*, 4679–4686. [[CrossRef](#)]
52. Britton, S.; Coates, J.; Jackson, S.P. A new method for high-resolution imaging of Ku foci to decipher mechanisms of DNA double-strand break repair. *J. Cell Biol.* **2013**, *202*, 579–595. [[CrossRef](#)]
53. Lee, K.J.; Saha, J.; Sun, J.; Fattah, K.R.; Wang, S.C.; Jakob, B.; Chi, L.; Wang, S.Y.; Taucher-Scholz, G.; Davis, A.J.; et al. Phosphorylation of Ku dictates DNA double-strand break (DSB) repair pathway choice in S phase. *Nucleic Acids Res.* **2016**, *44*, 1732–1745. [[CrossRef](#)]

54. Zhang, W.W.; Yaneva, M. Reduced sulphhydryl groups are required for DNA binding of Ku protein. *Biochem. J.* **1993**, *293* (Pt 3), 769–774. [[CrossRef](#)]
55. Weterings, E.; Gallegos, A.C.; Dominick, L.N.; Cooke, L.S.; Bartels, T.N.; Vagner, J.; Matsunaga, T.O.; Mahadevan, D. A novel small molecule inhibitor of the DNA repair protein Ku70/80. *DNA Repair* **2016**, *43*, 98–106. [[CrossRef](#)]
56. Gavande, N.S.; VanderVere-Carozza, P.S.; Pawelczak, K.S.; Mendoza-Munoz, P.; Vernon, T.L.; Hanakahi, L.A.; Summerlin, M.; Dynlacht, J.R.; Farmer, A.H.; Sears, C.R.; et al. Discovery and development of novel DNA-PK inhibitors by targeting the unique Ku-DNA interaction. *Nucleic Acids Res.* **2020**, *48*, 11536–11550. [[CrossRef](#)]
57. Sibanda, B.L.; Chirgadze, D.Y.; Ascher, D.B.; Blundell, T.L. DNA-PKcs structure suggests an allosteric mechanism modulating DNA double-strand break repair. *Science* **2017**, *355*, 520–524. [[CrossRef](#)]
58. Sibanda, B.L.; Chirgadze, D.Y.; Blundell, T.L. Crystal structure of DNA-PKcs reveals a large open-ring cradle comprised of HEAT repeats. *Nature* **2010**, *463*, 118–121. [[CrossRef](#)]
59. Williams, D.R.; Lee, K.J.; Shi, J.; Chen, D.J.; Stewart, P.L. Cryo-EM structure of the DNA-dependent protein kinase catalytic subunit at subnanometer resolution reveals alpha helices and insight into DNA binding. *Structure* **2008**, *16*, 468–477. [[CrossRef](#)]
60. Sharif, H.; Li, Y.; Dong, Y.; Dong, L.; Wang, W.L.; Mao, Y.; Wu, H. Cryo-EM structure of the DNA-PK holoenzyme. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 7367–7372. [[CrossRef](#)]
61. Kaczmarek, W.; Khan, S.A. Lupus autoantigen Ku protein binds HIV-1 TAR RNA in vitro. *Biochem. Biophys. Res. Commun.* **1993**, *196*, 935–942. [[CrossRef](#)] [[PubMed](#)]
62. Yoo, S.; Dynan, W.S. Characterization of the RNA binding properties of Ku protein. *Biochemistry* **1998**, *37*, 1336–1343. [[CrossRef](#)] [[PubMed](#)]
63. Anisenko, A.N.; Knyazhanskaya, E.S.; Zatsepin, T.S.; Gottikh, M.B. Human Ku70 protein binds hairpin RNA and double stranded DNA through two different sites. *Biochimie* **2017**, *132*, 85–93. [[CrossRef](#)] [[PubMed](#)]
64. Sabath, K.; Staubli, M.L.; Marti, S.; Leitner, A.; Moes, M.; Jonas, S. INTS10-INTS13-INTS14 form a functional module of Integrator that binds nucleic acids and the cleavage module. *Nat. Commun.* **2020**, *11*, 3422. [[CrossRef](#)] [[PubMed](#)]
65. Prasad, R.; Shock, D.D.; Beard, W.A.; Wilson, S.H. Substrate channeling in mammalian base excision repair pathways: Passing the baton. *J. Biol. Chem.* **2010**, *285*, 40479–40488. [[CrossRef](#)] [[PubMed](#)]
66. DeFazio, L.G.; Stansel, R.M.; Griffith, J.D.; Chu, G. Synapsis of DNA ends by DNA-dependent protein kinase. *EMBO J.* **2002**, *21*, 3192–3200. [[CrossRef](#)] [[PubMed](#)]
67. Zhao, B.; Watanabe, G.; Morten, M.J.; Reid, D.A.; Rothenberg, E.; Lieber, M.R. The essential elements for the noncovalent association of two DNA ends during NHEJ synapsis. *Nat. Commun.* **2019**, *10*, 3588. [[CrossRef](#)]
68. Ramsden, D.A.; Gellert, M. Ku protein stimulates DNA end joining by mammalian DNA ligases: A direct role for Ku in repair of DNA double-strand breaks. *EMBO J.* **1998**, *17*, 609–614. [[CrossRef](#)]
69. Wang, J.L.; Duboc, C.; Wu, Q.; Ochi, T.; Liang, S.; Tsutakawa, S.E.; Lees-Miller, S.P.; Nadal, M.; Tainer, J.A.; Blundell, T.L.; et al. Dissection of DNA double-strand-break repair using novel single-molecule forceps. *Nat. Struct. Mol. Biol.* **2018**, *25*, 482–487. [[CrossRef](#)]
70. Lees-Miller, J.P.; Cobban, A.; Katsonis, P.; Bacolla, A.; Tsutakawa, S.E.; Hammel, M.; Meek, K.; Anderson, D.W.; Lichtarge, O.; Tainer, J.A.; et al. Uncovering DNA-PKcs ancient phylogeny, unique sequence motifs and insights for human disease. *Prog. Biophys. Mol. Biol.* **2020**. [[CrossRef](#)]
71. Chen, L.; Trujillo, K.; Ramos, W.; Sung, P.; Tomkinson, A.E. Promotion of Dnl4-catalyzed DNA end-joining by the Rad50/Mre11/Xrs2 and Hdf1/Hdf2 complexes. *Mol. Cell* **2001**, *8*, 1105–1115. [[CrossRef](#)]
72. Daley, J.M.; Palmbo, P.L.; Wu, D.; Wilson, T.E. Nonhomologous end joining in yeast. *Annu. Rev. Genet.* **2005**, *39*, 431–451. [[CrossRef](#)]
73. Arosio, D.; Costantini, S.; Kong, Y.; Vindigni, A. Fluorescence anisotropy studies on the Ku-DNA interaction: Anion and cation effects. *J. Biol. Chem.* **2004**, *279*, 42826–42835. [[CrossRef](#)]
74. Roberts, S.A.; Ramsden, D.A. Loading of the nonhomologous end joining factor, Ku, on protein-occluded DNA ends. *J. Biol. Chem.* **2007**, *282*, 10605–10613. [[CrossRef](#)]
75. Li, X.; Tyler, J.K. Nucleosome disassembly during human non-homologous end joining followed by concerted HIRA- and CAF-1-dependent reassembly. *Elife* **2016**, *5*. [[CrossRef](#)]
76. Costantini, S.; Woodbine, L.; Andreoli, L.; Jeggo, P.A.; Vindigni, A. Interaction of the Ku heterodimer with the DNA ligase IV/Xrcc4 complex and its regulation by DNA-PK. *DNA Repair* **2007**, *6*, 712–722. [[CrossRef](#)]
77. Chaplin, A.K.; Blundell, T.L. Structural biology of multicomponent assemblies in DNA double-strand-break repair through non-homologous end joining. *Curr. Opin. Struct. Biol.* **2019**, *61*, 9–16. [[CrossRef](#)]
78. Calsou, P.; Frit, P.; Humbert, O.; Muller, C.; Chen, D.J.; Salles, B. The DNA-dependent protein kinase catalytic activity regulates DNA end processing by means of Ku entry into DNA. *J. Biol. Chem.* **1999**, *274*, 7848–7856. [[CrossRef](#)]
79. Mari, P.O.; Florea, B.I.; Persengiev, S.P.; Verkaik, N.S.; Bruggenwirth, H.T.; Modesti, M.; Giglia-Mari, G.; Bezstarosti, K.; Demmers, J.A.; Luidier, T.M.; et al. Dynamic assembly of end-joining complexes requires interaction between Ku70/80 and XRCC4. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 18597–18602. [[CrossRef](#)]
80. Kim, J.S.; Krasieva, T.B.; Kurumizaka, H.; Chen, D.J.; Taylor, A.M.; Yokomori, K. Independent and sequential recruitment of NHEJ and HR factors to DNA damage sites in mammalian cells. *J. Cell Biol.* **2005**, *170*, 341–347. [[CrossRef](#)]

81. Kochan, J.A.; Desclos, E.C.B.; Bosch, R.; Meister, L.; Vriend, L.E.M.; van Attikum, H.; Krawczyk, P.M. Meta-analysis of DNA double-strand break response kinetics. *Nucleic Acids Res.* **2017**, *45*, 12625–12637. [[CrossRef](#)]
82. Haince, J.F.; McDonald, D.; Rodrigue, A.; Dery, U.; Masson, J.Y.; Hendzel, M.J.; Poirier, G.G. PARP1-dependent kinetics of recruitment of MRE11 and NBS1 proteins to multiple DNA damage sites. *J. Biol. Chem.* **2008**, *283*, 1197–1208. [[CrossRef](#)]
83. Wu, D.; Topper, L.M.; Wilson, T.E. Recruitment and dissociation of nonhomologous end joining proteins at a DNA double-strand break in *Saccharomyces cerevisiae*. *Genetics* **2008**, *178*, 1237–1249. [[CrossRef](#)]
84. Mimitou, E.P.; Symington, L.S. Ku prevents Exo1 and Sgs1-dependent resection of DNA ends in the absence of a functional MRX complex or Sae2. *EMBO J.* **2010**, *29*, 3358–3369. [[CrossRef](#)]
85. Zhang, Y.; Hefferin, M.L.; Chen, L.; Shim, E.Y.; Tseng, H.M.; Kwon, Y.; Sung, P.; Lee, S.E.; Tomkinson, A.E. Role of Dnl4-Lif1 in nonhomologous end-joining repair complex assembly and suppression of homologous recombination. *Nat. Struct. Mol. Biol.* **2007**, *14*, 639–646. [[CrossRef](#)]
86. Garcia, V.; Phelps, S.E.; Gray, S.; Neale, M.J. Bidirectional resection of DNA double-strand breaks by Mre11 and Exo1. *Nature* **2011**, *479*, 241–244. [[CrossRef](#)]
87. Chapman, J.R.; Taylor, M.R.; Boulton, S.J. Playing the end game: DNA double-strand break repair pathway choice. *Mol. Cell* **2012**, *47*, 497–510. [[CrossRef](#)]
88. Reginato, G.; Cannavo, E.; Cejka, P. Physiological protein blocks direct the Mre11-Rad50-Xrs2 and Sae2 nuclease complex to initiate DNA end resection. *Genes Dev.* **2017**, *31*, 2325–2330. [[CrossRef](#)]
89. Myler, L.R.; Gallardo, I.F.; Soniat, M.M.; Deshpande, R.A.; Gonzalez, X.B.; Kim, Y.; Paull, T.T.; Finkelstein, I.J. Single-Molecule Imaging Reveals How Mre11-Rad50-Nbs1 Initiates DNA Break Repair. *Mol. Cell* **2017**, *67*, 891–898.e4. [[CrossRef](#)]
90. Deshpande, R.A.; Myler, L.R.; Soniat, M.M.; Makharashvili, N.; Lee, L.; Lees-Miller, S.P.; Finkelstein, I.J.; Paull, T.T. DNA-dependent protein kinase promotes DNA end processing by MRN and CtIP. *Sci. Adv.* **2020**, *6*, eaay0922. [[CrossRef](#)]
91. Liang, F.; Jasin, M. Ku80-deficient cells exhibit excess degradation of extrachromosomal DNA. *J. Biol. Chem.* **1996**, *271*, 14405–14411. [[CrossRef](#)] [[PubMed](#)]
92. Clerici, M.; Mantiero, D.; Guerini, I.; Lucchini, G.; Longhese, M.P. The Yku70-Yku80 complex contributes to regulate double-strand break processing and checkpoint activation during the cell cycle. *EMBO Rep.* **2008**, *9*, 810–818. [[CrossRef](#)] [[PubMed](#)]
93. Sun, J.; Lee, K.J.; Davis, A.J.; Chen, D.J. Human Ku70/80 protein blocks exonuclease 1-mediated DNA resection in the presence of human Mre11 or Mre11/Rad50 protein complex. *J. Biol. Chem.* **2012**, *287*, 4936–4945. [[CrossRef](#)] [[PubMed](#)]
94. Cheng, Q.; Barboule, N.; Frit, P.; Gomez, D.; Bombarde, O.; Couderc, B.; Ren, G.S.; Salles, B.; Calsou, P. Ku counteracts mobilization of PARP1 and MRN in chromatin damaged with DNA double-strand breaks. *Nucleic Acids Res.* **2011**, *39*, 9605–9619. [[CrossRef](#)] [[PubMed](#)]
95. Moldovan, G.L.; Pfander, B.; Jentsch, S. PCNA, the maestro of the replication fork. *Cell* **2007**, *129*, 665–679. [[CrossRef](#)]
96. Postow, L.; Ghenoiu, C.; Woo, E.M.; Krutchinsky, A.N.; Chait, B.T.; Funabiki, H. Ku80 removal from DNA through double strand break-induced ubiquitylation. *J. Cell Biol.* **2008**, *182*, 467–479. [[CrossRef](#)]
97. Ismail, I.H.; Gagne, J.P.; Genois, M.M.; Strickfaden, H.; McDonald, D.; Xu, Z.; Poirier, G.G.; Masson, J.Y.; Hendzel, M.J. The RNF138 E3 ligase displaces Ku to promote DNA end resection and regulate DNA repair pathway choice. *Nat. Cell Biol.* **2015**, *17*, 1446–1457. [[CrossRef](#)]
98. Schmidt, C.K.; Galanty, Y.; Sczaniecka-Clift, M.; Coates, J.; Jhujh, S.; Demir, M.; Cornwell, M.; Beli, P.; Jackson, S.P. Systematic E2 screening reveals a UBE2D-RNF138-CtIP axis promoting DNA repair. *Nat. Cell Biol.* **2015**, *17*, 1458–1470. [[CrossRef](#)]
99. Ishida, N.; Nakagawa, T.; Iemura, S.I.; Yasui, A.; Shima, H.; Katoh, Y.; Nagasawa, Y.; Natsume, T.; Igarashi, K.; Nakayama, K. Ubiquitylation of Ku80 by RNF126 Promotes Completion of Nonhomologous End Joining-Mediated DNA Repair. *Mol. Cell Biol.* **2017**, *37*. [[CrossRef](#)]
100. Feng, L.; Chen, J. The E3 ligase RNF8 regulates KU80 removal and NHEJ repair. *Nat. Struct. Mol. Biol.* **2012**, *19*, 201–206. [[CrossRef](#)]
101. Van den Boom, J.; Wolf, M.; Weimann, L.; Schulze, N.; Li, F.; Kaschani, F.; Riemer, A.; Zierhut, C.; Kaiser, M.; Iliakis, G.; et al. VCP/p97 Extracts Sterically Trapped Ku70/80 Rings from DNA in Double-Strand Break Repair. *Mol. Cell* **2016**, *64*, 189–198. [[CrossRef](#)]
102. Sharma, A.; Alswillah, T.; Kapoor, I.; Debjani, P.; Willard, B.; Summers, M.K.; Gong, Z.; Almasan, A. USP14 is a deubiquitinase for Ku70 and critical determinant of non-homologous end joining repair in autophagy and PTEN-deficient cells. *Nucleic Acids Res.* **2020**, *48*, 736–747. [[CrossRef](#)]
103. Brown, J.S.; Lukashchuk, N.; Sczaniecka-Clift, M.; Britton, S.; le Sage, C.; Calsou, P.; Beli, P.; Galanty, Y.; Jackson, S.P. Neddylation promotes ubiquitylation and release of Ku from DNA-damage sites. *Cell Rep.* **2015**, *11*, 704–714. [[CrossRef](#)]
104. Reid, D.A.; Keegan, S.; Leo-Macias, A.; Watanabe, G.; Strande, N.T.; Chang, H.H.; Oksuz, B.A.; Fenyo, D.; Lieber, M.R.; Ramsden, D.A.; et al. Organization and dynamics of the nonhomologous end-joining machinery during DNA double-strand break repair. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, E2575–E2584. [[CrossRef](#)]
105. Ropars, V.; Drevet, P.; Legrand, P.; Baconnais, S.; Amram, J.; Faure, G.; Marquez, J.A.; Pietrement, O.; Guerois, R.; Callebaut, I.; et al. Structural characterization of filaments formed by human Xrcc4-Cernunnos/XLF complex involved in nonhomologous DNA end-joining. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 12663–12668. [[CrossRef](#)]
106. Ryan, A.J.; Squires, S.; Strutt, H.L.; Johnson, R.T. Camptothecin cytotoxicity in mammalian cells is associated with the induction of persistent double strand breaks in replicating DNA. *Nucleic Acids Res.* **1991**, *19*, 3295–3300. [[CrossRef](#)]

107. Chanut, P.; Britton, S.; Coates, J.; Jackson, S.P.; Calsou, P. Coordinated nuclease activities counteract Ku at single-ended DNA double-strand breaks. *Nat. Commun.* **2016**, *7*, 12889. [[CrossRef](#)]
108. Britton, S.; Chanut, P.; Delteil, C.; Barboule, N.; Frit, P.; Calsou, P. ATM antagonizes NHEJ proteins assembly and DNA-ends synapsis at single-ended DNA double strand breaks. *Nucleic Acids Res.* **2020**, *48*, 9710–9723. [[CrossRef](#)]
109. Balestrini, A.; Ristic, D.; Dionne, I.; Liu, X.Z.; Wyman, C.; Wellinger, R.J.; Petrini, J.H. The Ku Heterodimer and the Metabolism of Single-Ended DNA Double-Strand Breaks. *Cell Rep.* **2013**, *3*, 2033–2045. [[CrossRef](#)]
110. Teixeira-Silva, A.; Ait Saada, A.; Hardy, J.; Iraqui, I.; Nocente, M.C.; Freon, K.; Lambert, S.A.E. The end-joining factor Ku acts in the end-resection of double strand break-free arrested replication forks. *Nat. Commun.* **2017**, *8*, 1982. [[CrossRef](#)]
111. Wellinger, R.J.; Zakian, V.A. Everything you ever wanted to know about *Saccharomyces cerevisiae* telomeres: Beginning to end. *Genetics* **2012**, *191*, 1073–1105. [[CrossRef](#)] [[PubMed](#)]
112. Boulton, S.J.; Jackson, S.P. Identification of a *Saccharomyces cerevisiae* Ku80 homologue: Roles in DNA double strand break rejoining and in telomeric maintenance. *Nucleic Acids Res.* **1996**, *24*, 4639–4648. [[CrossRef](#)] [[PubMed](#)]
113. Porter, S.E.; Greenwell, P.W.; Ritchie, K.B.; Petes, T.D. The DNA-binding protein Hdf1p (a putative Ku homologue) is required for maintaining normal telomere length in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* **1996**, *24*, 582–585. [[CrossRef](#)] [[PubMed](#)]
114. Lopez, C.R.; Ribes-Zamora, A.; Indiviglio, S.M.; Williams, C.L.; Haricharan, S.; Bertuch, A.A. Ku must load directly onto the chromosome end in order to mediate its telomeric functions. *PLoS Genet.* **2011**, *7*, e1002233. [[CrossRef](#)]
115. Gravel, S.; Larrivee, M.; Labrecque, P.; Wellinger, R.J. Yeast Ku as a regulator of chromosomal DNA end structure. *Science* **1998**, *280*, 741–744. [[CrossRef](#)]
116. Polotnianka, R.M.; Li, J.; Lustig, A.J. The yeast Ku heterodimer is essential for protection of the telomere against nucleolytic and recombinational activities. *Curr Biol.* **1998**, *8*, 831–834. [[CrossRef](#)]
117. Maringele, L.; Lydall, D. EXO1-dependent single-stranded DNA at telomeres activates subsets of DNA damage and spindle checkpoint pathways in budding yeast yku70Delta mutants. *Genes Dev.* **2002**, *16*, 1919–1933. [[CrossRef](#)]
118. Bonetti, C.; Surace, E.M. Mouse embryonic retina delivers information controlling cortical neurogenesis. *PLoS ONE* **2010**, *5*, e15211. [[CrossRef](#)]
119. Peterson, S.E.; Stellwagen, A.E.; Diede, S.J.; Singer, M.S.; Haimberger, Z.W.; Johnson, C.O.; Tzoneva, M.; Gottschling, D.E. The function of a stem-loop in telomerase RNA is linked to the DNA repair protein Ku. *Nat. Genet.* **2001**, *27*, 64–67. [[CrossRef](#)]
120. Roy, R.; Meier, B.; McAinsh, A.D.; Feldmann, H.M.; Jackson, S.P. Separation-of-function mutants of yeast Ku80 reveal a Yku80p-Sir4p interaction involved in telomeric silencing. *J. Biol. Chem.* **2004**, *279*, 86–94. [[CrossRef](#)]
121. Fisher, T.S.; Taggart, A.K.; Zakian, V.A. Cell cycle-dependent regulation of yeast telomerase by Ku. *Nat. Struct. Mol. Biol.* **2004**, *11*, 1198–1205. [[CrossRef](#)]
122. Chen, H.; Xue, J.; Churikov, D.; Hass, E.P.; Shi, S.; Lemon, L.D.; Luciano, P.; Bertuch, A.A.; Zappulla, D.C.; Geli, V.; et al. Structural Insights into Yeast Telomerase Recruitment to Telomeres. *Cell* **2018**, *172*, 331–343. [[CrossRef](#)]
123. Nandakumar, J.; Cech, T.R. Finding the end: Recruitment of telomerase to telomeres. *Nat. Rev. Mol. Cell Biol.* **2013**, *14*, 69–82. [[CrossRef](#)]
124. Taddei, A.; Schober, H.; Gasser, S.M. The budding yeast nucleus. *Cold Spring Harb. Perspect. Biol.* **2010**, *2*, a000612. [[CrossRef](#)]
125. Taddei, A.; Hediger, F.; Neumann, F.R.; Bauer, C.; Gasser, S.M. Separation of silencing from perinuclear anchoring functions in yeast Ku80, Sir4 and Esc1 proteins. *EMBO J.* **2004**, *23*, 1301–1312. [[CrossRef](#)]
126. Hsu, H.L.; Gilley, D.; Blackburn, E.H.; Chen, D.J. Ku is associated with the telomere in mammals. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 12454–12458. [[CrossRef](#)]
127. Wang, Y.; Ghosh, G.; Hendrickson, E.A. Ku86 represses lethal telomere deletion events in human somatic cells. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 12430–12435. [[CrossRef](#)]
128. Hsu, H.L.; Gilley, D.; Galande, S.A.; Hande, M.P.; Allen, B.; Kim, S.H.; Li, G.C.; Campisi, J.; Kohwi-Shigematsu, T.; Chen, D.J. Ku acts in a unique way at the mammalian telomere to prevent end joining. *Genes Dev.* **2000**, *14*, 2807–2812. [[CrossRef](#)]
129. Bianchi, A.; de Lange, T. Ku binds telomeric DNA in vitro. *J. Biol. Chem.* **1999**, *274*, 21223–21227. [[CrossRef](#)]
130. De Lange, T. Shelterin-Mediated Telomere Protection. *Annu. Rev. Genet.* **2018**, *52*, 223–247. [[CrossRef](#)]
131. Bombarde, O.; Boby, C.; Gomez, D.; Frit, P.; Giraud-Panis, M.J.; Gilson, E.; Salles, B.; Calsou, P. TRF2/RAP1 and DNA-PK mediate a double protection against joining at telomeric ends. *EMBO J.* **2010**, *29*, 1573–1584. [[CrossRef](#)]
132. Sfeir, A.; de Lange, T. Removal of shelterin reveals the telomere end-protection problem. *Science* **2012**, *336*, 593–597. [[CrossRef](#)]
133. Ting, N.S.; Yu, Y.; Pohorelic, B.; Lees-Miller, S.P.; Beattie, T.L. Human Ku70/80 interacts directly with hTR, the RNA component of human telomerase. *Nucleic Acids Res.* **2005**, *33*, 2090–2098. [[CrossRef](#)]
134. Giffin, W.; Torrance, H.; Rodda, D.J.; Prefontaine, G.G.; Pope, L.; Hache, R.J. Sequence-specific DNA binding by Ku autoantigen and its effects on transcription. *Nature* **1996**, *380*, 265–268. [[CrossRef](#)]
135. Hu, S.; Pluth, J.M.; Cucinotta, F.A. Putative binding modes of Ku70-SAP domain with double strand DNA: A molecular modeling study. *J. Mol. Model.* **2012**, *18*, 2163–2174. [[CrossRef](#)]
136. Chan, D.W.; Ye, R.; Veillette, C.J.; Lees-Miller, S.P. DNA-dependent protein kinase phosphorylation sites in Ku 70/80 heterodimer. *Biochemistry* **1999**, *38*, 1819–1828. [[CrossRef](#)]
137. Douglas, P.; Gupta, S.; Morrice, N.; Meek, K.; Lees-Miller, S.P. DNA-PK-dependent phosphorylation of Ku70/80 is not required for non-homologous end joining. *DNA Repair* **2005**, *4*, 1006–1018. [[CrossRef](#)]

138. Bouley, J.; Saad, L.; Grall, R.; Schellenbauer, A.; Biard, D.; Paget, V.; Morel-Altmeier, S.; Guipaud, O.; Chambon, C.; Salles, B.; et al. A new phosphorylated form of Ku70 identified in resistant leukemic cells confers fast but unfaithful DNA repair in cancer cell lines. *Oncotarget* **2015**, *6*, 27980–28000. [[CrossRef](#)]
139. Zhao, X.; Blobel, G. A SUMO ligase is part of a nuclear multiprotein complex that affects DNA repair and chromosomal organization. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 4777–4782. [[CrossRef](#)]
140. Cremona, C.A.; Sarangi, P.; Yang, Y.; Hang, L.E.; Rahman, S.; Zhao, X. Extensive DNA damage-induced sumoylation contributes to replication and repair and acts in addition to the mecl1 checkpoint. *Mol. Cell* **2012**, *45*, 422–432. [[CrossRef](#)] [[PubMed](#)]
141. Hang, L.E.; Lopez, C.R.; Liu, X.; Williams, J.M.; Chung, I.; Wei, L.; Bertuch, A.A.; Zhao, X. Regulation of Ku-DNA association by Yku70 C-terminal tail and SUMO modification. *J. Biol. Chem.* **2014**, *289*, 10308–10317. [[CrossRef](#)] [[PubMed](#)]
142. Hendriks, I.A.; Lyon, D.; Young, C.; Jensen, L.J.; Vertegaal, A.C.; Nielsen, M.L. Site-specific mapping of the human SUMO proteome reveals co-modification with phosphorylation. *Nat. Struct. Mol. Biol.* **2017**, *24*, 325–336. [[CrossRef](#)] [[PubMed](#)]
143. Li, B.; Navarro, S.; Kasahara, N.; Comai, L. Identification and biochemical characterization of a Werner's syndrome protein complex with Ku70/80 and poly(ADP-ribose) polymerase-1. *J. Biol. Chem.* **2004**, *279*, 13659–13667. [[CrossRef](#)] [[PubMed](#)]
144. Hohegger, H.; Dejsuphong, D.; Fukushima, T.; Morrison, C.; Sonoda, E.; Schreiber, V.; Zhao, G.Y.; Saberi, A.; Masutani, M.; Adachi, N.; et al. Parp-1 protects homologous recombination from interference by Ku and Ligase IV in vertebrate cells. *EMBO J.* **2006**, *25*, 1305–1314. [[CrossRef](#)]
145. Sawada, M.; Sun, W.; Hayes, P.; Leskov, K.; Boothman, D.A.; Matsuyama, S. Ku70 suppresses the apoptotic translocation of Bax to mitochondria. *Nat. Cell Biol.* **2003**, *5*, 320–329. [[CrossRef](#)]
146. Sawada, M.; Hayes, P.; Matsuyama, S. Cytoprotective membrane-permeable peptides designed from the Bax-binding domain of Ku70. *Nat. Cell Biol.* **2003**, *5*, 352–357. [[CrossRef](#)]
147. Cohen, H.Y.; Lavu, S.; Bitterman, K.J.; Hekking, B.; Imahiyerobo, T.A.; Miller, C.; Frye, R.; Ploegh, H.; Kessler, B.M.; Sinclair, D.A. Acetylation of the C terminus of Ku70 by CBP and PCAF controls Bax-mediated apoptosis. *Mol. Cell* **2004**, *13*, 627–638. [[CrossRef](#)]
148. Lamaa, A.; Le Bras, M.; Skuli, N.; Britton, S.; Frit, P.; Calsou, P.; Prats, H.; Cammas, A.; Millevoi, S. A novel cytoprotective function for the DNA repair protein Ku in regulating p53 mRNA translation and function. *EMBO Rep.* **2016**, *17*, 508–518. [[CrossRef](#)]
149. Andrews, B.J.; Lehman, J.A.; Turchi, J.J. Kinetic analysis of the Ku-DNA binding activity reveals a redox-dependent alteration in protein structure that stimulates dissociation of the Ku-DNA complex. *J. Biol. Chem.* **2006**, *281*, 13596–13603. [[CrossRef](#)]
150. Ayene, I.S.; Stamato, T.D.; Mauldin, S.K.; Biaglow, J.E.; Tuttle, S.W.; Jenkins, S.F.; Koch, C.J. Mutation in the glucose-6-phosphate dehydrogenase gene leads to inactivation of Ku DNA end binding during oxidative stress. *J. Biol. Chem.* **2002**, *277*, 9929–9935. [[CrossRef](#)]
151. Hanakahi, L.A.; Bartlett-Jones, M.; Chappell, C.; Pappin, D.; West, S.C. Binding of inositol phosphate to DNA-PK and stimulation of double-strand break repair. *Cell* **2000**, *102*, 721–729. [[CrossRef](#)]
152. Hanakahi, L.A.; West, S.C. Specific interaction of IP6 with human Ku70/80, the DNA-binding subunit of DNA-PK. *EMBO J.* **2002**, *21*, 2038–2044. [[CrossRef](#)]
153. Ma, Y.; Lieber, M.R. Binding of inositol hexakisphosphate (IP6) to Ku but not to DNA-PKcs. *J. Biol. Chem.* **2002**, *277*, 10756–10759. [[CrossRef](#)]
154. Cheung, J.C.; Salerno, B.; Hanakahi, L.A. Evidence for an inositol hexakisphosphate-dependent role for Ku in mammalian nonhomologous end joining that is independent of its role in the DNA-dependent protein kinase. *Nucleic Acids Res.* **2008**, *36*, 5713–5726. [[CrossRef](#)]
155. Doherty, A.J.; Jackson, S.P.; Weller, G.R. Identification of bacterial homologues of the Ku DNA repair proteins. *FEBS Lett.* **2001**, *500*, 186–188. [[CrossRef](#)]
156. Pitcher, R.S.; Brissett, N.C.; Doherty, A.J. Nonhomologous end-joining in bacteria: A microbial perspective. *Annu. Rev. Microbiol.* **2007**, *61*, 259–282. [[CrossRef](#)]
157. Milne, G.T.; Jin, S.; Shannon, K.B.; Weaver, D.T. Mutations in two Ku homologs define a DNA end-joining repair pathway in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* **1996**, *16*, 4189–4198. [[CrossRef](#)]
158. Boulton, S.J.; Jackson, S.P. *Saccharomyces cerevisiae* Ku70 potentiates illegitimate DNA double-strand break repair and serves as a barrier to error-prone DNA repair pathways. *EMBO J.* **1996**, *15*, 5093–5103. [[CrossRef](#)]
159. Manolis, K.G.; Nimmo, E.R.; Hartsuiker, E.; Carr, A.M.; Jeggo, P.A.; Allshire, R.C. Novel functional requirements for non-homologous DNA end joining in *Schizosaccharomyces pombe*. *EMBO J.* **2001**, *20*, 210–221. [[CrossRef](#)]
160. D'Adda di Fagagna, F.; Weller, G.R.; Doherty, A.J.; Jackson, S.P. The Gam protein of bacteriophage Mu is an orthologue of eukaryotic Ku. *EMBO Rep.* **2003**, *4*, 47–52. [[CrossRef](#)]
161. Bartlett, E.J.; Brissett, N.C.; Doherty, A.J. Ribonucleolytic resection is required for repair of strand displaced nonhomologous end-joining intermediates. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, E1984–E1991. [[CrossRef](#)] [[PubMed](#)]
162. Aravind, L.; Koonin, E.V. Prokaryotic homologs of the eukaryotic DNA-end-binding protein Ku, novel domains in the Ku protein and prediction of a prokaryotic double-strand break repair system. *Genome Res.* **2001**, *11*, 1365–1374. [[CrossRef](#)] [[PubMed](#)]
163. Bertrand, C.; Thibessard, A.; Bruand, C.; Lecointe, F.; Leblond, P. Bacterial NHEJ: A never ending story. *Mol. Microbiol.* **2019**, *111*, 1139–1151. [[CrossRef](#)] [[PubMed](#)]
164. Dupuy, P.; Sauviac, L.; Bruand, C. Stress-inducible NHEJ in bacteria: Function in DNA repair and acquisition of heterologous DNA. *Nucleic Acids Res.* **2019**, *47*, 1335–1349. [[CrossRef](#)] [[PubMed](#)]

165. Weller, G.R.; Kysela, B.; Roy, R.; Tonkin, L.M.; Scanlan, E.; Della, M.; Devine, S.K.; Day, J.P.; Wilkinson, A.; d'Adda di Fagagna, F.; et al. Identification of a DNA nonhomologous end-joining complex in bacteria. *Science* **2002**, *297*, 1686–1689. [[CrossRef](#)] [[PubMed](#)]
166. McGovern, S.; Baconnais, S.; Roblin, P.; Nicolas, P.; Drevet, P.; Simonson, H.; Pietrement, O.; Charbonnier, J.B.; Le Cam, E.; Noirot, P.; et al. C-terminal region of bacterial Ku controls DNA bridging, DNA threading and recruitment of DNA ligase D for double strand breaks repair. *Nucleic Acids Res.* **2016**, *44*, 4785–4806. [[CrossRef](#)]
167. Oz, R.; Wang, J.L.; Guerois, R.; Goyal, G.; Kk, S.; Ropars, V.; Sharma, R.; Koca, F.; Charbonnier, J.B.; Modesti, M.; et al. Dynamics of Ku and bacterial non-homologous end-joining characterized using single DNA molecule analysis. *Nucleic Acids Res.* **2021**. [[CrossRef](#)]
168. Shao, Z.; Davis, A.J.; Fattah, K.R.; So, S.; Sun, J.; Lee, K.J.; Harrison, L.; Yang, J.; Chen, D.J. Persistently bound Ku at DNA ends attenuates DNA end resection and homologous recombination. *DNA Repair* **2012**, *11*, 310–316. [[CrossRef](#)]
169. Marmignon, A.; Bischerour, J.; Silve, A.; Fojcik, C.; Dubois, E.; Arnaiz, O.; Kapusta, A.; Malinsky, S.; Betermier, M. Ku-mediated coupling of DNA cleavage and repair during programmed genome rearrangements in the ciliate *Paramecium tetraurelia*. *PLoS Genet.* **2014**, *10*, e1004552. [[CrossRef](#)]
170. Abello, A.; Regnier, V.; Arnaiz, O.; Le Bars, R.; Betermier, M.; Bischerour, J. Functional diversification of *Paramecium* Ku80 paralogs safeguards genome integrity during precise programmed DNA elimination. *PLoS Genet.* **2020**, *16*, e1008723. [[CrossRef](#)]
171. Betermier, M.; Borde, V.; de Villartay, J.P. Coupling DNA Damage and Repair: An Essential Safeguard during Programmed DNA Double-Strand Breaks? *Trends Cell Biol.* **2020**, *30*, 87–96. [[CrossRef](#)]
172. Jackson, S.P.; Bartek, J. The DNA-damage response in human biology and disease. *Nature* **2009**, *461*, 1071–1078. [[CrossRef](#)]