

Review

Secondary Metabolites of *Purpureocillium lilacinum*

Wei Chen  and Qiongbo Hu * 

Key Laboratory of Bio-Pesticide Innovation and Application of Guangdong Province, College of Plant Protection, South China Agricultural University, Guangzhou 510642, China; cw@stu.scau.edu.cn

* Correspondence: hqbscau@scau.edu.cn

Abstract: Fungi can synthesize a wealth of secondary metabolites, which are widely used in the exploration of lead compounds of pharmaceutical or agricultural importance. *Beauveria*, *Metarhizium*, and *Cordyceps* are the most extensively studied fungi in which a large number of biologically active metabolites have been identified. However, relatively little attention has been paid to *Purpureocillium lilacinum*. *P. lilacinum* are soil-habituated fungi that are widely distributed in nature and are very important biocontrol fungi in agriculture, providing good biological control of plant parasitic nematodes and having a significant effect on *Aphidoidea*, *Tetranychus cinnabarinus*, and *Aleyrodidae*. At the same time, it produces secondary metabolites with various biological activities such as anticancer, antimicrobial, and insecticidal. This review attempts to provide a comprehensive overview of the secondary metabolites of *P. lilacinum*, with emphasis on the chemical diversity and biological activity of these secondary metabolites and the biosynthetic pathways, and gives new insight into the secondary metabolites of medical and entomogenous fungi, which is expected to provide a reference for the development of medicine and agrochemicals in the future.

Keywords: entomogenous fungi; biosynthesis; biocontrol; nematodes



Citation: Chen, W.; Hu, Q. Secondary Metabolites of *Purpureocillium lilacinum*. *Molecules* **2022**, *27*, 18. <https://doi.org/10.3390/molecules27010018>

Academic Editor: Josphat Matasyoh

Received: 22 November 2021

Accepted: 17 December 2021

Published: 21 December 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The genus *Purpureocillium* in the Ophiocordycipitaceae family was structured by Luangsa-Ard et al. In 2011, based on the medical importance, the *Purpureocillium lilacinum* was designated as the type species of the *Paecilomyces* genus [1]. This species was nominated as *Penicillium lilacinum* by Thom in 1901, and then it was revised as *Paecilomyces lilacinus* by Samson in 1974 [2]. After comparing the 18S rRNA gene, internal transcribed spacer, and partial translation elongation factor 1- α sequences with *P. lilacinus*, Luangsa-Ard proposed a new genus name *Purpureocillium* and made the new combination *P. lilacinum* in 2011. The fungus was found in a wide range of land and marine environments [3–5]. They are often isolated from insects, nematodes, and the rhizosphere of many crops [6–8]. The species can grow in a wide range of temperatures from 8 to 38 °C with optimal temperatures of 26–30 °C [3]. It also has a wide pH tolerance and can grow on a variety of substrates [9]. This fungus has promising potential as a biocontrol agent to control crops' root-knot nematodes [10,11]. The parasitism of nematodes is that the hyphae directly invade the surface of nematodes' eggs and then produce appressoria on the surface, which infects the nematodes' eggs after adsorption. In the process of infection, *P. lilacinum* secretes a variety of enzymes, such as serine protease and chitinase, which can lead to the degradation of protein and chitin components of the nematode epidermis, which is conducive to the invasion of fungi and destruction of cell components. It has been shown that the fermentation filtrate of *P. lilacinum* can inhibit the mycelial growth of the pathogenic fungi *Helminthosporium maydis* and *Fusarium graminearum*, and has a significant inhibitory effect on the spore germination of *Fusarium oxysporum* [12,13]. Currently, there are eight registered pesticide products of *P. lilacinum* in China used to control root-knot nematodes (<http://www.chinapesticide.org.cn/hysj/index.jhtml>, accessed on 23 February 2021); similar pesticides are also registered in the USA (<https://iaspub.epa.gov/apex/pesticides>, accessed on 23 February 2021) and

European Union (<http://www.efsa.europa.eu/>, accessed on 23 February 2021). In addition, *P. lilacinum* has been shown to be effective against *Phyllotreta striolata*, *Thrips palmi*, and predatory mite [14,15]. However, the *P. lilacinum* strains used for biocontrol agents have a high identity with those strains causing infections in humans [16,17].

Secondary metabolites are produced in a certain growth period of plants and microorganisms. They are small molecules with complex chemical structures that are not necessary for growth and reproduction, such as pigments, hormones, toxins, and antibiotics [18]. Fungi are important organisms that produce active secondary metabolites. Different kinds of fungi produce different secondary metabolites. The discovery of fungal secondary metabolites has become an important source of new drugs and pesticides [19]. Fumosorinone was isolated from the *Isaria fumosorosea*, and it is a potential medicine for the treatment of type II diabetes and other associated metabolic syndromes [20]. Diorcinol K, D, and I were isolated from *Aspergillus*, displaying significant antibacterial activities against *Staphylococcus aureus* and methicillin-resistant *S. aureus* [21]. Pyrenocine A was produced by *Paecilomyces* and showed a significant antitrypanosomal activity against *Trypanosoma brucei* [22]. Paeciloxanthone was isolated from the extracts of *Paecilomyces* sp. and showed significant cytotoxicity against HepG2 cell lines [23]. The research on the synthesis and regulation of secondary metabolites is helpful to develop new active compounds and increase the output of active compounds. The common secondary metabolites of fungi are polyketones, nonribosome peptides, sterols, alkaloids, and terpenes.


Several interesting reviews related to various aspects of *Paecilomyces* species have been published. For example, Weng reviewed the risks of *Isaria fumosorosea* (formerly *P. fumosoroseus*) and *I. farinose* (formerly *P. farinosus*) and their mycotoxins, including their structures, bioactivities, and toxicities [24]. Zhang summarized the mycotoxins of *I. cicadae* (formerly *P. cicadae*) and *I. tenuipes* (formerly *P. tenuipes*), as well as their risk evaluation [25]. As far as we know, a comprehensive overview of *P. lilacinum*, with an emphasis on the chemical diversity, relevant biological activities, and biosynthesis of these metabolites, remains untouched. In this paper, according to the published research reports in recent years, we have sorted out the secondary metabolites of *P. lilacinum* and described the sources, chemical structures, and bioactivities of the reported compounds with particular emphasis on their potential use as drug lead compounds and general biosynthesis pathways. We provide a reference for the follow-up study and ensure more secure and efficient use of fungal products.

2. Leucinostatins

Leucinostatins is a class of peptide mycotoxins that have some potent effects on liver cells after oral administration, and they were used as both antimicrobial and antitumor agents through interactions with the membrane phospholipids [26].

In the early 1970s, Tadashi and his team isolated a new antibiotic from the culture medium of *P. lilacinum*, named leucinostatin (Figure 1, Table 1). It was found that leucinostatin A, B, C, and D were active against yeasts including pathogenic and nonpathogenic strains and filamentous fungi, and were moderately active against Gram-positive bacteria, but more important is its anticancer activity, which is cytotoxic to the HeLa cell culture and inhibits Ehrlich subcutaneous solid tumors and prostate cancer [27,28]. A new report indicated that leucinostatin has a significant antihypertensive effect on rabbit blood pressure, but does not affect the reaction of adrenaline and acetylcholine [29]. It is worth mentioning that leucinostatin is one of the active substances to kill nematodes. Manabu suggested that Leucinostatin A inhibited prostate cancer cell growth through the reduction in insulin-like growth factor I expression in prostate stromal cells [30]. In addition, the use of leucinostatin A-loaded nanospheres could represent a new promising therapeutic system against *Candida* infection [31]. Leucinostatin Y exhibited the promising importance of the C-terminal of leucinostatins on the preferential cytotoxicity and inhibition of mitochondrial function of cancer cells in the absence of glucose [32]. It is not difficult to see that leucinostatins are potential anticancer compounds, but it should also be noted that

the acute toxicities determined by intraperitoneal LD₅₀ values for leucinostatins A and B hydrochlorides were found to be 1.8 mg/kg in mice body weight. Thus, it should be evaluated more carefully when it comes to drug use [33,34]. In addition, Yuzuru separated to obtain leucinostatin A, B, C, D, H, and K in *P. lilacinum* and found that the different isolates all produced leucinostatins, which means that they may play an important role not only in the infection process of mycoses in the human case, but also with insects and nematodes [26].



- A: R = $-\text{CH}_2(\text{CH}_2\text{COCH}_2\text{CH}_3)\text{OOCCH}_2\text{CH}_2\text{COOH}$, R₁ = R₂ = R₅ = R₆ = $-\text{CH}_3$, R₃ = H, R₄ = $-\text{CH}(\text{CH}_3)_2$
 B: R = $-\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{COCH}_2\text{CH}_3$, R₁ = R₂ = R₅ = $-\text{CH}_3$, R₃ = R₆ = H, R₄ = $-\text{CH}(\text{CH}_2)_2$
 C: R = $-\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{COCH}_2\text{CH}_3$, R₁ = R₂ = R₃ = R₄ = $-\text{CH}_3$, R₅ = R₆ = H
 D: R = R₃ = R₄ = R₅ = $-\text{CH}_3$, R₁ = $-\text{CH}(\text{CH}_3)_2$, R₂ = R₆ = H

Figure 1. The structures of leucinostatins.

3. PK Metabolites

3.1. Acremonidins and Acremoxanthones

Acremonidins (Figure 2, Table 1) and acremoxanthones (Figure 3, Table 1) are kinds of polyketides (PKs), which contain alternating carbonyl and methylene groups, usually involved in cell defense or intercellular communication.




Figure 2. The structures of acremonidins.





Figure 3. The structures of acremoxanthones.

Acremoxanthone and acremonidin are xanthone-anthraquinone heterodimers, both calmodulin inhibitors produced by *P. lilacinum* [35]. Both of them exhibited anti-*Bacillus cereus*, antibacterial, antifungal, antiplasmodial, and cytotoxic activity [36]. In addition, acremonidin A and C, and acremoxanthone C and D were found to be moderate for 20 s proteasome inhibitory activity. Acremonidin A and Acremoxanthone C also have high affinity with human calmodulin biosensors. This means that they are also calmodulin inhibitors [37]. In addition, researchers found that acremoxanthone C has anti-oomycete activities [38]. Two new compounds acremoxanthones F and G represented antimalarial activity against the *Plasmodium falciparum* K1 strain, a multidrug-resistant strain [39]. These studies indicate that acremonidins and acremoxanthones all have high medicinal and economic value, but their toxicological effects need to be evaluated in more detail.

3.2. Paecilomide

Teles found a new α -pyridone Paecilomide (Figure 4, Table 1) by adding salmonella typhimurium to a flask containing *P. lilacinum* in two different concentrations and three different forms at two different stages of the development of *P. lilacinum*. He observed that adding a small amount of inoculum produced a living extract, while adding a large amount of inoculum led to the extraction of active extract, paecilomide [40]. It is an acetylcholinesterase inhibitor [41,42].



paecilomide

Figure 4. The structure of paecilomide.

3.3. Pyrones

Elbandy isolated 13 compounds from the sponge-derived fungus *P. lilacinum* in 2009, four of which were α -pyrones: paecilopyrone A, paecilopyrone B, phomapyrone B, and phomapyrone C; and one γ -pyrone: kojic acid (Figure 5, Table 1). Paecilopyrone A and its linear analog paecilopyrone B may be derived from pentaketide; unfortunately, the

detailed synthetic pathway is still unclear [43]. Kojic acid showed antibacterial activities and tyrosinase inhibitory activity [44,45].




Figure 5. The structures of pyrones.

3.4. Phomaligols

Phomaligol A, phomaligol A1, methylphomaligol A, acetylphomaligol A, phomaligol A hydroperoxide, phomaligol A1 hydroperoxide, phomaligol B, and phomaligol C were all isolated from the culture of *P. lilacinum* (Figure 6, Table 1). All of them are yellow oil. Phomaligol A and Methylphomaligol A both showed cytotoxicity against a small panel of human solid tumor cell lines [46]. Phomaligol A, phomaligol A1, phomaligol A hydroperoxide, and phomaligol A1 hydroperoxide were previously isolated from *Phoma lingam* and *Phoma wasabiae*, but their biological activity has not been reported yet [47,48].




Figure 6. The structures of phomaligols.

3.5. Pigment

Lenta investigated the ethyl acetate extract of the mycelium of *P. lilacinum* isolated from the pigment purpureone (Figure 7, Table 1), which was found to possess potent antileishmanial activity and antibacterial activity [49].




Figure 7. The structure of purpureone.

4. Other Compounds

4.1. Ergosterols

Professor Cui and his team isolated seven compounds from the metabolites of *P. lilacinum* ZBY-1 identified as 9(11)-dehydroergosterol peroxide, ergosterol peroxide (Figure 8, Table 1), (22E,24R)-5 α ,6 α -epoxy-3 β -hydroxyergosta-22-ene-7-one, and cerebrosides A, B, C, and D. 9(11)-dehydroergosterol peroxide, ergosterol peroxide, and (22E,24R)-5 α ,6 α -epoxy-3 β -hydroxyergosta-22-ene-7-one had the inhibitory effect of human cancer K562, MCF-7, HL-60, and BGC-823 cells, with their IC₅₀ values on these cell lines ranging from 9.5 mg/L to 59.6 mg/L [50]. 9(11)-dehydroergosterol peroxide and ergosterol peroxide were found to be useful for developing a therapeutic candidate for lung cancer complications [51,52]. These three compounds all show high medical value and provide good materials for the research and development of anti-cancer drugs.




Figure 8. The structures of ergosterols.

4.2. Cerebrosides

Cerebrosides A, B, C, and D (Figure 9, Table 1) were isolated from the *P. lilacinum* ZBY-1 strain by professor Cui in 2013, and they are all linear analogs. Cerebrosides function as a non-race-specific elicitor in a wide range of plant-phytopathogenic fungus interactions [53]; cerebrosides A and B have significant effects of analgesia and brain protection [54]. Cerebrosides C and D were known to potentiate the activity of cell wall-active antibiotics [55]. In addition, cerebrosides A and C play a role in cell growth, differentiation, and apoptosis in animals [56].




Figure 9. The structures of cerebrosides.

4.3. Paecilaminols and Others

Cui found 11 compounds in the year of 2012 identified as paecilaminol, paecilaminol hydrochloride, 1(2)-linolyl-2(1)-palmityl-glycero-*O*-4'-(*N,N,N*-trimethyl) homoserine, 1,2-dilinolylglycero-*O*-4'-(*N,N,N*-trimethyl) homoserine, Me myristate, Me linoleate, linoleate, oleic acid, indole-3-carboxaldehyde, indolyl-3-carboxylic acid, and 4-hydroxybenzoic acid (Figure 10, Table 1). Compounds paecilaminol and paecilaminol hydrochloride have the ability to inhibit human cancer K562, MCF-7, HL-60, and BGC-823 cells. Compound paecilaminol was a major antitumor metabolite of the strain *P. lilacinum* ZBY-1 and it is also a NADH-fumarate reductase inhibitor [57].




Figure 10. Cont.




Figure 10. The structures of paecilaminols and others.

Me myristate was promised to be exploited as a hydrophobic medical carrier [58]. Me linoleate exhibited cytotoxic antibacterial activities against *Bacillus subtilis* and *Staphylococcus aureus* [59,60]. Modern medicine often makes acute respiratory distress syndrome models by injecting oleic acid into rabbits or rats [61].

Indole-3-carboxaldehyde has useful antimicrobial properties, and it can inhibit the atopic dermatitis-like inflammation induced by MC903 [62,63]. Hydroxybenzoic acids such as 4-hydroxybenzoic acid and 3,4-dihydroxybenzoic acid have various functional biological properties, including anticancer, antibacterial, antiaging, antidiabetic, anti-inflammatory, and antiviral activities. They are widely used in food, cosmetic, and pharmaceutical industries [64]. In addition, 4-hydroxybenzoic acid increased after an organic intervention diet, and it may help with health-promoting qualities in the near future [65,66].

Table 1. SMs isolated from *Purpureocillium lilacinum* and their biological activities.

Metabolites	CAS. No	Material Source	Biological Activity
Leucinostatin A	76600-38-9	<i>P. lilacinus</i> ZBY-1 from deep sea water	Inhibited prostate cancer cells [29], nematocidal activity [26], activity against Gram-positive bacteria [27].
Leucinostatin B	159544-15-7	Culture medium of <i>P. lilacinm</i>	Treatment of systemic candidiasis, nematocidal activity [26], activity against Gram-positive bacteria [27].
Leucinostatin C	110483-88-0	Culture medium of <i>P. lilacinm</i>	Drug-related side-effects and adverse reactions activity against Gram-positive bacteria [27], nematocidal activity [26].
Leucinostatin D	100334-47-2	Cultivated, mycelia complex of <i>P. marquandii</i>	Activity against Gram-positive bacteria [27], nematocidal activity [26].
Leucinostatin F		Culture medium of <i>P. lilacinm</i>	Unknown
Leucinostatin H	109539-58-4	Culture medium of <i>P. lilacinm</i>	Nematocidal activity [26].
Leucinostatin K	109539-57-3	Culture medium of <i>P. lilacinm</i>	Nematocidal activity [26].
Leucinostatin Y		Mycelia, cultivated complex of <i>P. linacinus</i> 40-H-28	Preferential cytotoxicity to cancer cells under glucose-deprived conditions and inhibition of mitochondrial function [32].

Table 1. Cont.

Metabolites	CAS. No	Material Source	Biological Activity
Acremoxanthone C	1360445-63-1P	Cultivated, mycelia complex of <i>P. lilacinm</i>	Cytotoxicity and 20 s proteasome inhibitory activity; high affinity with human calmodulin biosensors [37]; anti-oomycete activities [38]; exhibited anti- <i>Bacillus cereus</i> , antibacterial, antifungal, antiplasmodial, and cytotoxic activity; Gram-positive bacteria [36].
Acremoxanthone D	1360445-62-0P	Cultivated, mycelia complex of <i>P. lilacinm</i>	Moderate 20 s proteasome inhibitory activity [37].
Acremoxanthone F	1882150-25-5P	Cultivated, mycelia complex of <i>P. lilacinm</i>	Antimalarial activity against plasmodium falciparum K1 strain and multidrug-resistant strain [39].
Acremoxanthone G	1882150-26-6P	Cultivated, mycelia complex of <i>P. lilacinm</i>	Antimalarial activity against plasmodium falciparum K1 strain and multidrug-resistant strain [39].
Acremonidin A	701914-77-4P	Cultivated, mycelia complex of <i>P. lilacinm</i>	Moderate activity Against Gram-positive bacteria [36].
Acremonidin C	701914-79-6P	Cultivated, mycelia complex of <i>P. lilacinm</i>	Antibacterial activity [36].
Acremonidin G	1882150-23-3P	<i>P. lilacinus</i> ZBY-1 from deep sea water	Anti-enterococcus faecium activity [39].
Paecilomide	1538575-22-2P	Cultivated, mycelia complex of <i>P. lilacinm</i>	Acetylcholinesterase inhibitor [41].
9(11)-dehydroergosterolperoxide	91579717	<i>P. lilacinus</i> ZBY-1 from deep sea water	Cytotoxic effect [51].
Ergosterol peroxide	2061-64-5	<i>P. lilacinus</i> ZBY-1 from deep sea water	Exhibits antimycobacterial, trypanocidal, and antineoplastic activities [51].
(22E,24R)-5 α , 6 α -epoxy-3 β -hydroxyergosta-22-ene-7-one		<i>P. lilacinus</i> ZBY-1 from deep sea water	Inhibitory effect of human cancer K562, MCF-7, HL-60, and BGC-823 cells [50].
Cerebroside A	115681-40-8	<i>P. lilacinus</i> ZBY-1 from deep sea water	Induction of cell growth, differentiation, and apoptosis in animals [56].
Cerebroside B	88642-46-0	<i>P. lilacinus</i> ZBY-1 from deep sea water	Causes disease such as fusariosis, colitis, and apnea
Cerebroside C	98677-33-9	<i>P. lilacinus</i> ZBY-1 from deep sea water	Activity of cell wall-active; antibiotics; induction of cell growth, differentiation, and apoptosis in animals [55].
Cerebroside D	113773-89-0	<i>P. lilacinus</i> ZBY-1 from deep sea water	Activity of cell wall-active antibiotics [55].
Paecilopyrone A	1173292-70-0	Cultivated, mycelia complex of <i>P. lilacinm</i>	Unknown
Paecilopyrone B	1173292-71-1	Same as above	Unknown
Phomapyrone B	157744-25-7	Same as above	Unknown
Micropyrone	54682570	Same as above	Unknown
Phomapyrone C	157744-26-8	Same as above	Unknown
Kojic acid	501-30-4	Same as above	Antibacterial activities; tyrosinase inhibitory activity [44].
Phomaligol A	152204-32-5	Same as above	Unknown
Phomaligol A1	152053-11-7	Same as above	Unknown
Methylphomaligol A	152159-01-8	Same as above	Unknown

Table 1. Cont.

Metabolites	CAS. No	Material Source	Biological Activity
Acetylphomaligol A	1173292-72-2	Same as above	Unknown
Phomaligol A hydroperoxide	181798-75-4	Same as above	Unknown
Phomaligol A1 hydroperoxide	182072-72-6	Same as above	Unknown
Phomaligol B	1173292-73-3	Same as above	Unknown
Phomaligol C	1173292-74-4	Same as above	Unknown
Paecilaminol	540770-33-0	Same as above	Inhibits human cancer cell K562, MCF-7, HL-60, and BGC-823 cells [50].
Paecilaminol Hydrochloride	1650570-79-8	Same as above	Inhibits human cancer cell K562, MCF-7, HL-60, and BGC-823 cells
Me myristate	124-10-7	Same as above	Medical carrier [50].
Me linoleate	112-63-0	Same as above	Exhibited cytotoxic antibacterial activities against <i>Bacillus subtilis</i> and <i>Staphylococcus aureus</i> [59].
Indole-3-carboxaldehyde	487-89-8	Same as above	Antimicrobial properties [62].
Indolyl-3-carboxylic acid	771-50-6	Same as above	Potential in vitro antimalarial, anticancer activity [63].
4-hydroxybenzoic acid	99-96-7	Same as above	Inhibits LPS-induced protein [64].
Purpureone	2231079-10-8P	Mycelium of <i>P. lilacinm</i>	Antileishmanial activity; antibacterial activity [49].

5. Biosynthesis of Secondary Metabolites in *Purpureocillium lilacinum*

In 2015, Prasad sequenced the TERIBC-1 strain of *P. lilacinus* with a genome size of 40.02 Mb by using Illumina HiSeq technology, and predicted 30 secondary metabolite synthesis genes: 12 polyketide synthases (PKs, details of all abbreviations are in Table S1), 2 PKs-like, 7 nonribosome peptide synthetases, 7 NRPSs-like, 1 PK-NRPS, and 1 dimethylallyl tryptophan synthases (DMATs) gene [67]. In 2016, Wang sequenced *P. lilacinus* PLBJ-1 and PLFJ-1 strains. The genome sizes of the two strains were 38.14 Mb and 38.53 Mb, respectively [18]. Using SMURF [68] and anti-SMASH [69] software to predict the secondary metabolite synthesis gene cluster, PLBJ-1 and PLFJ-1 strains were found to encode 13 PKs, 2 PKs-like, 10 NRPSs, 10 NRPSs-like, 1 PK-NRPS, 4 terpene synthases (TSs), and 1 DMAT genes. It can be seen that the secondary metabolites produced by different species of *P. lilacinum* are not identical, but in general, *P. lilacinum* has great potential in the synthesis of secondary metabolites.

We know that the typical NRPS modules have adenylation (A), thiolation (T) or peptidyl carrier (PCP), condensation (C), and thioesterase (TE) domains [70,71], which are, respectively, responsible for the activation of amino acids, the extension of peptide chains, the formation of amide bonds, and the release of peptide chains [72,73]. The synthesis mechanism of NRPSs is shown in Figure 11. Generally speaking, the A domain combines with the amino acid substrate under the action of ATP to form the corresponding aminoacyl AMP, and the aminoacyl AMP combines with the sulfhydryl group of the T domain to form the aminoacyl-s-carrier complex. Finally, the carriers carrying the aminoacyl group and the peptide acyl group combine with the specific region of the C domain, and the amino group on the aminoacyl-s-carrier complex attacks the acyl group on the peptidyl-s-carrier complex, forming a new peptide bond, and finally forming a complete peptide chain, through the action of multiple modules, wherein the amino acids in the peptide chain correspond to the modules in the NRPS one by one. Some NRPS modules also contain epimerization (E), formylation (F), methylation (M), heterocyclization (CY) [74], reduction (R), and oxidation (OX) domains, which are involved in the structural modification of peptide chains. Finally, mature peptide chains are released from the NRP assembly line under the action of the TE domain [75].




Figure 11. The biosynthesis of NRPSs [76].

In 2016, Wang sequenced and analyzed the whole genome of *P. lilacinum*, and predicted the knock-out of the NRPS synthetic gene (LcsA), PK synthetase (LcsC), Acyl CoA ligase (LcsD), and thioesterase (LcsE), using high-performance liquid chromatography (HPLC) to compare the crude extracts of wild-type and mutant strains of *P. lilacinum*. It was found that the crude extracts of Δ LcsA, Δ LcsC, Δ LcsD, and Δ LcsE had a lack of Leucinostatin A and Leucinostatin B, and then these enzymes were found to play a key role in the synthesis of Leucinostatin, and the synthesis of leucinostatin of *P. lilacinum* was suggested. This hypothetical biosynthesis is initiated by the assembly of 4-methylhex-2-enoic acid via reductive PKs. However, they were unable to estimate which PKs were responsible for 4-methylhex-2-enoic acid [18].

In microorganisms, PK comes from the independent hypothesis of a variety of compounds. Polyketide compounds are assembled by repeated Claisen condensations between the activated acyl initiation unit and the chain extender unit derived from malonyl-CoA. This process is catalyzed by the synergistic action of keto synthase (KS), acyltransferase (AT), and phosphopantetheinyl acyl carrier protein (ACP) or CoA linked to the primary chain. After each extension step, the functionality of β -keto can be reduced by further involved enzymes [76]. This general PK catalytic mechanism is realized by different enzyme mechanisms (Figure 12). Three types of PKs are described below, which are responsible for the biosynthesis of polyketide chains.

PKs can be divided into three types: type I PKs are modular enzymes composed of several functional domains, which are arranged linearly and covalently. Any functional domain is not reused in the process of chain synthesis and extension. They mainly synthesize polyether, polyene, and macrolides. PKs of type II are aromatic, starting from acetyl CoA. Polyketones with an aromatic ring structure are synthesized with malonyl coenzyme A as an extension unit. Type III PKs are chalcone synthetases, a kind of homologous dimer enzyme that can be reused. It catalyzes the condensation of acetyl-CoA molecules to synthesize one ring or multi-ring aromatic polyketones [74].




Figure 12. The biosynthesis of PKs [77].

It is generally believed that most α -pyrones are synthesized through the polyketide pathway [78]. Terpenoids are a kind of chain or cyclic secondary metabolites, which are composed of isoprene as the basic unit. Terpenoids are synthesized by terpene synthase and can be divided into: monoterpenes, with geranyl diphosphate as the synthetic precursor; sesquiterpenes, with farnesyl diphosphate as the synthetic precursor; diterpenes, with geranyl pyrophosphate as the precursor. According to the degree of reduction, it can be divided into reduced terpenoids and nonreduced terpenoids [79].

6. Problems and Perspectives

Among the more than 40 metabolites reviewed in this paper, we can see that most SMs of *P. lilacinus* that have been reported so far have the functions of anticancer activity, antimicrobial activity, insecticidal activity, cytotoxicity, drug carriers, and so on. Most importantly, some of the compounds showed potent activities compared to those of the positive controls, which indicates that they could be used to develop new medicines. These include the anticancer lead compound leucinostatins, ergosterol peroxide, (22E,24R)-5 α , 6 α -epoxy-3 β -hydroxyergosta-22-ene-7-one, and paecilaminol. Leucinostatins is cytotoxic to HeLa cells, Ehrlich subcutaneous solid tumors, and prostate cancer. However, it was found to be toxic to rats by intraperitoneal injection, so more attention should be paid to its safety assessment when developing the drug. The other three compounds have the ability to inhibit human cancer K562, MCF-7, HL-60, and BGC-823 cells, but their safety for other species is still unknown. Acremoxanthone and acremonidin were both calmodulin inhibitors; paecilomide is an acetylcholinesterase inhibitor and kojic acid showed tyrosinase inhibitory activity, indicating their potential as insecticides. These remarkable activities make many of these compounds suitable candidates for new drugs and insecticides discovery and may lead to future synthesis studies. However, some of the SMs of *P. lilacinus* are toxic to animals and humanity. Hocquette, Dr. Qian, Pastor, and others have reported infections caused by *P. lilacinus* in immunocompromised patients [80,81].

With the development of society, more and more attention has been paid to biological control, more and more fungal products will come out, and the safety of related products has also received great attention. Therefore, how to ensure the safety of fungal products has become particularly important.

Generally, in production and in life, there are six destinations (i.e., target organisms, nontarget organisms, soil, water, atmosphere, and humans) involved in the production and application of *P. lilacinum* pesticide formulations. The most important destination is target organisms, including pests and crops when preparations are released in fields. Soil is another important destination, especially when it is released through soil treatments for nematodes. Water and the atmosphere are the destinations of the drifting formulations. Humans contact *P. lilacinum* through direct and indirect pathways. There is no doubt that

the biosafety risks of *P. lilacinum* are closely related to the sources and fates of the SMs produced by entomopathogenic fungi [24].

Surveying the SMs will be beneficial to improving the safety of *P. lilacinum* fungal products. Thus, developing the discovery, structure, function, and synthesis pathway of secondary metabolites of *P. lilacinum* are of great significance to biomedicine, human health, and agricultural disease control. For a long time, due to the gene silencing or low expression of most gene clusters in common culture medium, the research of fungal secondary metabolites has been hindered to some extent. There are only a few kinds of research on SMs of *P. lilacinum*, which are leucinostatins, acremoxanthones, and paecilomides, and their synthetic pathway and regulatory mechanism are still unclear. Therefore, it is necessary to use a super-expressing transcription factor, to replace the promoter in the synthetic gene cluster with an inducible strong promoter, to modify the histone, to heterologously express the gene cluster to activate the silent gene cluster, and to further discover that the structure is novel and biologically active. The SMs production yield of *P. lilacinum* needs to be improved by changing the culture conditions. First, gene knockout methods need to be used to further clarify the synthesis mechanism of secondary metabolites. In addition, it is necessary to continuously improve the efficiency and precision of chemical separation detection, in order to be more conducive to the separation of secondary metabolites and the identification of structural functions.

Supplementary Materials: The following are available online. Table S1: Information of all abbreviations.

Author Contributions: Conceptualization, W.C. and Q.H.; methodology, W.C.; investigation, W.C. and Q.H.; data curation, W.C. and Q.H.; writing—original draft preparation, W.C. and Q.H.; writing—review and editing, Q.H.; supervision, Q.H.; funding acquisition, Q.H. All authors have read and agreed to the published version of the manuscript.

Funding: This project was supported by the Guangzhou Science and Technology Project (201903010067).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Luangsa-Ard, J.; Houbraken, J.; van Doorn, T.; Hong, S.B.; Borman, A.M.; Hywel-Jones, N.L.; Samson, R.A. *Purpureocillium*, a new genus for the medically important *Paecilomyces lilacinus*. *FEMS Microbiol. Lett.* **2011**, *321*, 141–149. [[CrossRef](#)]
2. Sampson, R.A. *Paecilomyces* and Some Allied Hyphomycetes. *Cent. Voor Schimmelcultures* **1975**, *64*, 174. [[CrossRef](#)]
3. Srilakshmi, A.; Sai Gopal, D.V.R.; Narasimha, G. Impact of bioprocess parameters on cellulase production by *Purpureocillium lilacinum* isolated from forest soil. *Int. J. Pharma Bio Sci.* **2017**, *8*, 157–165. [[CrossRef](#)]
4. Zhu, Y.; Ai, D.; Zhang, W. Difference of soil microbiota in perennial ryegrass turf before and after turning green using high-throughput sequencing technology. *Res. J. BioTechnol.* **2017**, *12*, 50–60.
5. Redou, V.; Navarri, M.; Meslet-Cladiere, L.; Barbier, G.; Burgaud, G. Species richness and adaptation of marine fungi from deep-subseafloor sediments. *Appl. Environ. Microbiol.* **2015**, *81*, 3571–3583. [[CrossRef](#)] [[PubMed](#)]
6. Liu, L.; Zhang, C.; Fan, H.; Guo, Z.; Yang, H.; Chen, M.; Han, J.; Cao, Y.; Xu, J.; Zhang, K.; et al. An efficient gene disruption system for the nematophagous fungus *Purpureocillium lavendulum*. *Fungal. Biol.* **2019**, *123*, 274–282. [[CrossRef](#)] [[PubMed](#)]
7. Silva, S.D.; Carneiro, R.M.D.G.; Faria, M.; Souza, D.A.; Monnerat, R.G.; Lopes, R.B. Evaluation of *Pochonia chlamydosporia* and *Purpureocillium lilacinum* for suppression of *Meloidogyne enterolobii* on tomato and banana. *J. Nematol.* **2017**, *49*, 77–85. [[CrossRef](#)] [[PubMed](#)]
8. Gine, A.; Sorribas, F.J. Effect of plant resistance and BioAct WG (*Purpureocillium lilacinum* strain 251) on *Meloidogyne incognita* in a tomato-cucumber rotation in a greenhouse. *Pest Manag. Sci.* **2017**, *73*, 880–887. [[CrossRef](#)]
9. Cavello, I.A.; Hours, R.A.; Rojas, N.L.; Cavalitto, S.F. Purification and characterization of a keratinolytic serine protease from *Purpureocillium lilacinum* LPS # 876. *Process. Biochem.* **2013**, *48*, 972–978. [[CrossRef](#)]
10. Desaegeer, J.A.; Watson, T.T. Evaluation of new chemical and biological nematicides for managing *Meloidogyne javanica* in tomato production and associated double-crops in Florida. *Pest Manag. Sci.* **2019**, *75*, 3363–3370. [[CrossRef](#)] [[PubMed](#)]

11. Jiao, Y.; Li, Y.; Li, Y.; Cao, H.; Mao, Z.; Ling, J.; Yang, Y.; Xie, B. Functional genetic analysis of the leucinostatin biosynthesis transcription regulator *lcsL* in *Purpureocillium lilacinum* using CRISPR-Cas9 technology. *Appl. Microbiol. Biotechnol.* **2019**, *103*, 6187–6194. [[CrossRef](#)]
12. Wang, M.; Zhou, H.; Fu, Y.; Wang, C. The Antifungal Activities of the Fungus 36–1 to Several Plant Pathogens. *Chin. J. Biol. Control.* **1996**, *12*, 20–23.
13. Li, F.; Chen, J.; Shi, H.; Liu, B. Antagonistic effect of biocontrol fungus, *Paecilomyces lilacinus* strain NH-PL-3 and its mechanism against *Fusarium oxysporum*. *J. Plant Prot.* **2005**, *32*, 373–378.
14. Hotaka, D.; Amnuaykanjanasin, A.; Maketon, C.; Siritutsoontorn, S.; Maketon, M. Efficacy of *Purpureocillium lilacinum* CKPL-053 in controlling *Thrips palmi* (Thysanoptera: Thripidae) in orchid farms in Thailand. *Appl. Entomol. Zool.* **2015**, *50*, 317–329. [[CrossRef](#)]
15. Yoder, J.A.; Fisher, K.A.; Dobrotka, C.J. A report on *Purpureocillium lilacinum* found naturally infecting the predatory mite, *Balaustium murorum* (Parasitengona: Erythraeidae). *Int. J. Acarol.* **2018**, *44*, 139–145. [[CrossRef](#)]
16. Deng, J.X.; Paul, N.C.; Sang, H.K.; Lee, J.H.; Hwang, Y.S.; Yu, S.H. First Report on Isolation of *Penicillium adametzoides* and *Purpureocillium lilacinum* from Decayed Fruit of Cheongsoo Grapes in Korea. *Mycobiology* **2012**, *40*, 66–70. [[CrossRef](#)] [[PubMed](#)]
17. Guo, L.-N.; Wang, H.; Hsueh, P.-R.; Meis, J.F.; Chen, H.; Xu, Y.-C. Endophthalmitis caused by *Purpureocillium lilacinum*. *J. Microbiol. Immunol. Infect.* **2019**, *52*, 170–171. [[CrossRef](#)] [[PubMed](#)]
18. Wang, G.; Liu, Z.; Lin, R.; Li, E.; Mao, Z.; Ling, J.; Yang, Y.; Yin, W.-B.; Xie, B. Biosynthesis of antibiotic leucinostatins in bio-control fungus *Purpureocillium lilacinum* and their inhibition on *Phytophthora* revealed by genome mining. *PLoS Pathog.* **2016**, *12*, e1005685. [[CrossRef](#)] [[PubMed](#)]
19. Bode, H.B.; Bethe, B.; Hofs, R.; Zeeck, A. Big effects from small changes: Possible ways to explore nature's chemical diversity. *Chembiochem* **2002**, *3*, 619–627. [[CrossRef](#)]
20. Liu, L.; Zhang, J.; Chen, C.; Teng, J.; Wang, C.; Luo, D. Structure and biosynthesis of fumosorinone, a new protein tyrosine phosphatase 1B inhibitor firstly isolated from the entomogenous fungus *Isaria fumosorosea*. *Fungal Genet. Biol.* **2015**, *81*, 191–200. [[CrossRef](#)]
21. Yurchenko, A.N.; Girich, E.V.; Yurchenko, E.A. Metabolites of Marine Sediment-Derived Fungi: Actual Trends of Biological Activity Studies. *Mar. Drugs* **2021**, *19*, 88. [[CrossRef](#)] [[PubMed](#)]
22. Li, X.-Q.; Xu, K.; Liu, X.-M.; Zhang, P. A Systematic Review on Secondary Metabolites of *Paecilomyces* Species: Chemical Diversity and Biological Activity. *Planta Medica* **2020**, *86*, 805–821. [[CrossRef](#)] [[PubMed](#)]
23. Wen, L.; Lin, Y.C.; She, Z.G.; Du, D.S.; Chan, W.L.; Zheng, Z.H. Paeciloxanthone, a new cytotoxic xanthone from the marine mangrove fungus *Paecilomyces* sp. (Tree1–7). *J. Asian Nat. Prod. Res.* **2008**, *10*, 133–137. [[CrossRef](#)] [[PubMed](#)]
24. Weng, Q.; Zhang, X.; Chen, W.; Hu, Q. Secondary Metabolites and the Risks of *Isaria fumosorosea* and *Isaria farinosa*. *Molecules* **2019**, *24*, 664. [[CrossRef](#)]
25. Zhang, X.; Hu, Q.; Weng, Q. Secondary metabolites (SMs) of *Isaria cicadae* and *Isaria tenuipes*. *RSC Adv.* **2019**, *9*, 172–184. [[CrossRef](#)]
26. Mikami, Y.; Yazawa, K.; Fukushima, K.; Arai, T.; Samson, R.A.J.M. Paecilotoxin production in clinical or terrestrial isolates of *Paecilomyces lilacinus* strains. *Mycopathologia* **1989**, *108*, 195–199. [[CrossRef](#)] [[PubMed](#)]
27. Rossi, C.; Tuttobello, L.; Ricci, M.; Casinovi, C.G.; Radios, L. Leucinostatin D, a novel peptide antibiotic from *Paecilomyces marquandii*. *J. Antibiot.* **1987**, *40*, 130–133. [[CrossRef](#)]
28. Momose, I.; Onodera, T.; Doi, H.; Adachi, H.; Iijima, M.; Yamazaki, Y.; Sawa, R.; Kubota, Y.; Igarashi, M.; Kawada, M. Leucinostatin Y: A Peptaibiotic Produced by the Entomoparasitic Fungus *Purpureocillium lilacinum* 40-H-28. *J. Nat. Prod.* **2019**, *82*, 1120–1127. [[CrossRef](#)]
29. Kawada, M.; Inoue, H.; Momose, I.; Masuda, T.; Ikeda, D. 188 POSTER Leucinostatins suppress prostate cancer cell growth through the tumour-stromal cell interactions. *Eur. J. Cancer Suppl.* **2008**, *6*, 59–60. [[CrossRef](#)]
30. Kawada, M.; Inoue, H.; Ohba, S.I.; Masuda, T.; Momose, I.; Ikeda, D. Leucinostatin A inhibits prostate cancer growth through reduction of insulin-like growth factor-I expression in prostate stromal cells. *Int. J. Cancer* **2010**, *126*, 810–818. [[CrossRef](#)]
31. Ricci, M.; Blasi, P.; Giovagnoli, S.; Perioli, L.; Vescovi, C.; Rossi, C. Leucinostatin-A loaded nanospheres: Characterization and in vivo toxicity and efficacy evaluation. *Int. J. Pharm.* **2004**, *275*, 61–72. [[CrossRef](#)] [[PubMed](#)]
32. Arai, T.; Mikami, Y.; Fukushima, K.; Utsumi, T.; Yazawa, K. A new antibiotic, leucinostatin, derived from *Penicillium lilacinum*. *J. Antibiot.* **1973**, *26*, 157–161. [[CrossRef](#)]
33. Fukushima, K.; Arai, T.; Mori, Y.; Tsuboi, M.; Suzuki, M. Studies on peptide antibiotics, leucinostatins. I. Separation, physico-chemical properties and biological activities of leucinostatins A and B. *J. Antibiot.* **1983**, *36*, 1606–1612. [[CrossRef](#)] [[PubMed](#)]
34. Ishiyama, A.; Otoguro, K.; Iwatsuki, M.; Namatame, M.; Nishihara, A.; Nonaka, K.; Kinoshita, Y.; Takahashi, Y.; Masuma, R.; Shiomi, K.; et al. In vitro and in vivo antitrypanosomal activities of three peptide antibiotics: Leucinostatin A and B, alamethicin I and tsushimycin. *J. Antibiot.* **2009**, *62*, 303–308. [[CrossRef](#)] [[PubMed](#)]
35. Madariaga-Mazón, A.; González-Andrade, M.; González, M.d.C.; Glenn, A.E.; Cerda-García-Rojas, C.M.; Mata, R. Absolute Configuration of Acremoxanthone C, a Potent Calmodulin Inhibitor from *Purpureocillium lilacinum*. *J. Nat. Prod.* **2013**, *76*, 1454–1460. [[CrossRef](#)] [[PubMed](#)]
36. Ayers, S.; Graf, T.N.; Adcock, A.F.; Kroll, D.J.; Shen, Q.; Swanson, S.M.; Matthew, S.; de Blanco, E.; Wani, M.C.; Darveaux, B.A.; et al. Cytotoxic xanthone-anthraquinone heterodimers from an unidentified fungus of the order Hypocreales (MSX 17022). *J. Antibiot.* **2012**, *65*, 3–8. [[CrossRef](#)]

37. Melendez-Gonzalez, C.; Muria-Gonzalez, M.J.; Anaya, A.L.; Hernandez-Bautista, B.E.; Hernandez-Ortega, S.; Gonzalez, M.C.; Glenn, A.E.; Hanlin, R.T.; Macias-Rubalcava, M.L. Acremoxanthone E, a Novel Member of Heterodimeric Polyketides with a Bicyclo[3.2.2]nonene Ring, Produced by *Acremonium camptosporum* W. Gams (Clavicipitaceae) Endophytic Fungus. *Chem. Biodivers.* **2015**, *12*, 133–147. [CrossRef]
38. Isaka, M.; Palasarn, S.; Auncharoen, P.; Komwijit, S.; Gareth Jones, E.B. Acremoxanthones A and B, novel antibiotic polyketides from the fungus *Acremonium* sp. BCC 31806. *Tetrahedron Lett.* **2009**, *50*, 284–287. [CrossRef]
39. He, H.; Bigelis, R.; Solum, E.H.; Greenstein, M.; Carter, G.T. Acremonidins, new polyketide-derived antibiotics produced by *Acremonium* sp., LL-Cyan 416. *J. Antibiot.* **2003**, *56*, 923–930. [CrossRef]
40. Teles, A.P.C.; Takahashi, J.A. Paecilomide, a new acetylcholinesterase inhibitor from *Paecilomyces lilacinus*. *Microbiol. Res.* **2013**, *168*, 204–210. [CrossRef] [PubMed]
41. Tumiatto, V.; Bolognesi, M.L.; Minarini, A.; Rosini, M.; Milelli, A.; Matera, R.; Melchiorre, C. Progress in acetylcholinesterase inhibitors for Alzheimer's disease: An update. *Expert Opin. Ther. Pat.* **2008**, *18*, 387–401. [CrossRef]
42. Kulshreshtha, A.; Piplani, P. Current pharmacotherapy and putative disease-modifying therapy for Alzheimer's disease. *Neurol. Sci.* **2016**, *37*, 1403–1435. [CrossRef]
43. Seibert, S.F.; Eguereva, E.; Krick, A.; Kehraus, S.; Voloshina, E.; Raabe, G.; Fleischhauer, J.; Leistner, E.; Wiese, M.; Prinz, H.; et al. Polyketides from the marine-derived fungus *Ascochyta salicorniae* and their potential to inhibit protein phosphatases. *Org. Biomol. Chem.* **2006**, *4*, 2233–2240. [CrossRef] [PubMed]
44. Song, L.; Xie, W.; Zhao, Y.; Lv, X.; Yang, H.; Zeng, Q.; Zheng, Z.; Yang, X. Synthesis, antimicrobial, moisture absorption and retention activities of kojic acid-grafted konjac glucomannan oligosaccharides. *Polymers* **2019**, *11*, 1979. [CrossRef]
45. Li, X.; Jeong, J.H.; Lee, K.T.; Rho, J.R.; Choi, H.D.; Kang, J.S.; Son, B.W. γ -pyrone derivatives, kojic acid methyl ethers from a marine-derived fungus *Altenaria* sp. *Arch. Pharmacol. Res.* **2003**, *26*, 532–534. [CrossRef]
46. Elbandy, M.; Shinde, P.B.; Hong, J.; Bae, K.S.; Kim, M.A.; Lee, S.M.; Jung, J.H. α -pyrones and yellow pigments from the sponge-derived fungus *Paecilomyces lilacinus*. *Bull. Korean Chem. Soc.* **2009**, *30*, 188–192. [CrossRef]
47. Pedras, M.S.C.; Taylor, J.L.; Morales, V.M. Phomaligin A and other yellow pigments in *Phoma lingam* and *P. wasabiae*. *Phytochemistry* **1995**, *38*, 1215–1222. [CrossRef]
48. Soga, O.; Iwamoto, H.; Hata, K.; Maeba, R.; Takuwa, A.; Fujiwara, T.; Hsu, Y.H.; Nakayama, M. New oxidation product of was abidienone-A. *Agric. Biol. Chem.* **1988**, *52*, 865–866. [CrossRef]
49. Lenta, B.N.; Ngatchou, J.; Frese, M.; Ladoh-Yemeda, F.; Voundi, S.; Nardella, F.; Michalek, C.; Wibberg, D.; Ngouela, S.; Tsamo, E.; et al. Purpureone, an antileishmanial ergochrome from the endophytic fungus *Purpureocillium lilacinum*. *Z. Für Nat. B J. Chem. Sci.* **2016**, *71*, 1159–1167. [CrossRef]
50. Cui, X.; Li, C.; Wu, C.; Hua, W.; Cui, C.; Zhu, T.; Gu, Q.-Q. Metabolites of *Paecilomyces lilacinus* zby-1 from deep-sea water and their antitumor activity. *J. Int. Pharm. Res.* **2013**, *40*, 177–186.
51. Wu, H.; Yang, F.; Li, L.; Rao, Y.K.; Ju, T.; Wong, W.; Hsieh, C.; Pivkin, M.V.; Hua, K.; Wu, S. Ergosterol peroxide from marine fungus *Phoma* sp. induces ROS-dependent apoptosis and autophagy in human lung adenocarcinoma cells. *Sci. Rep.* **2018**, *8*, 1–14. [CrossRef] [PubMed]
52. Lee, S.; Lee, S.; Roh, H.-S.; Song, S.-S.; Ryoo, R.; Pang, C.; Baek, K.-H.; Kim, K.H. Cytotoxic constituents from the sclerotia of *Poria cocos* against human lung adenocarcinoma cells by inducing mitochondrial apoptosis. *Cells* **2018**, *7*, 116. [CrossRef] [PubMed]
53. Umemura, K.; Tanino, S.; Nagatsuka, T.; Koga, J.; Iwata, M.; Nagashima, K.; Amemiya, Y. Cerebroside elicitor confers resistance to Fusarium disease in various plant species. *Phytopathology* **2004**, *94*, 813–818. [CrossRef] [PubMed]
54. Hua, W.; Miao, Y.; Chen, J.; You, Y.; Sun, C.; Yuan, L.; Xu, L. Identification and growth characterization of a cerebroside producing *Termitomyces clypeatus* CTM-1. *Shengwu Jiagong Guocheng* **2015**, *13*, 67–73. [CrossRef]
55. Wicklow, D.T.; Joshi, B.K.; Gamble, W.R.; Gloer, J.B.; Dowd, P.F. Antifungal metabolites (monorden, monocillin IV, and cerebroside) from *Humicola fuscoatra* traena NRRL 22980, a mycoparasite of *Aspergillus flavus* sclerotia. *Appl. Environ. Microbiol.* **1998**, *64*, 4482–4484. [CrossRef] [PubMed]
56. Koga, J. Induction of Rice Disease Resistance by Fungal Sphingolipids and Bile Acids. *Am. Chem. Soc.* **2009**, *237*, 100.
57. Ui, H.; Shiomi, K.; Suzuki, H.; Hatano, H.; Morimoto, H.; Yamaguchi, Y.; Masuma, R.; Sakamoto, K.; Kita, K.; Miyoshi, H.; et al. Paecilaminol, a new NADH-fumarate reductase inhibitor, produced by *Paecilomyces* sp. FKI-0550. *J. Antibiot.* **2006**, *59*, 591–596. [CrossRef] [PubMed]
58. Hu, Y.; Guo, J.; Liu, W. The synthesis and property research of nanoparticles based on hydrophobic modified inulin. *Shandong Huagong* **2015**, *44*, 10–15.
59. Khentoul, H.; Bensouici, C.; Reyes, F.; Albanese, D.; Sarri, D.; Ratiba, M.; Fadila, B.; Seghiri, R.; Boumaza, O. Chemical constituents and HRESI-MS analysis of an Algerian endemic plant—*Verbascum atlanticum* Batt—extracts and their antioxidant activity. *Nat. Prod. Res.* **2019**, *34*, 3008–3012. [CrossRef]
60. Abd-Ellatif, A.E.S.; Abdel-Razek, A.S.; Hamed, A.; Soltan, M.M.; Soliman, H.S.M.; Shaaban, M. Bioactive compounds from marine *Streptomyces* sp.: Structure identification and biological activities. *Vietnam. J. Chem.* **2019**, *57*, 628–635. [CrossRef]
61. Goncalves-de-Albuquerque, C.F.; Silva, A.R.; Burth, P.; Castro-Faria, M.V.; Castro-Faria-Neto, H.C. Acute Respiratory Distress Syndrome: Role of Oleic Acid-Triggered Lung Injury and Inflammation. *Mediat. Inflamm.* **2015**, *2015*, 1–9. [CrossRef]
62. Balaz, M.; Kudlickova, Z.; Vilkova, M.; Imrich, J.; Balazova, L.; Daneu, N. Mechanochemical synthesis and isomerization of N-substituted indole-3-carboxaldehyde oximes. *Molecules* **2019**, *24*, 3347. [CrossRef] [PubMed]

63. Li, W.; Yu, J.; Yao, X. Application of Indole-3-Carboxaldehyde in Preparation of Cosmetic or Drug for Treating Atopic Dermatitis. CN110368385A, 25 October 2019. China National Intellectual Property Administration, Beijing, China.
64. Kim, H.; Kim, S.Y.; Sim, G.Y.; Ahn, J.-H. Synthesis of 4-Hydroxybenzoic Acid Derivatives in *Escherichia coli*. *J. Agric. Food Chem.* **2020**, *68*, 9743–9749. [[CrossRef](#)]
65. Hurtado-Barroso, S.; Quifer-Rada, P.; Marhuenda-Munoz, M.; de Alvarenga, J.F.R.; Tresserra-Rimbau, A.; Lamuela-Raventos, R.M. Increase of 4-hydroxybenzoic, a bioactive phenolic compound, after an organic intervention diet. *Antioxidants* **2019**, *8*, 340. [[CrossRef](#)] [[PubMed](#)]
66. Maia, N.J.L.; Correa, J.A.F.; Rigotti, R.T.; da Silva Junior, A.A.; Luciano, F.B. Combination of natural antimicrobials for contamination control in ethanol production. *World J. Microbiol. Biotechnol.* **2019**, *35*, 1–9. [[CrossRef](#)] [[PubMed](#)]
67. Prasad, P.; Varshney, D.; Adholeya, A. Whole genome annotation and comparative genomic analyses of bio-control fungus *Purpureocillium lilacinum*. *BMC Genom.* **2015**, *16*, 1004. [[CrossRef](#)]
68. Khaldi, N.; Seifuddin, F.T.; Turner, G.; Haft, D.; Nierman, W.C.; Wolfe, K.H.; Fedorova, N.D. SMURF: Genomic mapping of fungal secondary metabolite clusters. *Fungal Genet. Biol.* **2010**, *47*, 736–741. [[CrossRef](#)] [[PubMed](#)]
69. Weber, T.; Blin, K.; Duddela, S.; Krug, D.; Kim, H.U.; Brucoleri, R.; Lee, S.Y.; Fischbach, M.A.; Muller, R.; Wohlleben, W.; et al. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res.* **2015**, *43*, W237–W243. [[CrossRef](#)]
70. Takeda, K.; Kemmoku, K.; Satoh, Y.; Ogasawara, Y.; Shinya, K.; Dairi, T.J.A.C.B. N-Phenylacetylation and Nonribosomal Peptide Synthetases with Substrate Promiscuity for Biosynthesis of Heptapeptide Variants, JBIR-78 and JBIR-95. *ACS Chem. Biol.* **2017**, *12*, 1813. [[CrossRef](#)]
71. Han, M.; Chen, J.; Qiao, Y.; Zhu, P. Advances in the nonribosomal peptide synthetases. *Yaoxue Xuebao* **2018**, *53*, 1080–1089. [[CrossRef](#)]
72. Sung, C.T.; Chang, S.; Entwistle, R.; Ahn, G.; Lin, T.; Petrova, V.; Yeh, H.; Praseuth, M.B.; Chiang, Y.; Oakley, B.R.; et al. Overexpression of a three-gene conidial pigment biosynthetic pathway in *Aspergillus nidulans* reveals the first NRPS known to acetylate tryptophan. *Fungal Genet. Biol.* **2017**, *101*, 1–6. [[CrossRef](#)] [[PubMed](#)]
73. Payne, J.A.E.; Schoppet, M.; Hansen, M.H.; Cryle, M.J. Diversity of nature’s assembly lines—recent discoveries in non-ribosomal peptide synthesis. *Mol. Biosyst.* **2017**, *13*, 9–22. [[CrossRef](#)]
74. Sun, Y.-H.; Deng, Z.-X. Polyketides and combinatorial biosynthetic approaches. *Zhongguo Kangshengsu Zazhi* **2006**, *31*, 6.
75. Bloudoff, K.; Fage, C.D.; Marahiel, M.A.; Schmeing, T.M. Structural and mutational analysis of the nonribosomal peptide synthetase heterocyclization domain provides insight into catalysis. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 95–100. [[CrossRef](#)] [[PubMed](#)]
76. Strieker, M.; Tanovic, A.; Marahiel, M.A. Nonribosomal peptide synthetases: Structures and dynamics. *Curr. Opin. Struct. Biol.* **2010**, *20*, 234–240. [[CrossRef](#)] [[PubMed](#)]
77. Hertweck, C. The Biosynthetic Logic of Polyketide Diversity. *Angew. Chem. Int. Ed.* **2009**, *48*, 4688–4716. [[CrossRef](#)] [[PubMed](#)]
78. Schaeberle, T.F. Biosynthesis of alpha-pyrone. *Beilstein J. Org. Chem.* **2016**, *12*, 571–588. [[CrossRef](#)]
79. Sun, J.; Bu, J.; Cui, G.; Ma, Y.; Zhao, H.; Mao, Y.; Zeng, W.; Guo, J.; Huang, L. Accumulation and biosynthetic of curcuminoids and terpenoids in turmeric rhizome in different development periods. *Zhongguo Zhong Yao Za Zhi* **2019**, *44*, 927–934. [[CrossRef](#)] [[PubMed](#)]
80. Hocquette, A.; Grondin, M.; Bertout, S.; Mallié, M. Les champignons des genres *Acremonium*, *Beauveria*, *Chrysosporium*, *Fusarium*, *Onychocola*, *Paecilomyces*, *Penicillium*, *Scedosporium* et *Scopulariopsis* responsables de hyalohyphomycoses. *J. De Mycol. Médicale* **2005**, *15*, 136–149. [[CrossRef](#)]
81. Okhravi, N.; Lightman, S. Clinical manifestations, treatment and outcome of *Paecilomyces lilacinus* infections. *Clin. Microbiol. Infect.* **2007**, *13*, 554. [[CrossRef](#)] [[PubMed](#)]