


## Protocol

# Purification and cryo-EM structure determination of *Arabidopsis thaliana* GLR3.4



Ionotropic glutamate receptors (iGluRs) are ligand-gated ion channels that play crucial roles in the central nervous system. iGluR homologs, termed glutamate receptor-like channels (GLRs), have been found in plants. Investigating the structural and functional relationship between iGluRs and GLRs was limited by GLR protein expression, purification, and structural characterization. Here, we provide a detailed protocol for *Arabidopsis thaliana* GLR3.4 (AtGLR3.4) expression in a mammalian cell line and purification for structure determination by cryogenic electron microscopy (cryo-EM).

Shanti Pal Gangwar,  
Marriah N. Green,  
Maria V.  
Yelshanskaya,  
Alexander I.  
Sobolevsky

as4005@cumc.columbia.edu

### Highlights

Protocol for expression of plant glutamate receptor-like channel (GLR) in HEK cells

Purification protocol that yields pure monodisperse tetrameric GLR protein

Purified protein subjected to cryo-EM analysis produces high-resolution GLR structure

Gangwar et al., STAR  
Protocols 2, 100855  
December 17, 2021 © 2021  
The Author(s).  
<https://doi.org/10.1016/j.xpro.2021.100855>



## Protocol

Purification and cryo-EM structure determination of *Arabidopsis thaliana* GLR3.4Shanti Pal Gangwar,<sup>1,3</sup> Marriah N. Green,<sup>1,2,3</sup> Maria V. Yelshanskaya,<sup>1</sup> and Alexander I. Sobolevsky<sup>1,4,5,\*</sup><sup>1</sup>Department of Biochemistry and Molecular Biophysics, Columbia University, 650 West 168<sup>th</sup> Street, New York, NY 10032, USA<sup>2</sup>Institute of Human Nutrition, Columbia University, 630 West 168<sup>th</sup> Street, New York, NY 10032, USA<sup>3</sup>These authors contributed equally<sup>4</sup>Technical contact<sup>5</sup>Lead contact\*Correspondence: [as4005@cumc.columbia.edu](mailto:as4005@cumc.columbia.edu)  
<https://doi.org/10.1016/j.xpro.2021.100855>

## SUMMARY

Ionotropic glutamate receptors (iGluRs) are ligand-gated ion channels that play crucial roles in the central nervous system. iGluR homologs, termed glutamate receptor-like channels (GLRs), have been found in plants. Investigating the structural and functional relationship between iGluRs and GLRs was limited by GLR protein expression, purification, and structural characterization. Here, we provide a detailed protocol for *Arabidopsis thaliana* GLR3.4 (AtGLR3.4) expression in a mammalian cell line and purification for structure determination by cryogenic electron microscopy (cryo-EM).

For the complete details on the use and execution of this protocol, please refer to Green et al. (2021).

## BEFORE YOU BEGIN

This protocol was used in a recent publication (Green et al., 2021) to purify *Arabidopsis thaliana* GLR3.4 (AtGLR3.4) recombinant protein expressed in a mammalian cell line for high-resolution full-length structure determination by single-particle cryo-EM. Before the experiment, prepare all the buffers to be used in subsequent steps one day in advance unless otherwise mentioned.

## Buffer preparation

⌚ Timing: 1 day


Prepare 1L of T buffer, 100 mL of Cell Lysis buffer, 25 mL of Solubilization buffer, 300 mL of Size-Exclusion Chromatography (SEC) buffer and 20 mL of Strep Elution buffer.

## Construct preparation

⌚ Timing: 4 days

1. Subclone DNA for full-length AtGLR3.4 (Met1-Thr959) into a pEG BacMam vector (Goehring et al., 2014) using gene-specific primers. Introduce the thrombin cleavage site (LVPRGS), followed by an eGFP and a streptavidin (strep) affinity tag (WSHPQFEK) at the C-terminus (Figure 1).
2. Design the gene-specific forward primer with the Sall and reverse primer with the NotI site to amplify the AtGLR3.4 DNA flanked by Kozak sequence (KZK) and thrombin site to clone in pEG BacMam. Conduct polymerase chain reaction (PCR) using the information described in the tables below.





**Figure 1. AtGLR3.4 construct schematic**

▮▮▮ **Pause point:** PCR reaction can be stored at 4°C for 24 h or at –20°C for long-term storage.

3. Run a 1% agarose gel to verify the size and purity the PCR products.
4. Add 1  $\mu$ L of DpnI enzyme and incubate the reaction at 37°C for 1 h.
5. Perform DNA cleanup using any commercially available kit. We use Monarch New England Bio Labs' PCR and DNA Cleanup Kit following manufactures protocol.

▮▮▮ **Pause point:** DNA can be stored at 4°C for 24 h or at –20°C for long-term storage.

6. Digest PCR DNA and pEG BacMam plasmid in two separate 1.5-mL tubes following the instruction in the table below. Incubate the digest reaction at 37°C for 1 h.
7. Perform a DNA cleanup using any commercially available kit as mentioned above.
8. Ligate both the PCR and linearized plasmid following the instruction in the table below. Incubate the ligation reaction at ~20°C–25°C for 10–15 min. Use the insert and vector DNA at 3:1 molar ratio.
9. Transform DH5 $\alpha$  competent cells and select the transformants on LB agar plates containing 100  $\mu$ g/mL ampicillin.
  - a. Incubate the plates at 37°C for 12–15 h.
  - b. Inoculate a single colony in LB media supplemented with 100  $\mu$ g/mL ampicillin and perform a miniprep to extract and purify the plasmid DNA.
  - c. Sequence the DNA to confirm that the construct sequence is correct before proceeding.

**Note:** The circular plasmid that we used was engineered to have a Kozak sequence, thrombin recognition site, eGFP, and Strep-tag as described in [Figure 1](#).

## KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
Tris-HCl	Fisher Scientific	Cat# BP152-1
NaCl	Fisher Scientific	Cat# BP358-212
L-Glutamate	Sigma	Cat# 49621
PMSF	Acros Organics	Cat# 215740500
2-Mercaptoethanol ( $\beta$ ME)	Acros Organics	Cat# 125470100
Thrombin	Haematologic Technologies	Cat# HCT-0020
D-Desthiobiotin	Sigma-Aldrich	Cat#D1411
Fetal bovine serum	Gibco	Cat# 16140071
Sf-900 III SFM	Gibco	Cat# 12658027
Freestyle 293 expression medium	Gibco	Cat# 12338018
Sodium butyrate	Acros Organics	Cat# 263191000
Digitonin	Cayman Chemical Company	Cat#14952
Aprotinin	Sigma-Aldrich	Cat#A1153
Leupeptin	Sigma-Aldrich	Cat#L0649
Pepstatin A	Sigma-Aldrich	Cat#P4265
Kanamycin	Fisher Scientific	Cat# BP906-5

(Continued on next page)

<b>Continued</b>		
REAGENT or RESOURCE	SOURCE	IDENTIFIER
Gentamicin sulfate	Sigma-Aldrich	Cat#G1914
Tetracycline	Fisher Scientific	Cat# BP912
Bluo-Gal	Life Technologies	Cat# 15519028
IPTG	Zymo Research	Cat# I1001-5
Sall-HF	New England Biolabs	Cat# R3138S
NotI-HF	New England Biolabs	Cat#3189S
Quick ligase	New England Biolabs	Cat#M2200
Monarch Plasmid Miniprep Kit	New England Biolabs	Cat #T1010L
Monarch DNA and PCR Cleanup Kit	New England Biolabs	Cat #T1030L
Nitrocellulose membrane filter (0.22 μm)	Merck Millipore	Cat#SA1J789H5
Wizard Plus Minipreps DNA Purification System	Promega	Cat#A7510
Phenol:chloroform:isoamylalcohol	Life Technologies	Cat# 15593031
Cellfectin II Reagent	Gibco	Cat#58760
Ethanol 200 Proof	Decon Labs, Inc.	Cat #2701
Centrifugal Filter Unit (Amicon Ultra-15)	Sigma-Aldrich	Cat#UFC910024
<b>Deposited data</b>		
Coordinates of full-length AtGLR3.4	(Green et al., 2021)	PDB: 7LZH
Cryo-EM map of full-length AtGLR3.4	(Green et al., 2021)	EMDB: EMD-23606
Coordinates of AtGLR3.4-S1S2 <sub>Glu</sub>	(Green et al., 2021)	PDB: 7LZO
Crystal structure of GluA2	(Sobolevsky et al., 2009)	PDB: 3KG2
<b>Experimental models: cell lines</b>		
HEK 293S GnTI <sup>-</sup>	ATCC	Cat#CRL-3022
Sf9	Gibco	Cat#12659017
DH10Bac	Life Technologies	Cat#10361012
DH5α	Zymo Research	Cat#T3007
<b>Oligonucleotides</b>		
AtGLR3.4 amplification primer: 5'-gtcgactccgccaccatgggatttttggtgatgataagag -3'	This paper	N/A
AtGLR3.4 amplification primer: 5'-cggcaccagagtaatttcgcatgtgtgattgtga -3'	This paper	N/A
<b>Software and algorithms</b>		
gCTF	(Zhang, 2016)	<a href="http://www.mrc-lmb.cam.ac.uk/kzhang/">http://www.mrc-lmb.cam.ac.uk/kzhang/</a>
Motioncor2	(Zheng et al., 2017)	<a href="https://msg.ucsf.edu/software">https://msg.ucsf.edu/software</a>
RELION 3.0	(Zivanov et al., 2018)	<a href="http://www2.mrc-lmb.cam.ac.uk/relion/">http://www2.mrc-lmb.cam.ac.uk/relion/</a>
UCSF Chimera	(Pettersen et al., 2004)	<a href="https://www.cgl.ucsf.edu/chimera/">https://www.cgl.ucsf.edu/chimera/</a>
cryoSPARC	(Punjani et al., 2017)	<a href="https://cryosparc.com/">https://cryosparc.com/</a>
PHENIX	(Adams et al., 2010)	<a href="https://www.phenix-online.org/">https://www.phenix-online.org/</a>
Coot	(Emsley and Cowtan, 2004)	<a href="http://www2.mrc-lmb.cam.ac.uk/Personal/pemsley/coot">http://www2.mrc-lmb.cam.ac.uk/Personal/pemsley/coot</a>
SWISS-MODEL	(Waterhouse et al., 2018)	<a href="https://swissmodel.expasy.org/">https://swissmodel.expasy.org/</a>
<b>Other</b>		
CF-1.2/1.3-2Au 200 mesh holey carbon grids	Protochips	Cat#CF-1.2/1.3-2Au
Gold wire	Ted Pella, Inc.	Cat#21-10
Superose6 10/300 column	GE Healthcare	Cat# 17-5172-01

## MATERIALS AND EQUIPMENT

<b>T buffer</b>		
Reagent	Final concentration	Amount
Tris-HCl pH 8.0 (1 M)	20 mM	20 mL
NaCl (5 M)	150 mM	30 mL

(Continued on next page)

**Continued**

Reagent	Final concentration	Amount
Milli-Q H <sub>2</sub> O	n/a	950 mL
<b>Total</b>	<b>n/a</b>	<b>1 L</b>

Run T buffer through a 0.22 µm filter and store at 4°C for up to one month.

**Cell Lysis buffer**

Reagent	Final concentration	Amount
T buffer	n/a	100 mL
β-mercaptoethanol (βME) (14.3 M)	1 mM	7.1 µL
Aprotinin (0.8 mM)	0.8 µM	100 µL
Leupeptin (4.3 mM)	4.3 µM	100 µL
Pepstatin A (2 mM)	2 µM	100 µL
Phenylmethylsulphonyl fluoride (1 M)	1 mM	100 µL
<b>Total</b>	<b>n/a</b>	<b>~100 mL</b>

**Note:** Protease inhibitors and βME should be added right before buffer use. Keep at 4°C, if needed can be stored at 4°C for up to 2 hours.

△ **CRITICAL:** Protein degradation during the purification procedure can have detrimental effects on the resulting protein quality. It is, therefore, important that fresh reducing agents and protease inhibitors are added at the steps indicated in the protocol.

**Solubilization buffer**

Reagent	Final concentration	Amount
T buffer	n/a	25 mL
β-mercaptoethanol (βME) (14.3 M)	1 mM	1.78 µL
Aprotinin (0.8 mM)	0.8 µM	25 µL
Leupeptin (4.3 mM)	4.3 µM	25 µL
Pepstatin A (2 mM)	2 µM	25 µL
Phenylmethylsulphonyl fluoride (1 M)	1 mM	25 µL
Digitonin	2%	0.5 g
<b>Total</b>	<b>n/a</b>	<b>~25 mL</b>

**Note:** Digitonin is not readily soluble at room temperature. Therefore, add digitonin to the T buffer, heat the mixture up to 70°C–75°C and stir at medium-low speed until completely dissolved. Keep at 4°C, if needed can be stored at 4°C for up to 2 hours.

△ **CRITICAL:** Protease inhibitors and βME should be added to the chilled digitonin plus T buffer mixture right before membrane solubilization.

**SEC buffer**

Reagent	Final concentration	Amount
T buffer	n/a	300 mL
Digitonin	0.05%	150 mg
<b>Total</b>	<b>n/a</b>	<b>300 mL</b>

**Note:** Digitonin is not readily soluble at room temperature. Therefore, add digitonin to the T buffer, heat the mixture up to 70°C–75°C and stir at medium-low speed until completely dissolved. After digitonin is completely dissolved, cool to ~4°C–20°C. Once

cooled, run SEC buffer through 0.22  $\mu\text{M}$  filter. Keep at 4°C, if needed can be stored at 4°C for up to 24 hours.

### Strep Elution buffer

Reagent	Final concentration	Amount
SEC buffer	n/a	20 mL
D-desthiobiotin	2.5 mM	10.8 mg
<b>Total</b>	<b>n/a</b>	<b>20 mL</b>

If needed, adjust the pH of Strep Elution buffer to 8.0, run it through 0.22  $\mu\text{M}$  filter and store at 4°C for up to 12 h.

### PCR reaction

Reagents	Final concentration	Volume
Autoclaved Milli-Q H <sub>2</sub> O	n/a	32 $\mu\text{L}$
5 $\times$ Q5 Reaction Buffer	1 $\times$	10 $\mu\text{L}$
Forward Primer (10 $\mu\text{M}$ )	0.5 $\mu\text{M}$	2.5 $\mu\text{L}$
Reverse Primer (10 $\mu\text{M}$ )	0.5 $\mu\text{M}$	2.5 $\mu\text{L}$
Template DNA	1–10 ng	1 $\mu\text{L}$
dNTPs (10 mM)	200 $\mu\text{M}$	1 $\mu\text{L}$
Q5 High-Fidelity DNA Polymerase	0.04 U/ $\mu\text{L}$	1 $\mu\text{L}$
<b>Total</b>	<b>n/a</b>	<b>50 <math>\mu\text{L}</math></b>

### Thermocycler settings included lid temperature at 105°C and volume of 50 $\mu\text{L}$ .

#### PCR cycling conditions

Steps	Temperature	Time	Number of cycles
Initial Denaturation	98°C	3 min	1
Denaturation	98°C	10 s	25–35 cycles
Annealing	55°C	30 s	
Extension	72°C	2 min	
Final extension	72°C	5 min	1
Hold	4°C	Forever	

### Digest reaction

Reagents	Volume
PCR DNA/Plasmid	80 $\mu\text{L}$
CutSmart Buffer 10 $\times$	10 $\mu\text{L}$
Sall-HF	2 $\mu\text{L}$
NotI-HF	2 $\mu\text{L}$
Milli-Q H <sub>2</sub> O	6 $\mu\text{L}$
<b>Total</b>	<b>100 <math>\mu\text{L}</math></b>

### Ligation reaction

Reagents	Volume
PCR DNA	40–50 ng
Plasmid DNA	40–60 ng
Quick ligase buffer (2 $\times$ )	5 $\mu\text{L}$
Quick ligase	0.5 $\mu\text{L}$
Milli-Q H <sub>2</sub> O	—
<b>Total</b>	<b>10 <math>\mu\text{L}</math></b>

## STEP-BY-STEP METHOD DETAILS


Below we provide a detailed step-by-step protocol about the AtGLR3.4 protein purification, sample preparation, and cryo-EM structure determination (Figure 2).

### Bacmid preparation and isolation

⌚ Timing: 4 days

According to the manufacturer protocol, bacmid preparation and isolation were carried out using DH10Bac competent cells (Bac-to-Bac, Invitrogen). The steps are described here.

1. For transposition, prepare LB agar plates containing 50  $\mu\text{g}/\text{mL}$  kanamycin, 7  $\mu\text{g}/\text{mL}$  gentamicin, 10  $\mu\text{g}/\text{mL}$  tetracycline, 200  $\mu\text{g}/\text{mL}$  Bluo-gal and 40  $\mu\text{g}/\text{mL}$  IPTG.
2. Transformation of bacmid
  - a. Thaw DH10Bac (Life Technologies Cat#10361012) competent cells on ice and dispense 25–30  $\mu\text{L}$  to a prechilled 1.5-mL polypropylene tube.
  - b. Add approximately 1–10 ng (1–2  $\mu\text{L}$ ) of plasmid DNA (pEG BacMam\_AtGLR3.4) and mix gently by flicking/tapping the tube several times.



**Figure 2. Overview of the protocol of AtGLR3.4 protein expression, purification, sample preparation, and cryo-EM structure determination**  
The image has been created using BioRender (<https://biorender.com/>).

- c. Incubate on ice for 15–30 min.
  - d. Transfer the tube to a water bath at 42°C for 45 s.
  - e. After heat shock, immediately transfer the tube back to ice and let it cool on ice for 2–3 min.
  - f. Add 200  $\mu$ L SOC media to the tube and transfer to an incubator shaker set at 220 rpm and incubate for 4 h at 37°C.
  - g. Pipette 20–100  $\mu$ L of transformation reaction on the plates and spread evenly.
3. Grow transformed cells
- a. Incubate the plates for at least 24 h at 37°C. Blue and white colonies will appear; for better results, incubate the plates for 48 h to distinguish the blue from the white colonies.
  - b. Carefully inoculate a single white colony in 5–6 mL LB broth supplemented with 50  $\mu$ g/mL kanamycin, 7  $\mu$ g/mL gentamicin, and 10  $\mu$ g/mL tetracycline.
  - c. Incubate for 12–15 h at 37°C with continuous rotation at 220–250 rpm.

▮▮ **Pause point:** Overnight (12–16 h)

4. Isolate bacmid DNA
- a. Centrifuge the culture at  $\sim$ 3,100 g for 10 min. Then, decant to discard the supernatant and save the cell pellet.

**Note:** If needed, the pellet can be stored at  $-20^{\circ}\text{C}$  for up to a couple of weeks prior to bacmid DNA isolation. For bacmid isolation, we use reagents from the Promega mini prep kit. Reagents from other mini prep kits can also be used.

- b. Resuspend the cell pellet in 200  $\mu$ L resuspension buffer until homogeneous, via pipetting up and down and/or by vortexing. Transfer the suspension to a 1.5-mL Eppendorf tube.
- c. Add 200  $\mu$ L lysis buffer and mix by gently inverting the tube several ( $\sim$ 15–20) times. The suspension in the tube turns slightly viscous.
- d. Add 200  $\mu$ L neutralization buffer and mix by gentle inversion several ( $\sim$ 20) times. A white precipitate will be observed.
- e. Centrifuge the tube at  $\sim$ 21,100 g for 10 min at  $\sim$ 20°C–25°C.
- f. Avoiding the pellet, carefully collect and transfer the supernatant to a fresh 1.5-mL Eppendorf tube. In the fume hood, add an equal volume (typically around 500–600  $\mu$ L) of Tris equilibrated Phenol:Chloroform: Isoamyl-alcohol (25:24:1) and mix by inverting gently several times for 1 min.
- g. Spin at  $\sim$ 21,100 g for 5 min at  $\sim$ 20°C–25°C. In the fume hood, gently collect the top layer supernatant ( $\sim$ 500–600  $\mu$ L) and transfer it to a new 1.5-mL Eppendorf tube. Add 2 $\times$  volume ( $\sim$ 1–1.2 mL) of ice-cold 100% ethanol, mix well by gently inverting for 30 s, and incubate at  $-20^{\circ}\text{C}$  for a minimum of 30 min.
- h. Centrifuge at  $\sim$ 21,100 g for 20–30 min at 4°C. A white pellet is observed at the bottom of the tube. Carefully decant to discard the supernatant and add 1 mL ice-cooled 70% ethanol.
- i. Invert the tube gently several times to wash the pellet. Centrifuge for 5 min at  $\sim$ 21,100 g at 4°C.
- j. Discard the supernatant by gentle aspiration (make sure to not discard the pellet as it may dislodge from the bottom).
- k. To avoid traces of remaining ethanol, air dry the pellet for 15–20 min at  $\sim$ 20°C–25°C in a fume hood. Add 40–50  $\mu$ L of autoclaved milli-Q water and dissolve the pellet by gentle tapping/flicking. Once the pellet is dissolved, place the tube with bacmid DNA on ice. Now the bacmid DNA is ready to be used for baculovirus production and can be stored at  $-80^{\circ}\text{C}$  for an extended period.

### Transfection of Sf9 cells with recombinant bacmid DNA

⌚ **Timing:** 4–6 days

Production of the first-generation baculovirus (P1 virus).



**Note:** The following steps should be performed under sterile conditions in a laminar hood.

5. In a 6-well plate, add 2 mL of Sf9 cells per well, with the Sf9 cell density of  $0.5\text{--}0.75 \times 10^6/\text{mL}$  ( $1\text{--}1.5 \times 10^6$  cells per P1 virus). Allow the cells to adhere for 30 min to 1 h in a dark incubator at 27°C.
6. In the meantime, take a 1.5-mL Eppendorf tube and add:
  - a. 180  $\mu\text{L}$  of SF900 media
  - b. 10  $\mu\text{L}$  of Cellfectin II
  - c. 10  $\mu\text{L}$  of thawed bacmid DNA
7. Close and tap the tube to mix. Let mixture sit for 20–30 min in the hood.
8. Add the mixture of SF900 media, bacmid DNA, and Cellfectin II dropwise to the cells in the well.
9. Incubate the plate at 27°C for 4–6 days. Observe the plate under the microscope to monitor cell behavior and fluorescence if the gene of interest is fused with a fluorescent marker gene.

**Note:** It is always better to have a fluorescent marker fused to the protein of interest to monitor baculovirus preparation and protein expression. The AtGLR3.4 P1 virus is typically harvested on the 5<sup>th</sup> day post-transfection because most of the cells already express the protein (monitored by eGFP fluorescence), while very little cell death is observed.

10. To harvest the P1 virus, filter the 2 mL medium using a 0.2  $\mu\text{m}$  sterile syringe filter in a laminar hood and store the filtered virus at 4°C protected from light. The P1 virus can be stored for 4–6 months.

## P2 virus production

⌚ Timing: 5 days

**Note:** The following steps (except for centrifugation and resuspension) should be performed under sterile conditions in a laminar hood.

11. In a laminar hood and sterile conditions, infect 500 mL of Sf9 cells at a cell density of  $1.5 \times 10^6$  cells/mL in a 1-L non-baffled Erlenmeyer flask by adding 500  $\mu\text{L}$  of P1 virus. Incubate the Sf9 cell culture in a shaker at 27°C and 115 rpm for 72–120 h. Keep the cell culture protected from light.
12. Centrifuge the culture at  $\sim 5,000 g$  for 15 min at 4°C in a Sorvall centrifuge. Discard the cell pellet and save the supernatant containing the P2 virus.
13. Ultracentrifuge the supernatant at 4°C for 1 h at  $\sim 61,000 g$  in a preparative ultracentrifuge with a Type 45Ti Beckman Coulter rotor.
14. A small translucent pellet is observed at the bottom of the tube. Discard the supernatant and resuspend the pellet in 50 or 25 mL Gibco FreeStyle 293 expression media supplemented with 2% of  $\gamma$ -irradiated fetal bovine serum (FBS) to get 10 $\times$  or 20 $\times$  concentrated P2 virus, respectively.

**Note:** Always run the FBS through a 0.22  $\mu\text{m}$  filter before adding to the media (perform this procedure under sterile conditions in a laminar hood).

15. Pass the resuspension through a 0.22  $\mu\text{m}$  filter into a sterile 50-mL tube and store at 4°C well protected from light. The P2 virus stock can be stored for 1–2 months at 4°C while protected from light.

**Note:** To fully dissolve the pellet, allow it to stay in the resuspension media in the dark at 4°C for no longer than 24 hours prior to filtration.

## Transduction of HEK 293S GnTI<sup>-</sup> cells with P2 baculovirus

⌚ Timing: 3 days

16. In a laminar hood and sterile conditions, prepare 800 mL of HEK 293S GnT1<sup>-</sup> cell culture at the density of  $2.5\text{--}3.5 \times 10^6$  cells/mL maintained in Freestyle 293 media supplemented with 2% FBS in 2-L baffled Erlenmeyer flask and add 8 mL of 20 $\times$  concentrated P2 virus stock. Incubate the cell culture at 37°C in an orbital shaker at 110 rpm and 5% CO<sub>2</sub>.
17. To enhance the protein expression, add 10 mM of sodium butyrate to the culture 12–20 h post-transduction and decrease the incubator temperature to 30°C.
18. After 72–74 h post-transduction, harvest the cells by centrifugation at  $\sim 5,000$  g (in a Sorvall RC 5B Plus centrifuge) for 15 min at 4°C. Discard the supernatant, wash the cell pellet by resuspension in phosphate-buffered saline (PBS, pH 8.0), and transfer the suspension to a 50-mL Falcon tube. Centrifuge at  $\sim 3,100$  g for 10 min at 4°C. Discard the supernatant and store the cell pellet at  $-80^\circ\text{C}$  until further use.

### Purification of AtGLR3.4

⌚ Timing: 2 days

**Note:** The buffers should be filtered using 0.22  $\mu\text{m}$  filter (Merck-Millipore) and chilled to 4°C prior to utilization. When required, add  $\beta$ -mercaptoethanol ( $\beta$ ME) and protease inhibitors right before buffer usage. All protein purification steps are to be carried out at 4°C or on ice, unless otherwise noted.

19. Resuspend the cell pellet in ice-cold Cell Lysis buffer. Add a stir bar and  $\sim 40$  mL of the Cell Lysis buffer to the 50-mL falcon tube with the frozen cell pellet till it reaches 45-mL volume and resuspend the pellet by rocking on a platform or by vortexing until the cell pellet becomes detached and dissolved.
20. Disrupt the cells by sonication using Misonix Sonicator with a preset program with six cycles (3 min total process time) at the amplitude of 8, 15 s pulse on time, and 15 s pulse off time. The sonication steps must be carried out on a stir plate (at medium speed stirring) on ice or at 4°C to avoid heating and denaturing of the protein. Repeat this process 2 more times or until optimal cell lysis.

**Note:** For optimal cell lysis, check a small droplet of the lysate under a light microscope to confirm that cells are lysed post-sonication, as the appearance will be distinctly different from cells prior to sonication.

21. Centrifuge the cell lysate suspension at  $\sim 3,100$  g (using an Eppendorf Centrifuge 5810) for 10 min at 4°C to remove cell debris and unbroken cells.
22. Collect the clarified supernatant by decanting the supernatant into a prechilled polycarbonate bottle assembly for ultracentrifugation. Ultracentrifuge in a Type 45Ti Beckman Coulter fixed-angle rotor at  $\sim 186,000$  g for 1 h at 4°C to obtain the membrane fraction.
23. Discard the supernatant and mechanically homogenize the obtained membrane fraction pellet in Cell Lysis buffer ( $\sim 25$  mL) using a homogenizer.

**Note:** We use a prechilled dounce homogenizer and move a plunger forth and back at least 6 times to fully homogenize the pellet in the Cell Lysis buffer.

24. Solubilize the membrane protein from the homogenate by adding an equal volume of solubilization buffer to the homogenate (totaling  $\sim 50$  mL with a final  $\sim 1\%$  concentration of the detergent, digitonin in our case) and stir/rotate (low-medium speed to prevent bubble formation) at 4°C for  $\sim 2$  h (1 h minimum).
25. Ultracentrifuge the solubilize at  $\sim 186,000$  g in a Type 45Ti Beckman Coulter fixed-angle rotor at 4°C for 1 h to remove insoluble material.

26. Collect the supernatant and add 2 mL of strep resin prewashed and equilibrated in T buffer. Rotate the mix for 12–14 h at 4°C.

▮▮ **Pause point:** overnight (12–15 h)

27. Purify the protein using affinity chromatography.
- Equilibrate 30-mL chromatography gravity column by rinsing column with milli-Q water and allowing SEC buffer to flow through.
  - Transfer the AtGLR3.4 protein-bound strep resin to the empty pre-equilibrated gravity chromatography column and collect the flow-through.
  - Wash the resin by pouring 25 mL of SEC buffer into the column.
  - Elute the AtGLR3.4 protein with ~10–15 mL of freshly made Strep Elution buffer. Observe the eGFP-tagged protein eluting from the column by the color of the collected protein appearing green and the resin becoming white. No more Strep Elution buffer is needed once the strep resin is white, stripped of its eGFP green color, indicating that the protein has been eluted from the column.

**Note:** When adding buffer to the column, wait until the buffer from the previous step is almost finished flowing through the column but do not let the column run dry.

28. Measure the concentration of the protein, add 1/300 (w/w) thrombin and incubate at 22°C for 90 min to cleave off eGFP and the strep tag. Protein concentration can be measured using a spectrophotometer set to  $A_{280}$  (absorbance at 280 nm) and blanked with the elution buffer.

**Note:** Successful thrombin cleavage can be monitored by fluorescence-detection size-exclusion chromatography (FSEC) (Kawate and Gouaux, 2006) and SDS-PAGE (Figure 3).

29. Concentrate the thrombin digest reaction to ~500  $\mu$ L using 100-kDa NMWL centrifugal filter.
30. Transfer the concentrated protein to a new 1.5-mL tube and centrifuge the concentrated protein at ~86,500 g for 30 min at 4°C using a Sorvall MTX150 Micro-Ultracentrifuge (Thermo Fisher Scientific) and a S100AT4 rotor.
31. Avoiding the pellet, inject the supernatant into a 500- $\mu$ L loop connected to a Superose™ 6 10/300 GL SEC column attached to an AKTA FPLC (GE Healthcare) and pre-equilibrated in SEC Buffer.
32. At the end of the SEC column run at 0.5 mL/min, pool the peak fractions corresponding to AtGLR3.4 tetramer (Figure 4) and concentrate to 3–4 mg/mL using 100-kDa NMWL centrifugal filter. Typically, ~800 mL of HEK 293S GnTI<sup>-</sup> cell culture yields ~1 mg of purified protein. The concentrated, purified protein can be stored at 4°C, structurally stable, and functionally active for one week.
33. Inspect the purity of the protein by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE).


**Optional:** Protein purity can also be assessed by FSEC.

### Grid sample preparation

⌚ **Timing:** 2–3 h

34. For cryo-EM grid sample preparation, use UltrAuFoil CF-1.2/1.3-2Au 200 mesh holey carbon grids covered with a thin layer of gold according to the published method (Russo and Passmore, 2014).

**Optional:** The user can optimize various grid types and vitrification parameters however they see fit.



**Figure 3. On the left, FSEC traces for AtGLR3.4 before (blue) and after (red) thrombin cleavage**

The chromatograms were recorded using Superose 6 column at the flow rate of 0.5 mL/min using eGFP fluorescence (excitation, 488 nm; emission, 507 nm). On the right, SDS-PAGE demonstrating AtGLR3.4 bands before and after thrombin digest at two (1/1,000 and 1/300) thrombin/AtGLR3.4-eGFP ratios.


- a. Deposit ~50 nm of gold onto the grids using gold wire and the Edwards Auto 306 evaporator.

**Note:** This grid gold-coating step is only necessary for carbon-coated grids.

- b. Then remove the carbon with the Gatan Solarus (model 950) Advanced Plasma Cleaning System Ar/O<sub>2</sub> treatment (4 min, 50 watts, 35.0 sccm Ar, 11.5 sccm O<sub>2</sub>).

**Note:** This carbon removal step is only necessary after the previous grid gold-coating step.

- c. Before applying the purified protein sample to the gold-coated side of the grid, subject the grids to a H<sub>2</sub>/O<sub>2</sub> plasma treatment using the Gatan Solarus (model 950) Advanced Plasma Cleaning System (20 s, 10 watts, 6.4 sccm H<sub>2</sub>, 27.5 sccm O<sub>2</sub>) with the gold-coated side facing up.



**Figure 4. On the left, SEC profile for purified AtGLR3.4**

The chromatogram was recorded using Superose 6 column at the flow rate of 0.5 mL/min using A<sub>280</sub> absorbance. In the middle, SDS-PAGE for the protein peak fractions outlined by the red dashed lines on the SEC plot. On the right, normalized FSEC traces for purified AtGLR3.4, rat GluA2-γ2 fusion (Twomey et al., 2016, 2017), mouse TRPV3 (Nadezhdin et al., 2021; Singh et al., 2018) and human TRPV6 (Bhardwaj et al., 2020; McGoldrick et al., 2018). The latter three membrane proteins represent molecular weight markers and confirm the tetrameric assembly of AtGLR3.4. The chromatograms were recorded using Superose 6 column at the flow rate of 0.5 mL/min using tryptophan fluorescence (excitation, 280 nm; emission, 334 nm).

**Note:** Glow discharging step could be used as an alternative to plasma cleaning.

35. Prior to applying the purified protein to the grid, ultracentrifuge the protein at  $\sim 86,500$  g for 30 min at  $4^{\circ}\text{C}$  using a SorvallMTX150 Micro-Ultracentrifuge (Thermo Fisher Scientific) and a S100AT4 rotor. Transfer the supernatant to a prechilled tube, avoiding any possible protein precipitation.
36. For vitrification, set the vitrobot to  $4^{\circ}\text{C}$  with 100% humidity, a 15 s wait time, and a 4–5 s blot time with a 4–5 blot force. Apply  $3\ \mu\text{L}$  of purified AtGLR3.4 protein at 3–3.5 mg/mL to the gold-coated side of the grid and plunge freeze the grid into liquid ethane cooled to  $\sim -190^{\circ}\text{C}$  in liquid nitrogen using a Mark IV vitrobot (Thermo fisher scientific).

**Note:** After grids are plunge frozen, they must remain at cryogenic temperatures and avoid contamination as well as exposure to humidity or condensation.

**▮▮ Pause point:** The prepared grids can stay in liquid nitrogen for long-term storage.

### Grid screening, cryo-EM data collection, and processing

**⌚ Timing:** 2–4 weeks

37. Test the grids on screening transmission electron microscopes (TEMs), such as Glacios or F20 (FEI Thermo Scientific), and assess particle distribution, orientation, and ice quality.

**Note:** The grids can be loaded onto the F20 microscope straight from the grid storage box. However, the grids need to be clipped before being screened on the Glacios microscope.

38. Collect cryo-EM data from the prescreened grids on a FEI Titan Krios TEM (Thermo Fisher Scientific) operating at 300 kV and equipped with a direct electron detection (DED) camera in counting mode with  $\sim 50$  frames per movie and a total dose of  $\sim 58$  electrons per  $\text{\AA}^2$ , a physical pixel size of  $\sim 0.83\ \text{\AA}$ , and a defocus value between  $-0.5$  to  $-2.5\ \mu\text{m}$ . We used Krios with a post-column GIF Quantum energy filter and a Gatan K3 Summit DED camera, Gatan, Pleasanton, CA, USA.
39. Process the data in RELION 3.1 (Zivanov et al., 2018) or CryoSparc (Punjani et al., 2017). Perform beam-induced motion correction using MotionCor2 (Dose per frame of 1.16, EER fractionation -32, B Factor -150, number of patches  $5 \times 5$ ) (Zheng et al., 2017) and contrast transfer function (CTF) estimation using CTFFIND4.1 (Zhang, 2016) in RELION 3.1.


**Note:** All the processing steps can also be performed in cryoSPARC.

40. Manually pick  $\sim 3,000$  particles by selecting any particle shape that commonly appears in the micrographs, recognizing reoccurring shapes as well as any shape that potentially resembles an iGluR.

**Note:** We used the particle/mask diameter of  $230\ \text{\AA}$ .

41. 2D-classify these manually selected particles to generate templates for further template-based auto picking in RELION 3.1.
42.  $4 \times 4$  bin the picked particle images to a pixel size of  $\sim 3.32\ \text{\AA}/\text{pixel}$  and subject them to 3D classification into 10 classes (C1 symmetry). Select the best classes, which corresponded to the Y shape of iGluR (Figure 5).

**Note:** A model-generated iGluR map (for example, using the GluA2 model, PDB: 3KG2 (Sobolevsky et al., 2009)), low-pass filtered to  $40\ \text{\AA}$ , can be used as an initial reference.



**Figure 5. Cryo-EM processing workflow**

43. Extract particles for the best 3D classes without binning to an original pixel size ( $\sim 0.83$  Å/pixel).

Perform 3D refinement using C2 symmetry in RELION 3.1. At this step, the postprocessed map from RELION had the resolution of  $\sim 3.98$  Å (FSC = 0.143) from 277,615 particles (Figure 5).

44. Carry out micelle subtraction. For micelle subtraction, first create a mask using 3D refined map as input from the last 3D refinement step in Relion. Then run Particle subtraction using optimizer.star file from the last 3D refinement step and the mask as an input. Carry out multiple rounds of 3D classification (C1 symmetry) without angular sampling to reduce particle heterogeneity followed by 3D refinement. Also, perform Bayesian polishing and CTF refinement.

**Note:** Vary the regularization parameter (T) and iterations for 3D classification. We used T= 4 and iterations = 40 in 3D classification without angular sampling. Multiple rounds of 3D classification into 8 classes followed by 3D refinement on the particles from the best class using C2 symmetry and postprocessing yielded a map with resolution of 3.73Å from 118,592 particles (FSC=0.143) in RELION. Every time we used the 3D refined map from the previous job as an input reference in the subsequent 3D classification job.

45. To improve the map quality, import the particles from the last 3D refinement job from RELION to cryoSPARC and clean them up by 2D classification.

46. Select the best 2D classes and subject the corresponding particles to homogeneous and non-uniform refinement in cryoSPARC applying C2 symmetry. The map after non-uniform refinement in cryoSPARC had a resolution of 3.57 Å (FSC = 0.143) (Chen et al., 2013) from 110,630 particles.

47. Estimate the local resolution using the unfiltered half-maps and Resmap (Kucukelbir et al., 2014) (Figure 5).

**Note:** Local resolution can also be estimated in cryoSPARC.

48. Use UCSF Chimera ([Pettersen et al., 2004](#)) to visualize the EM density maps.

### Model building and refinement

⌚ Timing: 1–2 weeks

Once the high-resolution cryo-EM map is obtained, build atomic models of AtGLR3.4.

49. As guides, use the crystal structure of AtGLR3.4-S1S2 (PDB: 7LZ0) ([Green et al., 2021](#)) to build the ligand-binding domain and homology modeling in SWISS-MODEL ([Waterhouse et al., 2018](#)) using the GluA2 crystal structure as a guide (PDB: 3KG2, ([Sobolevsky et al., 2009](#))) to build the amino terminal and transmembrane domains. Finalize the model building manually using COOT ([Emsley and Cowtan, 2004](#)).

**Note:** Since AtGLR3.4 is a tetramer, it is advisable to first build two subunits, A and B, and then duplicate the AB dimer following the C2 symmetry of the map to make the CD dimer and assemble the ABCD tetramer.

50. Refine the obtained atomic model of AtGLR3.4 in real-space using PHENIX ([Adams et al., 2010](#)).  
51. Validate the model quality. For this purpose, we used the validation programs MolProbity (score = 1.69, clash score = 4.49, poor rotamers = 0.44%) ([Williams et al., 2018](#)) and EMRinger (EMRinger score = 2.41, optimal Threshold = 0.67, Rotamer-ratio = 0.77) ([Barad et al., 2015](#)) in Phenix.

### EXPECTED OUTCOMES

Approximately ~ 1 mg of purified AtGLR3.4 can be obtained from 800 mL HEK cells.

### LIMITATIONS

The multidomain architecture and flexibility of AtGLR3.4 could limit the resolution.

### TROUBLESHOOTING

#### Problem 1

No PCR fragment amplified or incorrect size of the PCR product (step 3 of the Construct preparation).

#### Potential solution

Optimize the primer annealing temperature by running a gradient PCR and optimize the primer extension time.

#### Problem 2

Few or no transformants (step 9 of the Construct preparation and step 3 of the main protocol).

#### Potential solution

Check the efficiency of the competent cells by transforming a control circular plasmid. Also check the temperature of the water bath for heat shock.

#### Problem 3

For different membrane proteins, one may need to use different detergents for solubilization (step 24 of the main protocol).

#### Potential solution

Check which detergent best extracts the protein of interest by running a detergent screen via FSEC on crude cell samples solubilized in different detergents.

### Problem 4

Inefficient thrombin cleavage and eGFP removal (step 28 of the main protocol).

### Potential solution

Optimize the concentration of the protein and thrombin amount in the reaction mixture. Sometimes, extremely high concentration of the protein results in incomplete thrombin digestion.

### Problem 5

Too high or too low particle density on the grid (step 37 of the main protocol).

### Potential solution

Adjust the protein concentration accordingly or vary the blot time and blot force when using the Vitrobot.

## RESOURCE AVAILABILITY

### Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Dr. Alexander I. Sobolevsky ([as4005@cumc.columbia.edu](mailto:as4005@cumc.columbia.edu)).

### Materials availability

This study did not generate any new unique reagents. Further information requests about materials and reagents should be directed to the lead contact, Dr. Alexander I. Sobolevsky ([as4005@cumc.columbia.edu](mailto:as4005@cumc.columbia.edu)).

### Data and code availability

No new code has been generated during this study. The EMDB and PDB codes generated by this data have already been deposited and reported in our previous paper ([Green et al., 2021](#)).

## ACKNOWLEDGMENTS

We are thankful to Kirill Nadezhdin and Arthur Neuberger for providing MW membrane protein standards. A.I.S. was supported by the National Institutes of Health (CA206573, NS083660, and NS107253) and the National Science Foundation (1818086).

## AUTHOR CONTRIBUTIONS

S.P.G., M.N.G., and M.V.Y. performed the experiments. M.N.G., S.P.G., and A.I.S. wrote the manuscript.

## DECLARATION OF INTERESTS

The authors declare no competing interests.

## REFERENCES

- Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr. D Biol. Crystallogr.* *66*, 213–221.
- Barad, B.A., Echols, N., Wang, R.Y., Cheng, Y., DiMaio, F., Adams, P.D., and Fraser, J.S. (2015). EMRinger: side chain-directed model and map validation for 3D cryo-electron microscopy. *Nat. Methods* *12*, 943–946.
- Bhardwaj, R., Lindinger, S., Neuberger, A., Nadezhdin, K.D., Singh, A.K., Cunha, M.R., Derler, I., Gyimesi, G., Reymond, J.L., Hediger, M.A., et al. (2020). Inactivation-mimicking block of the epithelial calcium channel TRPV6. *Sci. Adv.* *6*, eabe1508.
- Chen, S., McMullan, G., Faruqi, A.R., Murshudov, G.N., Short, J.M., Scheres, S.H., and Henderson, R. (2013). High-resolution noise substitution to measure overfitting and validate resolution in 3D structure determination by single particle electron cryomicroscopy. *Ultramicroscopy* *135*, 24–35.
- Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. *Acta Crystallogr. D Biol. Crystallogr.* *60*, 2126–2132.
- Goehring, A., Lee, C.H., Wang, K.H., Michel, J.C., Claxton, D.P., Bacongus, I., Althoff, T., Fischer, S., Garcia, K.C., and Gouaux, E. (2014). Screening and large-scale expression of membrane proteins in mammalian cells for structural studies. *Nat. Protoc.* *9*, 2574–2585.
- Green, M.N., Gangwar, S.P., Michard, E., Simon, A.A., Portes, M.T., Barbosa-Caro, J., Wudick, M.M., Lizzio, M.A., Klykov, O., Yelshanskaya, M.V., et al. (2021). Structure of the Arabidopsis thaliana glutamate receptor-like channel GLR3.4. *Mol. Cell* *81*, 3216–3226.
- Kawate, T., and Gouaux, E. (2006). Fluorescence-detection size-exclusion chromatography for precrystallization screening of integral membrane proteins. *Structure* *14*, 673–681.



- Kucukelbir, A., Sigworth, F.J., and Tagare, H.D. (2014). Quantifying the local resolution of cryo-EM density maps. *Nat. Methods* *11*, 63–65.
- McGoldrick, L.L., Singh, A.K., Saotome, K., Yelshanskaya, M.V., Twomey, E.C., Grassucci, R.A., and Sobolevsky, A.I. (2018). Opening of the human epithelial calcium channel TRPV6. *Nature* *553*, 233–237.
- Nadezhdin, K.D., Neuberger, A., Trofimov, Y.A., Krylov, N.A., Sinica, V., Kupko, N., Vlachova, V., Zakharian, E., Efremov, R.G., and Sobolevsky, A.I. (2021). Structural mechanism of heat-induced opening of a temperature-sensitive TRP channel. *Nat. Struct. Mol. Biol.* *28*, 564–572.
- Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and Ferrin, T.E. (2004). UCSF Chimera—a visualization system for exploratory research and analysis. *J. Comput. Chem.* *25*, 1605–1612.
- Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat. Methods* *14*, 290–296.
- Russo, C.J., and Passmore, L.A. (2014). Electron microscopy: Ultrastable gold substrates for electron cryomicroscopy. *Science* *346*, 1377–1380.
- Singh, A.K., McGoldrick, L.L., and Sobolevsky, A.I. (2018). Structure and gating mechanism of the transient receptor potential channel TRPV3. *Nat. Struct. Mol. Biol.* *25*, 805–813.
- Sobolevsky, A.I., Rosconi, M.P., and Gouaux, E. (2009). X-ray structure, symmetry and mechanism of an AMPA-subtype glutamate receptor. *Nature* *462*, 745–756.
- Twomey, E.C., Yelshanskaya, M.V., Grassucci, R.A., Frank, J., and Sobolevsky, A.I. (2016). Elucidation of AMPA receptor-stargazin complexes by cryo-electron microscopy. *Science* *353*, 83–86.
- Twomey, E.C., Yelshanskaya, M.V., Grassucci, R.A., Frank, J., and Sobolevsky, A.I. (2017). Channel opening and gating mechanism in AMPA-subtype glutamate receptors. *Nature* *549*, 60–65.
- Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., Heer, F.T., de Beer, T.A.P., Rempfer, C., Bordoli, L., et al. (2018). SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Res.* *46*, W296–W303.
- Williams, C.J., Headd, J.J., Moriarty, N.W., Prisant, M.G., Videau, L.L., Deis, L.N., Verma, V., Keedy, D.A., Hintze, B.J., Chen, V.B., et al. (2018). MolProbity: more and better reference data for improved all-atom structure validation. *Protein Sci.* *27*, 293–315.
- Zhang, K. (2016). Gctf: real-time CTF determination and correction. *J. Struct. Biol.* *193*, 1–12.
- Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017). MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat. Methods* *14*, 331–332.
- Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and Scheres, S.H. (2018). New tools for automated high-resolution cryo-EM structure determination in RELION-3. *Elife* *7*, e42166.