

# Triaqua(cyclohex-4-ene-1,2-dicarboxylato- $\kappa^1 O^1$ )(1,10-phenanthroline- $\kappa^2 N,N'$ )-cobalt(II)

Hai-Ye Li, Cheng Wang, Fu-Ping Huang and Yi-Min Jiang\*

College of Chemistry and Chemical Engineering, Guangxi Normal University, Guilin, Guangxi 541004, People's Republic of China  
Correspondence e-mail: huangfp2006@163.com


Received 6 September 2012; accepted 22 September 2012

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$ ;  $R$  factor = 0.061;  $wR$  factor = 0.122; data-to-parameter ratio = 12.2.

In the title compound,  $[\text{Co}(\text{C}_8\text{H}_8\text{O}_4)(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_3]$ , the  $\text{Co}^{II}$  atom is coordinated by two N atoms from a bidentate 1,10-phenanthroline ligand, one O atom from a monodentate 4-cyclohexene-1,2-dicarboxylate ligand and three water O atoms in a distorted octahedral geometry. The mononuclear molecules are engaged in extensive intra- and intermolecular O–H···O hydrogen-bonding interactions and  $\pi$ – $\pi$  stacking interactions [centroid–centroid distance = 3.784 (3) Å], forming a three-dimensional supramolecular network.

## Related literature

For background to compounds with metal-organic framework structures, see: Huang *et al.* (2010); Ockwig *et al.* (2005); Rao *et al.* (2004). For a description of the Cambridge Structural Database (CSD), see: Allen (2002). For 4-cyclohexene-1,2-dicarboxylates, see: Kim *et al.* (2004); Lee *et al.* (2006). For related structures, see: Baruah *et al.* (2007); Hou *et al.* (2007); Zhang *et al.* (2008).



## Experimental

### Crystal data

$[\text{Co}(\text{C}_8\text{H}_8\text{O}_4)(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_3]$   
 $M_r = 461.33$

Monoclinic,  $P2_1/c$   
 $a = 8.1730 (16)\text{ \AA}$

$b = 20.210 (4)\text{ \AA}$   
 $c = 12.068 (2)\text{ \AA}$   
 $\beta = 91.44 (3)^\circ$   
 $V = 1992.7 (7)\text{ \AA}^3$   
 $Z = 4$

Mo  $K\alpha$  radiation  
 $\mu = 0.91\text{ mm}^{-1}$   
 $T = 293\text{ K}$   
 $0.40 \times 0.20 \times 0.08\text{ mm}$

### Data collection

Bruker SMART CCD area-detector diffractometer  
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)  
 $(SADABS$ ; Sheldrick, 1996)  
 $R_{\min} = 0.801$ ,  $T_{\max} = 0.945$

15996 measured reflections  
3606 independent reflections  
2982 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.073$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$   
 $wR(F^2) = 0.122$   
 $S = 0.99$   
3606 reflections  
295 parameters  
9 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.34\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O5–H5A···O3 <sup>i</sup>	0.85 (3)	1.85 (3)	2.695 (4)	171 (4)
O5–H5B···O4	0.86 (4)	2.06 (4)	2.912 (4)	172 (4)
O6–H6A···O4 <sup>i</sup>	0.86 (3)	1.86 (4)	2.716 (4)	174 (5)
O6–H6B···O3	0.86 (3)	1.99 (3)	2.835 (4)	165 (3)
O7–H7A···O2	0.86 (3)	1.77 (4)	2.610 (4)	165 (4)
O7–H7B···O4 <sup>ii</sup>	0.86 (3)	1.87 (3)	2.734 (4)	175 (4)

Symmetry codes: (i)  $-x + 1, -y + 1, -z$ ; (ii)  $x - 1, y, z$ .

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We gratefully acknowledge the Guangxi Natural Science Foundation of China (2010GXNSFD013017, 2012GXNSFBA053017) and the Natural Science Foundation of Guangxi Normal University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZQ2181).

## References

- Allen, F. H. (2002). *Acta Cryst. B* **58**, 380–388.
- Baruah, A. M., Karmakar, A. & Baruah, J. B. (2007). *Polyhedron*, **26**, 4479–4488.
- Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Hou, Y.-J., Yu, Y.-H., Sun, Z.-Z., Li, B.-Y. & Hou, G.-F. (2007). *Acta Cryst. E* **63**, m1884.
- Huang, F.-P., Tian, J.-L., Li, D.-D., Chen, G.-J., Gu, W., Yan, S.-P., Liu, X., Liao, D.-Z. & Cheng, P. (2010). *Inorg. Chem.* **49**, 2525–2529.
- Kim, D. S., Forster, P. M., Delgado, G. D., Park, S. E. & Cheetham, A. K. (2004). *Dalton Trans.* pp. 3365–3369.
- Lee, C., Mellot-Draznieks, C., Slater, B., Wu, G., Harrison, W. T. A., Rao, C. N. R. & Cheetham, A. K. (2006). *Chem. Commun.* pp. 2687–2689.
- Ockwig, N. W., Delgado-Friedrichs, O., O'Keeffe, M. & Yaghi, O. M. (2005). *Acc. Chem. Res.* **38**, 176–182.

## metal-organic compounds

---

- Rao, C. N. R., Natarajan, S. & Vaidhyanathan, R. (2004). *Angew. Chem. Int. Ed.* **43**, 1466–1496.  
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
Zhang, B.-Y., Nie, J.-J. & Xu, D.-J. (2008). *Acta Cryst. E* **64**, m986.

# supplementary materials

*Acta Cryst.* (2012). E68, m1375–m1376 [doi:10.1107/S160053681204024X]

## Triaqua(cyclohex-4-ene-1,2-dicarboxylato- $\kappa O^1$ )(1,10-phenanthroline- $\kappa^2 N,N'$ )cobalt(II)

Hai-Ye Li, Cheng Wang, Fu-Ping Huang and Yi-Min Jiang

### Comment

Recently, carboxylic acid (as well as carboxylate) has been widely applied in engineering studies of inorganic-organic hybrid materials and the construction of metal-organic coordination supramolecular complexes (Huang *et al.*, 2010; Ockwig *et al.*, 2005; Rao *et al.*, 2004). Although the Cambridge Structural Database (Allen, 2002) contains a great number of transition metal derivatives of carboxylic acids, the metal derivatives of *cis*-4-cyclohexene-1,2-dicarboxylate acid documented are surprisingly rare (Kim *et al.*, 2004; Lee *et al.*, 2006). As part of this ongoing work, the title complex,  $[\text{Co}(\text{C}_8\text{H}_8\text{O}_4)(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_3]$ , has been prepared and structurally characterized (Fig. 1).

In the title compound, the  $\text{Co}^{II}$  atom is coordinated by two N atoms from a bidentate 1,10-phenanthroline ligand (*phen*), one O atom from a monodentate *cis*-4-cyclohexene-1,2-dicarboxylate ligand, and three O atoms from water molecules in a distorted octahedral geometry. The coordinating Co—N and Co—O bond lengths [ $\text{Co}—\text{N}$  2.107 (3)–2.122 (3) Å;  $\text{Co}—\text{O}$  2.071 (3)–2.152 (3) Å] agree well with those observed in analogous complexes (Baruah *et al.*, 2007; Hou *et al.*, 2007; Zhang *et al.*, 2008). The crystal packing (Fig. 2) exhibits intra- and inter-molecular O—H $\cdots$ O hydrogen bonds (Table 1) and  $\pi$ – $\pi$  stacking interactions [Cg1 $\cdots$ Cg2<sup>iii</sup> distance is 3.784 (3) Å ( $iii = -x, 1-y, 1-z$ ) between the centroids of the (N1-C9-C10-C11-C12-C20) and (C12-C13-C14-C15-C19-C20) six-membered rings] forming a three-dimensional supramolecular network.

### Experimental


For the preparation of the title complex, *cis*-4-cyclohexene-1,2-dicarboxylate acid (0.085 g, 0.5 mmol),  $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  (0.12 g, 0.5 mmol), *phen* (0.10 g, 0.5 mmol) and  $\text{KHCO}_3$  (0.10 g, 1 mmol) were dissolved in a water/ethanol solution (20 ml, 1:1). The solution was stirred for 3 h at room temperature and filtered. Red block-shaped crystals were obtained from the filtrate after 4 d.

### Refinement


H atoms of water molecules were located in a difference Fourier map and refined with distance restraints of  $\text{O}—\text{H} = 0.85$  (2) Å and  $\text{H}—\text{H} = 1.39$  (2) Å. All other H atoms were positioned geometrically and refined using a riding model, with  $\text{C}—\text{H} = 0.93$  Å for  $\text{C}—\text{H}_{\text{aromatic}}$  and  $\text{C}—\text{H} = 0.97$  Å for  $\text{C}—\text{H}_{\text{aliphatic}}$  [ $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ ].

### Computing details

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT* (Bruker, 1998); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

**Figure 1**

The asymmetric unit of the title compound with the atom-numbering scheme and displacement ellipsoids at the 30% probability level.

**Figure 2**

A packing diagram of the title compound. Hydrogen bonds are shown as dotted lines.

### Triaqua(cyclohex-4-ene-1,2-dicarboxylato- $\kappa$ O¹)(1,10-phenanthroline- $\kappa^2$ N,N')cobalt(II)

#### Crystal data



$M_r = 461.33$

Monoclinic,  $P2_1/c$

Hall symbol: -P 2ybc

$a = 8.1730 (16)$  Å

$b = 20.210 (4)$  Å

$c = 12.068 (2)$  Å

$\beta = 91.44 (3)^\circ$

$V = 1992.7 (7)$  Å<sup>3</sup>

$Z = 4$

$F(000) = 956$

$D_x = 1.538 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 2567 reflections

$\theta = 1.5\text{--}25.3^\circ$

$\mu = 0.91 \text{ mm}^{-1}$

$T = 293$  K

Block, red

$0.40 \times 0.20 \times 0.08$  mm

#### Data collection

Bruker SMART CCD area-detector

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and  $\omega$  scans

Absorption correction: multi-scan  
 (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.801$ ,  $T_{\max} = 0.945$   
 15996 measured reflections  
 3606 independent reflections  
 2982 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.073$   
 $\theta_{\max} = 25.3^\circ$ ,  $\theta_{\min} = 3.1^\circ$   
 $h = -9 \rightarrow 9$   
 $k = -24 \rightarrow 24$   
 $l = -14 \rightarrow 14$

#### Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.061$   
 $wR(F^2) = 0.122$   
 $S = 0.99$   
 3606 reflections  
 295 parameters  
 9 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0338P)^2 + 5.P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{\AA}^{-3}$

#### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

#### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.23393 (6)	0.53534 (3)	0.19466 (4)	0.02844 (17)
C1	0.3182 (5)	0.67605 (19)	0.1389 (3)	0.0319 (9)
C2	0.4497 (5)	0.71600 (19)	0.0814 (3)	0.0327 (9)
H2A	0.4249	0.7143	0.0016	0.039*
C3	0.4475 (6)	0.7890 (2)	0.1160 (4)	0.0463 (12)
H3A	0.5051	0.8151	0.0620	0.056*
H3B	0.3352	0.8045	0.1169	0.056*
C4	0.5250 (6)	0.7988 (2)	0.2267 (5)	0.0567 (14)
H4A	0.5006	0.8372	0.2655	0.068*
C5	0.6267 (6)	0.7561 (3)	0.2732 (4)	0.0545 (14)
H5C	0.6676	0.7657	0.3440	0.065*
C6	0.6815 (5)	0.6932 (2)	0.2206 (4)	0.0400 (11)
H6C	0.6419	0.6560	0.2632	0.048*
H6D	0.8001	0.6916	0.2228	0.048*
C7	0.6202 (5)	0.68620 (19)	0.1002 (3)	0.0319 (9)
H7C	0.6947	0.7123	0.0553	0.038*
C8	0.6262 (5)	0.61588 (19)	0.0556 (3)	0.0305 (9)
C9	0.3023 (5)	0.5544 (2)	0.4478 (3)	0.0387 (10)
H9A	0.3474	0.5952	0.4298	0.046*

C10	0.3035 (6)	0.5346 (2)	0.5592 (4)	0.0460 (12)
H10A	0.3491	0.5619	0.6138	0.055*
C11	0.2385 (6)	0.4760 (2)	0.5869 (4)	0.0446 (12)
H11A	0.2392	0.4628	0.6608	0.054*
C12	0.1695 (5)	0.4345 (2)	0.5047 (3)	0.0369 (10)
C13	0.0959 (6)	0.3723 (2)	0.5252 (4)	0.0446 (11)
H13A	0.0915	0.3568	0.5976	0.054*
C14	0.0322 (6)	0.3351 (2)	0.4419 (4)	0.0457 (12)
H14A	-0.0140	0.2942	0.4577	0.055*
C15	0.0348 (5)	0.3577 (2)	0.3300 (3)	0.0359 (10)
C16	-0.0345 (5)	0.3215 (2)	0.2400 (4)	0.0426 (11)
H16A	-0.0839	0.2807	0.2518	0.051*
C17	-0.0280 (5)	0.3472 (2)	0.1365 (4)	0.0429 (11)
H17A	-0.0732	0.3242	0.0764	0.051*
C18	0.0467 (5)	0.4080 (2)	0.1210 (3)	0.0357 (10)
H18A	0.0505	0.4246	0.0493	0.043*
C19	0.1057 (5)	0.41830 (19)	0.3069 (3)	0.0295 (9)
C20	0.1743 (5)	0.4581 (2)	0.3947 (3)	0.0306 (9)
N1	0.2392 (4)	0.51715 (16)	0.3679 (3)	0.0312 (8)
N2	0.1127 (4)	0.44351 (16)	0.2022 (3)	0.0295 (8)
O1	0.3622 (3)	0.62360 (13)	0.1889 (2)	0.0315 (6)
O2	0.1735 (3)	0.69566 (14)	0.1286 (3)	0.0439 (8)
O3	0.5410 (3)	0.60345 (14)	-0.0292 (2)	0.0382 (7)
O4	0.7164 (3)	0.57339 (13)	0.1038 (2)	0.0356 (7)
O5	0.4681 (4)	0.48671 (15)	0.1938 (2)	0.0349 (7)
O6	0.2453 (4)	0.53756 (15)	0.0198 (2)	0.0334 (7)
O7	0.0207 (4)	0.59085 (15)	0.1985 (3)	0.0378 (7)
H5A	0.465 (6)	0.4552 (16)	0.147 (3)	0.062 (17)*
H6A	0.258 (5)	0.5009 (13)	-0.015 (4)	0.061 (17)*
H7A	0.054 (5)	0.6283 (14)	0.174 (4)	0.052 (15)*
H5B	0.538 (5)	0.5151 (17)	0.172 (4)	0.059 (17)*
H6B	0.324 (4)	0.5630 (16)	0.000 (4)	0.054 (16)*
H7B	-0.077 (3)	0.584 (2)	0.172 (4)	0.059 (16)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Co1	0.0293 (3)	0.0295 (3)	0.0264 (3)	-0.0004 (2)	0.0010 (2)	0.0015 (2)
C1	0.037 (3)	0.025 (2)	0.033 (2)	0.0006 (18)	0.0004 (18)	-0.0067 (18)
C2	0.039 (2)	0.028 (2)	0.031 (2)	0.0023 (18)	0.0014 (18)	0.0018 (18)
C3	0.044 (3)	0.027 (2)	0.069 (3)	0.001 (2)	0.008 (2)	0.004 (2)
C4	0.053 (3)	0.038 (3)	0.080 (4)	0.002 (2)	0.005 (3)	-0.028 (3)
C5	0.055 (3)	0.058 (3)	0.050 (3)	-0.009 (3)	-0.001 (2)	-0.030 (3)
C6	0.039 (3)	0.040 (3)	0.040 (3)	-0.005 (2)	-0.005 (2)	-0.008 (2)
C7	0.031 (2)	0.031 (2)	0.034 (2)	-0.0034 (18)	0.0025 (18)	0.0011 (18)
C8	0.028 (2)	0.030 (2)	0.034 (2)	-0.0042 (18)	0.0074 (18)	-0.0021 (18)
C9	0.045 (3)	0.038 (2)	0.033 (2)	-0.005 (2)	0.000 (2)	-0.0021 (19)
C10	0.059 (3)	0.047 (3)	0.032 (2)	-0.002 (2)	-0.010 (2)	-0.010 (2)
C11	0.057 (3)	0.050 (3)	0.027 (2)	0.002 (2)	-0.003 (2)	0.001 (2)
C12	0.039 (3)	0.042 (3)	0.030 (2)	0.002 (2)	0.0023 (19)	0.0049 (19)

C13	0.054 (3)	0.050 (3)	0.030 (2)	-0.004 (2)	-0.001 (2)	0.011 (2)
C14	0.054 (3)	0.043 (3)	0.040 (3)	-0.009 (2)	0.002 (2)	0.011 (2)
C15	0.035 (2)	0.038 (2)	0.035 (2)	-0.0007 (19)	-0.0006 (19)	0.0013 (19)
C16	0.040 (3)	0.037 (3)	0.051 (3)	-0.011 (2)	-0.001 (2)	0.002 (2)
C17	0.043 (3)	0.049 (3)	0.036 (3)	-0.012 (2)	-0.005 (2)	-0.004 (2)
C18	0.038 (2)	0.043 (3)	0.026 (2)	-0.003 (2)	-0.0022 (18)	-0.0007 (19)
C19	0.027 (2)	0.033 (2)	0.028 (2)	-0.0013 (17)	0.0009 (17)	0.0007 (17)
C20	0.030 (2)	0.035 (2)	0.026 (2)	0.0045 (18)	0.0017 (17)	0.0014 (18)
N1	0.033 (2)	0.0333 (19)	0.0272 (19)	0.0009 (15)	0.0018 (15)	-0.0008 (15)
N2	0.0291 (18)	0.0335 (19)	0.0261 (18)	-0.0026 (15)	0.0024 (14)	-0.0007 (14)
O1	0.0310 (15)	0.0284 (15)	0.0349 (16)	-0.0012 (12)	-0.0008 (12)	0.0018 (12)
O2	0.0292 (17)	0.0342 (17)	0.068 (2)	0.0063 (13)	-0.0034 (15)	0.0008 (15)
O3	0.0409 (17)	0.0386 (17)	0.0349 (17)	0.0004 (13)	-0.0023 (14)	-0.0103 (13)
O4	0.0337 (16)	0.0298 (16)	0.0432 (18)	0.0029 (13)	-0.0031 (13)	-0.0023 (13)
O5	0.0360 (18)	0.0356 (17)	0.0329 (17)	0.0033 (14)	0.0010 (13)	-0.0023 (14)
O6	0.0376 (18)	0.0334 (17)	0.0293 (16)	-0.0010 (14)	0.0023 (12)	-0.0015 (14)
O7	0.0280 (17)	0.0381 (19)	0.0474 (19)	0.0029 (14)	0.0000 (14)	0.0003 (15)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

Co1—O1	2.071 (3)	C9—H9A	0.9300
Co1—O7	2.074 (3)	C10—C11	1.345 (6)
Co1—N2	2.107 (3)	C10—H10A	0.9300
Co1—O6	2.115 (3)	C11—C12	1.407 (6)
Co1—N1	2.122 (3)	C11—H11A	0.9300
Co1—O5	2.152 (3)	C12—C20	1.412 (5)
C1—O2	1.250 (5)	C12—C13	1.418 (6)
C1—O1	1.268 (5)	C13—C14	1.348 (6)
C1—C2	1.525 (6)	C13—H13A	0.9300
C2—C7	1.530 (6)	C14—C15	1.427 (6)
C2—C3	1.533 (6)	C14—H14A	0.9300
C2—H2A	0.9800	C15—C19	1.387 (6)
C3—C4	1.477 (7)	C15—C16	1.415 (6)
C3—H3A	0.9700	C16—C17	1.355 (6)
C3—H3B	0.9700	C16—H16A	0.9300
C4—C5	1.315 (7)	C17—C18	1.386 (6)
C4—H4A	0.9300	C17—H17A	0.9300
C5—C6	1.494 (6)	C18—N2	1.319 (5)
C5—H5C	0.9300	C18—H18A	0.9300
C6—C7	1.531 (6)	C19—N2	1.364 (5)
C6—H6C	0.9700	C19—C20	1.433 (5)
C6—H6D	0.9700	C20—N1	1.349 (5)
C7—C8	1.521 (5)	O5—H5A	0.852 (19)
C7—H7C	0.9800	O5—H5B	0.854 (19)
C8—O3	1.250 (5)	O6—H6A	0.860 (19)
C8—O4	1.264 (5)	O6—H6B	0.859 (19)
C9—N1	1.319 (5)	O7—H7A	0.859 (18)
C9—C10	1.403 (6)	O7—H7B	0.858 (19)
O1—Co1—O7		87.78 (12)	N1—C9—H9A
			119.0

O1—Co1—N2	177.61 (12)	C10—C9—H9A	119.0
O7—Co1—N2	94.53 (13)	C11—C10—C9	119.8 (4)
O1—Co1—O6	85.03 (11)	C11—C10—H10A	120.1
O7—Co1—O6	93.97 (12)	C9—C10—H10A	120.1
N2—Co1—O6	95.43 (12)	C10—C11—C12	120.3 (4)
O1—Co1—N1	100.63 (12)	C10—C11—H11A	119.9
O7—Co1—N1	93.88 (12)	C12—C11—H11A	119.9
N2—Co1—N1	78.61 (12)	C11—C12—C20	116.2 (4)
O6—Co1—N1	170.49 (12)	C11—C12—C13	124.7 (4)
O1—Co1—O5	86.69 (11)	C20—C12—C13	119.0 (4)
O7—Co1—O5	174.34 (13)	C14—C13—C12	121.4 (4)
N2—Co1—O5	90.99 (12)	C14—C13—H13A	119.3
O6—Co1—O5	86.76 (11)	C12—C13—H13A	119.3
N1—Co1—O5	85.96 (12)	C13—C14—C15	120.9 (4)
O2—C1—O1	124.7 (4)	C13—C14—H14A	119.6
O2—C1—C2	117.6 (4)	C15—C14—H14A	119.6
O1—C1—C2	117.7 (4)	C19—C15—C16	117.6 (4)
C1—C2—C7	112.0 (3)	C19—C15—C14	119.3 (4)
C1—C2—C3	111.9 (3)	C16—C15—C14	123.1 (4)
C7—C2—C3	110.8 (3)	C17—C16—C15	119.0 (4)
C1—C2—H2A	107.3	C17—C16—H16A	120.5
C7—C2—H2A	107.3	C15—C16—H16A	120.5
C3—C2—H2A	107.3	C16—C17—C18	119.4 (4)
C4—C3—C2	111.6 (4)	C16—C17—H17A	120.3
C4—C3—H3A	109.3	C18—C17—H17A	120.3
C2—C3—H3A	109.3	N2—C18—C17	123.8 (4)
C4—C3—H3B	109.3	N2—C18—H18A	118.1
C2—C3—H3B	109.3	C17—C18—H18A	118.1
H3A—C3—H3B	108.0	N2—C19—C15	122.9 (4)
C5—C4—C3	123.3 (4)	N2—C19—C20	116.8 (3)
C5—C4—H4A	118.3	C15—C19—C20	120.3 (4)
C3—C4—H4A	118.3	N1—C20—C12	123.0 (4)
C4—C5—C6	124.8 (5)	N1—C20—C19	117.9 (3)
C4—C5—H5C	117.6	C12—C20—C19	119.1 (4)
C6—C5—H5C	117.6	C9—N1—C20	118.6 (3)
C5—C6—C7	112.8 (4)	C9—N1—Co1	128.3 (3)
C5—C6—H6C	109.0	C20—N1—Co1	113.0 (3)
C7—C6—H6C	109.0	C18—N2—C19	117.3 (3)
C5—C6—H6D	109.0	C18—N2—Co1	129.1 (3)
C7—C6—H6D	109.0	C19—N2—Co1	113.6 (2)
H6C—C6—H6D	107.8	C1—O1—Co1	126.8 (2)
C8—C7—C2	110.7 (3)	Co1—O5—H5A	109 (3)
C8—C7—C6	114.2 (3)	Co1—O5—H5B	107 (3)
C2—C7—C6	112.3 (3)	H5A—O5—H5B	108 (3)
C8—C7—H7C	106.4	Co1—O6—H6A	119 (3)
C2—C7—H7C	106.4	Co1—O6—H6B	110 (3)
C6—C7—H7C	106.4	H6A—O6—H6B	107 (3)
O3—C8—O4	123.2 (4)	Co1—O7—H7A	101 (3)
O3—C8—C7	117.1 (4)	Co1—O7—H7B	133 (3)

O4—C8—C7	119.7 (4)	H7A—O7—H7B	108 (3)
N1—C9—C10	122.1 (4)		
O2—C1—C2—C7	-179.6 (3)	N2—C19—C20—N1	0.4 (5)
O1—C1—C2—C7	-3.0 (5)	C15—C19—C20—N1	-178.9 (4)
O2—C1—C2—C3	55.3 (5)	N2—C19—C20—C12	179.7 (3)
O1—C1—C2—C3	-128.1 (4)	C15—C19—C20—C12	0.4 (6)
C1—C2—C3—C4	77.6 (5)	C10—C9—N1—C20	-0.1 (6)
C7—C2—C3—C4	-48.1 (5)	C10—C9—N1—Co1	-177.0 (3)
C2—C3—C4—C5	20.1 (7)	C12—C20—N1—C9	0.5 (6)
C3—C4—C5—C6	1.7 (9)	C19—C20—N1—Co1	179.7 (4)
C4—C5—C6—C7	5.8 (7)	C12—C20—N1—Co1	177.9 (3)
C1—C2—C7—C8	59.8 (4)	C19—C20—N1—Co1	-2.9 (4)
C3—C2—C7—C8	-174.5 (3)	O1—Co1—N1—C9	2.4 (4)
C1—C2—C7—C6	-69.1 (4)	O7—Co1—N1—C9	-86.0 (4)
C3—C2—C7—C6	56.6 (5)	N2—Co1—N1—C9	-179.9 (4)
C5—C6—C7—C8	-161.8 (4)	O5—Co1—N1—C9	88.3 (4)
C5—C6—C7—C2	-34.8 (5)	O1—Co1—N1—C20	-174.6 (3)
C2—C7—C8—O3	34.0 (5)	O7—Co1—N1—C20	96.9 (3)
C6—C7—C8—O3	161.9 (4)	N2—Co1—N1—C20	3.1 (3)
C2—C7—C8—O4	-146.7 (4)	O5—Co1—N1—C20	-88.7 (3)
C6—C7—C8—O4	-18.9 (5)	C17—C18—N2—C19	-0.1 (6)
N1—C9—C10—C11	-0.2 (7)	C17—C18—N2—Co1	178.9 (3)
C9—C10—C11—C12	0.1 (7)	C15—C19—N2—C18	0.7 (6)
C10—C11—C12—C20	0.3 (7)	C20—C19—N2—C18	-178.6 (4)
C10—C11—C12—C13	-179.0 (5)	C15—C19—N2—Co1	-178.4 (3)
C11—C12—C13—C14	-180.0 (5)	C20—C19—N2—Co1	2.3 (4)
C20—C12—C13—C14	0.7 (7)	O7—Co1—N2—C18	85.1 (4)
C12—C13—C14—C15	-0.8 (7)	O6—Co1—N2—C18	-9.4 (4)
C13—C14—C15—C19	0.7 (7)	N1—Co1—N2—C18	178.1 (4)
C13—C14—C15—C16	-178.1 (4)	O5—Co1—N2—C18	-96.2 (4)
C19—C15—C16—C17	0.4 (6)	O1—Co1—N2—C19	69 (3)
C14—C15—C16—C17	179.2 (4)	O7—Co1—N2—C19	-95.9 (3)
C15—C16—C17—C18	0.2 (7)	O6—Co1—N2—C19	169.6 (3)
C16—C17—C18—N2	-0.4 (7)	N1—Co1—N2—C19	-2.9 (3)
C16—C15—C19—N2	-0.9 (6)	O5—Co1—N2—C19	82.8 (3)
C14—C15—C19—N2	-179.8 (4)	O2—C1—O1—Co1	34.6 (6)
C16—C15—C19—C20	178.4 (4)	C2—C1—O1—Co1	-141.7 (3)
C14—C15—C19—C20	-0.5 (6)	O7—Co1—O1—C1	-36.1 (3)
C11—C12—C20—N1	-0.6 (6)	O6—Co1—O1—C1	58.1 (3)
C13—C12—C20—N1	178.8 (4)	N1—Co1—O1—C1	-129.6 (3)
C11—C12—C20—C19	-179.8 (4)	O5—Co1—O1—C1	145.1 (3)
C13—C12—C20—C19	-0.4 (6)		

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
O5—H5A···O3 <sup>i</sup>	0.85 (3)	1.85 (3)	2.695 (4)	171 (4)
O5—H5B···O4	0.86 (4)	2.06 (4)	2.912 (4)	172 (4)
O6—H6A···O4 <sup>i</sup>	0.86 (3)	1.86 (4)	2.716 (4)	174 (5)

## supplementary materials

---

O6—H6B···O3	0.86 (3)	1.99 (3)	2.835 (4)	165 (3)
O7—H7A···O2	0.86 (3)	1.77 (4)	2.610 (4)	165 (4)
O7—H7B···O4 <sup>ii</sup>	0.86 (3)	1.87 (3)	2.734 (4)	175 (4)

Symmetry codes: (i)  $-x+1, -y+1, -z$ ; (ii)  $x-1, y, z$ .