

POSTER PRESENTATION

Open Access

Family history scoring system shows Familial Colorectal Cancer Type X syndrome individuals and Amsterdam I individuals have comparable risk for developing colon cancer

Matthew A Strohacker^{1*}, James M Church², Janet Shenal^{1†}, Ellen McGannon¹

From 14th Annual Meeting of the Collaborative Group of the Americas on Inherited Colorectal Cancer Dallas, TX, USA. 12-13 October 2010

Background

A points-based scoring system defines levels of familial risk for colorectal neoplasia. More than 7 points defines a high-risk group that should undergo colonoscopy every 3 years. Family History Scoring was applied to families fulfilling Amsterdam I criteria and familial colorectal cancer type X to compare colon cancer risk.

Methods

Amsterdam I (AmI) and familial type X (famX) families identified from Jagelman Registry database were scored. Affected probands (AP), unaffected probands (UP), affected sibling (AS), and unaffected sibling (US) of the proband, and a child of each sibling were scored. Median, range, and standard deviation were compared for each syndrome to determine level of risk.

Results

200 patients were included. 181 matched AmI criteria with genetic testing, 19 matched criteria for famX. 37 families with an AP and 12 with an UP matched AmI criteria. 4 with an AP and 3 with an UP that matched criteria for famX. Level of risk for the two syndromes were compared (Table 1). Median family history score was 12 points for individuals fulfilling AmI criteria and 14 for famX. Total mean was 12.8 for AmI and 13.5 for famX. Median was compared between the two syndromes (Figure 1), to assess level of risk.

Conclusions

Family history scoring system has shown that families fulfilling Amsterdam I criteria and familial type X share similar familial risk level for developing colon cancer. Both show dominant inheritance, but molecular pathways differ.

Table 1 Comparing level of risk for AmI and famX (X) syndrome using Family History Scoring


	AP	AP/AS	AP/US	AP/AS Child	AP/US Child	UP	UP/AS	UP/US	UP/AS Child	UP/US Child	Total Median	
Median	Aml	17	18	17	12	5	12	16	12	9	4	12
	X	14	17	14	14	4	12	15	12	N/A	N/A	14
Mean	Aml	18.0	21.0	17.3	13.1	5.2	12.4	15.2	12.5	9.0	4.4	12.8
	X	13.5	17.0	13.5	14.0	4.0	11.3	15.0	11.3	N/A	N/A	13.5
Range	Aml	9-42	14-44	9-30	9-21	4-13	6-19	10-19	6-21	6-12	2-12	
	X	11-15	17	11-15	14	4	7-15	15	7-15	N/A	N/A	
STDEV	Aml	6.3	7.0	4.7	3.4	2.0	4.4	3.5	4.7	2.9	3.4	4.0
	X	1.9	N/A	1.9	N/A	0.0	4.0	N/A	4.0	N/A	N/A	1.9

* Correspondence: strohma@mail.uc.edu

† Contributed equally

¹Taussig Cancer Institute, Cleveland Clinic, Cleveland, OH, USA

Full list of author information is available at the end of the article



Author details

¹Taussig Cancer Institute, Cleveland Clinic, Cleveland, OH, USA. ²Colorectal Surgery, Sanford R. Weiss Center for Hereditary Colorectal Neoplasia, Digestive Disease Institute, Cleveland Clinic, Cleveland, OH, USA.

Published: 10 March 2011

doi:10.1186/1897-4287-9-S1-P35

Cite this article as: Strohacker et al.: Family history scoring system shows Familial Colorectal Cancer Type X syndrome individuals and Amsterdam I individuals have comparable risk for developing colon cancer. *Hereditary Cancer in Clinical Practice* 2011 **9**(Suppl 1):P35.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 BioMed Central