

degradation. The Ki-67/MIB-1 index decreased and the number of macrophages increased after chemotherapy. Moreover, the ratio of GSCs to total tumor cells increased after chemotherapy. GSCs and macrophages constitute the mechanism of resistance to and recurrence after alkylating agent chemotherapy in oligodendrogliomas.

Key words: 1p/19q | neoadjuvant chemotherapy | glioma stem cell

GENETICS/EPIGENETICS (GEN)

GEN-7

LIQUID BIOPSY IN BRAIN TUMOR PATIENTS -THE PRESENT AND FUTURE-

Manabu Natsumeda¹, Jyotaro On¹, Jun Watanabe¹, Yoshihiro Tsukamoto¹, Masayasu Okada¹, Makoto Oishi¹, Yukihiko Fujii¹; ¹Department of Neurosurgery, Brain Research Institute, Niigata University

We have previously published liquid biopsy for the diagnosis of brain tumors including PCNSL (JCO Precision Oncology, 2019; Leukemia and Lymphoma, 2019) and diffuse midline gliomas (DMG) (Diagnostics, 2021). We used the Maxwell RSC cDNA extraction kit to extract circulating tumor DNA (ctDNA from) 1 milliliter of cerebrospinal fluid (CSF), and droplet digital PCR to detect MYD88 L265P mutations in PCNSL and H3F3A K27M mutations in DMG. From our initial experience, we were able to detect a high rate of MYD88 mutations in PCNSL, but not H3F3A mutations in DMG. We also observed that higher concentrations of ctDNA were obtained when prompt centrifugation and storage were done after obtaining CSF. Application of liquid biopsy to early detection of relapse and monitoring of treatment relapse are highly anticipated. In cases of PCNSL, we perform liquid biopsy when relapse is suspected on post-contrast MRI. However interestingly, the rate of MYD88 mutations detected is lower than that of newly-diagnosed cases. We would also like to share our experience performing liquid biopsy in conjunction with CSF cytology in brain tumor patients with evidence of leptomeningeal disease. From our initial experience, we would like to discuss the present limitations and future prospects of liquid biopsy in brain tumor patients.

Key words: Liquid biopsy | MYD88 | H3F3A K27M

GEN-8

REAL-TIME PCR BASED INTRAOPERATIVE GENETIC ANALYSIS FOR GLIOMAS

Katsuhiko Takabayashi¹, Kensuke Tateishi¹, Takahiro Hayashi¹, Jo Sasame¹, Masataka Isoda¹, Youhei Miyake¹, Akito Oshima¹, Hirokuni Honma¹, Tetsuya Yamamoto¹; ¹Yokohama City University Hospital, Department of Neurosurgery, Yokohama, Japan

An individual therapeutic strategy based on the genetic characterization is important in gliomas. However, it has been difficult to obtain genetic features during surgery. In this study, we present an overview of intraoperative genetic analysis using modified real-time PCR method. The tumor specimen was crushed with liquid nitrogen, then extract DNA within 60 minutes. Reagents of real-time PCR for detecting IDH, TERT, and BRAF hot spot mutations were stocked and real-time PCR was performed after mixing the extracted DNA. We used PNA and LNA to detect single nucleotide variant (SNV). The average time from tumor extraction to intraoperative tentative judgement was approximately 100 minutes. Using this system, we preliminarily performed intraoperative genomic analysis in 10 glioma patients. We confirmed that 8 of 10 cases (80%) of intraoperative genomic diagnosis were consistent with post-operative diagnosis by Sanger sequencing. However, we experienced 2 (20%) unmatched cases due to low allele of SNV, which indicates that more advanced system is required for clinical application.

Key words: glioma | Real-time PCR | intraoperative genomic analysis

EXPERIMENTAL THERAPEUTICS (ET)

ET-1

TRANSLATIONAL RESEARCH PLATFORM FOR MALIGNANT BRAIN TUMORS

Kensuke Tateishi¹, Yohei Miyake¹, Taishi Nakamura¹, Jo Sasame¹, Takahiro Hayashi¹, Akito Oshima¹, Hirokuni Honma¹, Naoki Ikegaya¹, Tetsuya Yamamoto¹; ¹Department of Neurosurgery, Yokohama City University, Yokohama, Japan

Introduction: The standard therapy for malignant brain tumors includes surgery and combination therapy with radiation and chemotherapy, but to provide individualized treatment based on the biological and molecular genetic background of the tumor, integrate genetic information with

various functional data are required. In this study, we present an overview of our integrated approaches for translational research and clinical management. Methods: In glioma, pre-and intra-operative clinical information, including intraoperative genetic diagnosis, and intraoperative rapid immunohistochemistry is obtained, then a multidisciplinary treatment approach is started based on these integrated data. Specimens collected intraoperatively are cryopreserved for future analysis, and primary cultured cells are routinely collected. The cultured cells are transplanted into the brain of immunodeficient mice to establish patient-derived xenograft model (PDX). Genetic screening, such as IDH, TERT, BRAF, H3F3A mutation and MGMT methylation analysis are routinely assessed within a few days after surgery and used as information for integrated diagnosis. In case of PDX establishment or recurrence, we perform whole exon sequencing or comprehensive genomic assessment to identify genetic abnormalities. If genomic alterations for possible molecular targeted therapy are identified, we assess drug sensitivity test in vitro and in vivo, which are utilized for research to develop optimal molecular targeted therapy. The results, such as the therapeutic effects of molecular targeted drugs, are used for clinical applications. Results: Since the platform was established, we have treated a total of 286 patients, including 189 gliomas and 37 central nervous system lymphomas based on the integrated information. We are currently collecting clinical data to examine if this integrated approach could provide clinical benefit. Conclusion: The translational research system for malignant brain tumors plays an important role in the promotion of clinical and basic research.

Key words: translational research | brain tumors | research platform

ET-5

BIOLOGICAL EFFECTS OF SIMULTANEOUS USE OF MULTIPLE DRUGS IN NEUTRON CAPTURE THERAPY USING RAT BRAIN TUMOR MODEL

Shinji Kawabata¹, Hideki Kashiwagi¹, Kohei Yoshimura¹, Yusuke Fukuo¹, Ryo Hiramatsu¹, Naosuke Nonoguchi¹, Motomasa Furuse¹, Shin-Ichi Miyatake², Masahiko Wanibuchi¹; ¹Department of Neurosurgery, Osaka Medical and Pharmaceutical University, Osaka, Japan ²Kansai BNCT Medical Center, Osaka Medical College, Osaka, Japan

The world's first clinical trial of boron neutron capture therapy (BNCT), which treats malignant brain tumors with a single dose of neutron irradiation using multiple boron drugs simultaneously, was performed at our institution, and its excellent results have stimulated BNCT research around the world. BNCT is a particle irradiation therapy that biologically targets cancer cells, and is expected to be a "new option for cancer treatment" because it can deliver a dose of radiation at the cellular level. In the case of BNCT using a combination of multiple drugs, a method to appropriately consider the biological effects of the combination in the dose calculation has not been established. At present, BNCT based on an accelerator-based irradiation system and a boron drug (BPA) based on essential amino acids has been approved by the regulatory approval for head and neck cancer and has shown good results in brain tumors. As basic research, we have continued to develop new boron drugs, which will be essential in the future, and have explored the interpretation of the biological effects of multiple boron drugs in combination and the optimal conditions required for drug development. The survival curve of BNCT in a rat brain tumor model showed that the effect of the new drug alone was comparable to that of BPA, and the effect of the combination was improved, but the effect of the combination did not match the prediction of the combined biological effect derived from each drug. However, it has been found that the effect of the combination does not match the prediction based on the combination of biological effects derived from each drug. In other words, even if the equivalent X-ray equivalent dose (Gy-Eq) is calculated, the combined effect of some drugs exceeds the prediction, while the combined effect of other drugs is poor.

Key words: glioma | neutron capture therapy | biological effectiveness

ET-6

GEMCITABINE RADIOSENSITIZATION PRIMES IRRADIATED MALIGNANT MENINGIOMA CELLS FOR SENOLYTIC ELIMINATION BY NAVITOCLOX

Masahiro Yamamoto¹, Chifumi Kitanaka¹; ¹Department of Molecular Cancer Science, Yamagata University, Yamagata, Japan

BACKGROUND: Malignant meningioma is an aggressive tumor that requires adjuvant radiotherapy after surgery, yet there has been no standard systemic therapy established so far. We have demonstrated that malignant meningioma cells are exquisitely sensitive to gemcitabine due to their increased expression of hENT1 and dCK, which play critical roles in the intracellular transport and activation of gemcitabine, respectively (Takeda et al. Oncotarget 8:90996, 2017; Yamamoto et al., Neuro-Oncol 23:945, 2021). Significantly, in support of our findings, the efficacy and safety of gemcitabine have recently been documented in a small case series of patients with recurrent meningiomas, which has further led to a phase 2 clinical trial to evaluate the efficacy of gemcitabine in recurrent high-grade