

## REVIEW

# Coronavirus genomic nsp14-ExoN, structure, role, mechanism, and potential application as a drug target

Mohammed Tahir 

Department of Biology, University of Sulaimani, Sulaimanyah, Kurdistan, Iraq

**Correspondence**Mohammed Tahir, Department of Biology, University of Sulaimani, Kurdistan, Sulaimanyah, Iraq.  
Email: [mohammed.tahir@univsul.edu.iq](mailto:mohammed.tahir@univsul.edu.iq)**Abstract**

The recent coronavirus disease 2019 (COVID-19), causing a global pandemic with devastating effects on healthcare and social-economic systems, has no special antiviral therapies available for human coronaviruses (CoVs). The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) possesses a nonstructural protein (nsp14), with amino-terminal domain coding for proofreading exoribonuclease (ExoN) that is required for high-fidelity replication. The ability of CoVs during genome replication and transcription to proofread and exclude mismatched nucleotides has long hindered the development of anti-CoV drugs. The resistance of SARS-CoV-2 to antivirals, especially nucleoside analogs (NAs), shows the need to identify new CoV inhibition targets. Therefore, this review highlights the importance of nsp14-ExoN as a target for inhibition. Also, nucleoside analogs could be used in combination with existing anti-CoV therapeutics to target the proofreading mechanism.

**KEYWORDS**


coronavirus, exoribonuclease, nonstructural protein 14, proofreading, RNA recombination

## 1 | INTRODUCTION

Coronaviruses (CoVs) are enveloped, single-stranded positive-sense RNA viruses with some of the largest identified RNA viral genomes, about 30 kb. CoVs are a set of related viruses that belong to the *Coronaviridae* family in the order *Nidovirales*.<sup>2</sup> Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the causative agent of coronavirus disease 2019 (COVID-19) and belongs to the genus *Betacoronavirus* along with the severe acute respiratory syndrome coronavirus (SARS-CoV) and the Middle East Respiratory Syndrome coronavirus (MERS-CoV). All three of these viruses raised as novel coronaviruses, are causing respiratory diseases and transmitted from animals to humans (zoonotic viruses), and cause respiratory diseases.<sup>3</sup> SARS-CoV-2 is closest (88%) to two SARS-like bat-derived CoVs (bat-SL-CoVZC45 and bat-SL-CoVZXC21), with (79%) overall SARS-CoV sequence identity and (50%) MERS-CoV identity<sup>4</sup>

The SARS-CoV polycistronic RNA genome encodes 14 open reading frames (ORFs), some of which overlap.<sup>1</sup> The genomes 5' end is capped and has a 3' poly(A) tail with short 5' and 3' untranslated region sequences forming regulatory stem-loop structures and a single leader sequence proximal to the 5' end. Two-thirds of the genome encodes two large polyproteins, pp1a and pp1ab, that are cleaved into 16 nonstructural proteins. The last one-third of the genome encodes structure and additional proteins (Figure 1). The bifunctional nsp14 contains an N-terminal 3'-to-5' exoribonuclease (ExoN) and a C-terminal N7-methyltransferase (N7-MTase) domain. In the 3'-to-5' direction, nsp14 catalyzes nucleoside monophosphate excision from nucleic acids, using a mechanism that relies on two divalent metal ions and a reactive water molecule. This exonucleolytic activity is crucial for the Coronavirus replication proofreading activity, a property lacking in other RNA viruses that improve its fidelity

**Abbreviations:** COVID-19, coronavirus disease 2019; CoVs, coronaviruses; ExoN, exoribonuclease; MERS-CoV, Middle East Respiratory Syndrome coronavirus; MHV, murine hepatitis virus; N7-MTase, N7-methyltransferase; NAs, nucleoside analogs; nsp14, nonstructural protein; ORFs, open reading frames; RdRp, RNA-dependent RNA polymerase; RTC, replication-transcription complex; SARS-CoV, severe acute respiratory syndrome coronavirus; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; sgmRNAs, subgenomic mRNAs; TRSs, transcription regulatory sequences.



**FIGURE 1** Single-stranded RNA genome of severe acute respiratory syndrome coronavirus 2. Two-thirds of the genome encodes two major polyproteins, pp1a and pp1ab, which are divided into 16 nonstructural proteins<sup>40</sup>

replication and has played an essential role in *nidoviral* evolution and genome expansion.<sup>1</sup> The first proofreading protein identified in an RNA virus is the coronavirus ExoN, and targeting ExoN is a promising strategy to develop new coronavirus inhibitors and antiviral sensitizers.<sup>5</sup> The development of anti-CoV drugs has long been hampered by the ability of CoVs during genome replication and transcription to proofread and eliminate mismatched nucleotides (Table 1).

## 2 | CORONAVIRUS PROOFREADING MECHANISM

The replication of RNA virus has a high error rate (or low viral fidelity), resulting in the virus living as diverse genome mutant populations or “quasispecies.” RNA viruses can adapt to various replicative environments and selective stresses due to low replicative fidelity, however, a disastrous error could lead to viral extinction. This factor threatens the

**TABLE 1** Role of nonstructural proteins of SARS-CoV-2

Protein	Function	Reference
Nsp1	It plays a role in host-range restriction, countering the innate antiviral response of the host, and suppressing apoptosis induction during the early stages of infection to promote viral growth.	Jauregui et al. <sup>6</sup> and Narayanan et al. <sup>7</sup>
Nsp2	It is involved in disrupting intracellular host signaling.	Cornillez-Ty et al. <sup>8</sup>
Nsp3	It is suggested that it facilitates mRNA transcript translation while suppressing host protein synthesis.	Lu et al. <sup>9</sup> , Anand et al. <sup>10</sup> , Stobart et al. <sup>11</sup> , Cottam et al. <sup>12</sup> , and Benvenuto et al. <sup>13</sup>
Nsp4	Vital role is replication and the assembly of the replicative structures.	Oostra et al. <sup>14</sup>
Nsp5	Protease activity	Mielech et al. <sup>15</sup>
Nsp6	Autophagosome formation for immunomodulatory protein degradation	Cottam et al. <sup>12</sup>
Nsp7	Primer-Independent RNA polymerase Activity	Xiao et al. <sup>16</sup>
Nsp8	Primase activity	Tan et al. <sup>17</sup>
Nsp9	It has been suggested that it works as a single-stranded RNA binding protein.	Bouvet et al. <sup>18</sup>
Nsp10	It serves as a cofactor for both nsp16's 2'-O-methyltransferase and nsp14's N7-guaninemethyltransferase/exoribonuclease activities.	Bouvet et al. <sup>19</sup> , Deming et al. <sup>20</sup>
Nsp11	Necessary for replication.	Fang et al. <sup>21</sup>
Nsp12	RNA-dependent RNA polymerase.	te Velthuis et al. <sup>22</sup>
Nsp13	RTPase activity hydrolyzes the RNA strand's at 5' end.	Bouvet et al. <sup>18</sup>
Nsp14	Exoribonuclease activity and Methyl transferase.	Minskaia et al. <sup>23</sup> and Chen et al. <sup>24</sup>
Nsp15	Endoribonuclease	Bhardwaj et al. <sup>25</sup>
Nsp16	2'-O-methyltransferase methylates the RNA strand at position 2'.	Chen et al. <sup>26</sup> and Bouvet et al. <sup>18</sup>

Abbreviation: SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

development of antivirals against CoVs and other RNA viruses as they can easily establish drug resistance while preserving viral replicative fitness.<sup>27</sup> The distinctive 3' to 5' exoribonuclease (ExoN) proofreading feature is an additional obstacle to the production of nucleoside analogs (NAs) as antivirals against CoVs. Proofreading activities that protect the virus from mutagenesis are provided by nonstructural protein 14.<sup>5</sup> Within the family *Coronaviridae*, nsp14 is strongly conserved.<sup>28</sup> In Group 2 coronaviruses and possibly all coronaviruses, nsp14-ExoN is involved in RNA proofreading, and the acquisition of ExoN function is necessary for the achievement and preservation of broad genomes of coronaviruses.<sup>29</sup>

### 3 | BIFUNCTIONAL NSP14 PROTEIN

Nsp14 is a 60 kDa bifunctional enzyme involved in replication fidelity with an N-terminal ExoN domain and N7-methyltransferase (N7-MTase) C-terminal domain that is involved in messenger RNA (mRNA) capping.<sup>26</sup> SARS-CoV-2 and SARS-CoV nsp14 share more than 95% similarity in amino acid sequence.<sup>30</sup> Also, nsp14 is involved in many other virus life cycles and pathogenicity processes, including recombination of the viral genome and control of innate immune responses.<sup>31</sup> The exoribonucleolytic activity of nsp14 3'-5' relies on metal ions, preferably Mg<sup>2+</sup> over Mn<sup>2+</sup>, Co<sup>2+</sup>, and Zn<sup>2+</sup> promote residual activity, while catalysis was not supported by Ca<sup>2+</sup>, Ni<sup>2+</sup>, and Cu<sup>2+</sup>.<sup>32</sup>

#### 3.1 | N7-Methyltransferase domain

The C-terminal N7-MTase domain contains a DxG S-adenosyl-L-methionine (SAM)-a binding motif conserved among CoVs and is involved in mRNA capping and host immune response evasion.<sup>33</sup> In pre-mRNA splicing, mRNA export, RNA stability, this cap structure plays essential roles by blocking degradation by the 5'-3' exoribonuclease. Translational initiation by promoting the 4E (eIF4E) binding initiation factor of eukaryotic translation,<sup>34</sup> and escapes from the cellular innate immune system recognition.<sup>35</sup> In cytoplasmic granular compartments, host and viral RNA molecules missing the 5' cap structure are degraded.<sup>36</sup> In SARS-CoV nsp14, mutations in the DxG motif specifically eliminate the activity of N7-MTase that is essential for the formation of mRNA caps, and they have an exact detrimental effect on replication without disrupting the activity of ExoN, demonstrating the importance of this enzymatic function of nsp14.<sup>33</sup>

#### 3.2 | Exoribonuclease domain

The activity of nsp14 exoribonuclease resides in the N terminal.<sup>37</sup> By removing mismatched nucleotides from the 3' end of the rising RNA strand, the ExoN domain is proposed to correct errors produced by the RdRp.<sup>38</sup> CoV nsp14-ExoN is a member of the DEDD superfamily of DNA and RNA exonucleases, similar to the proofreading subunit ( $\epsilon$ ) of *Escherichia coli* DNA polymerase III.<sup>23</sup> This superfamily contains four preserved D-E-D-D acidic residues necessary for enzymatic

action, and mutation in these residues within CoV ExoN significantly decreases the activity or ablates the ExoN.<sup>23</sup> A recent study found that the ExoN domain plays a role in the frequency of subgenomic microRNA (sgmRNA) recombination and recombination patterns in virally infected cells and virions in nsp14. Inactivating nsp14-ExoN in murine hepatitis virus (MHV-CoV), significantly altered recombination patterns and decreased recombination frequency compared with wild-type MHV-CoV.<sup>31</sup> In both MHV-CoV model and SARS-CoV model, ExoN activity is vital for high-fidelity replication.<sup>29</sup>

### 4 | DEDDH SUPERFAMILY

The protein's ExoN domain was first described by comparative sequence analysis that demonstrated a distant association between the N-terminal portion of the protein and the cellular exonuclease DEDD protein superfamily.<sup>1</sup> DEDD refers to four preserved catalytic aspartic acid (D)/glutamic acid (E) (Asp/Glu) amino acids that are needed by the protein to form two metal-binding sites that drive nucleotide excision through a two-metal ion-assisted process.<sup>39</sup> X-ray crystallography of the nsp14/nsp10 complex and its functional ligands were able to identify a range of main factors that highlighted distinct differences between nsp14 and its DEDD superfamily homologs and shed light on several characteristics of the structure and function of the protein.<sup>28</sup>

Ma et al.<sup>28</sup> compared the nsp14 ExoN active sites with proofreading homologs of *E. coli* DNA polymerase I Klenow fragment and the polymerase III subunit  $\epsilon$ . They reported that Asp90, Glu92, Glu191, His268, and Asp2733 were the catalytic residues in nsp14. Glu191 replaced the Asp424 of the Klenow fragment and Asp103 of the  $\epsilon$  subunit. This makes nsp14 a "DEED outlier." Furthermore, Asp243, a catalytic center, is highly preserved and mutating it inhibit nsp14 activity, although its function has yet to be deciphered.<sup>40</sup> nsp14 has exoribonuclease activity and is capable of degrading ssRNA in 3'-5' directions.<sup>23</sup> The nsp14 activity requires divalent metal ions as a cofactor and it is essential to have a secondary structure of the RNA substrate. Furthermore, mutating the conserved D/E residues showed severe defects in viral RNA synthesis, and no viable virus could be recovered.<sup>23</sup> Two Mg<sup>2+</sup> ions are expected to coordinate the five catalytic DEEDh residues that aid in removing misincorporated nucleotides. To activate a nucleophilic attack on the phosphate of the substrate nucleotide, MgA must activate a water molecule, and MgB would then promote the product's exit.<sup>40</sup>

### 5 | THE INTERACTION BETWEEN NSP14-EXON AND THE NSP10

In many processes during the viral infection cycle, interactions between viral proteins play crucial roles. A yeast two-hybrid screen of intraviral protein-protein interactions revealed nsp10 as an interacting partner of nsp14.<sup>41</sup> Experiments in vitro in *E. coli* has identified that dsRNA with a 30 mismatch is the preferred substrate of the nsp10-bound nsp14 that mimics a faulty replication product.

Although nsp14 alone has exoribonuclease activity, this study also showed more than a 35-fold increase in nsp14 activity when bound to nsp10, indicating that nsp10 has a role in stabilizing the active site of ExoN in the correct conformation for catalysis of the substrate.<sup>18</sup> The interaction between nsp10 and nsp14 is figuratively equivalent to a hand (nsp14) over a fist (nsp10) (Figure 2). During exoribonuclease assays, the addition of nsp10 to the complex increased the ExoN activity, indicating that nsp10 improves ExoN activity in situ in the active complex.<sup>42</sup> In the presence of nsp10, ExoN active site residues are adequately positioned and form a highly active ExoN, as determined by the distance observed between catalytic residues.<sup>38</sup> Mutations in the nsp10 domain that interact with nsp14 resulted in a decrease in viral replication fidelity. The complexed protein structure showed that nsp10 confers structural integrity and stability to nsp14's ExoN domain.<sup>18</sup> In the absence of nsp10, the ExoN catalytic pocket partly collapses, accounting for the poor ExoN behavior of nsp14 alone.<sup>38</sup>

## 6 | THE ROLE OF ZINC FINGER MOTIFS DURING VIRAL REPLICATION

A study revealed significant differences between nsp14 ExoN and other members of the DEDD exonuclease superfamily in addition to the revised catalytic motif.<sup>39</sup> Although the core structural elements are similar, nsp14 ExoN has two zinc fingers essential for exoribonuclease activity and a site of interaction with nsp10. Zinc finger 2 is located near the catalytic core, and the domain is disrupted by the action of the enzyme via C261A or H264R mutations, meaning that it has a role in catalysis. On the other hand, zinc finger 1 mutation result in insoluble proteins of nsp14 when expressed in *E. coli*, and this, together with its composition, indicated the role for SARS-CoV nsp14 in structural stability.<sup>40</sup>

The proximity of zinc finger 1 to the interaction site of nsp10 could influence the interaction between nsp10 and nsp14 and thus


decrease the catalytic activity of ExoN. Zinc finger 2 mutations have eliminated viral viability, consistent with previous studies.<sup>30</sup> The induction of a particular mutation in zinc finger 1 of the nsp14-ExoN domain alphacoronavirus created a viable virus that induced a decrease in dsRNA intermediate aggregation and decreased antiviral response and apoptosis compared with the wild-type virus.<sup>43</sup>

## 7 | THE ACTIVITY OF SARS-COV-2 NSP14 EXON IS METAL DEPENDENT

It is also recognized that the nsp14 ExoN domain activity relies on divalent cations.<sup>19</sup> In the presence of both Mg<sup>2+</sup> and Mn<sup>2+</sup>, SARS-CoV-2 ExoN nsp14 is active, with more noticeable activity in the presence of Mg<sup>2+</sup>. In agreement with the data obtained for SARS-CoV nsp14 and other proteins from the DEDD family, Ca<sup>2+</sup>, Ni<sup>2+</sup>, and Cu<sup>2+</sup> did not help catalysis.<sup>19</sup> nsp14 residual activity also depends on the presence of Zn<sup>2+</sup> ion.<sup>39</sup> However, the addition of the chelating agent EDTA to the reaction entirely blocks nsp14 ExoN activity, confirming the significance for this activity of divalent ions, namely Mg<sup>2+</sup> and Mn<sup>2+</sup>, close to that described for the SARS-CoV counterpart.<sup>19</sup>

## 8 | THE ROLE OF NSP14 IN RNA RECOMBINATION

Studies comparing CoV strains closely related to SARS-CoV-2 have indicated that SARS-CoV-2 has acquired the capacity to infect human cells by recombining within the spike protein series.<sup>44</sup> CoV recombination has been reported to be associated with the increase in spread and severe illness, resulting in multiple livestock CoVs vaccine failures.<sup>45</sup> On virus-specific transcription regulatory sequences (TRSs), the CoV replication-transcription complex (RTC) performs intramolecular recombination to produce a set of subgenomic




**FIGURE 2** Structure model of severe acute respiratory syndrome coronavirus nsp10/nsp14 complex. From a side view and 90° rotation concerning the side view, ribbon structures of nsp10 (green) and nsp14 (purple) Each protein's amino-terminal and carboxyl-terminal extremities are indicated by the letters N and C with their respective colors<sup>38</sup>

mRNAs (sgmRNAs) with typical ends of 5' and 3'.<sup>46</sup> Therefore, as a natural part of their replication, CoVs, perform recombination, creating complex populations of recombined RNA molecules.<sup>31</sup>

Gribble et al.<sup>31</sup> reveal that the CoVs nsp14-ExoN is necessary for native recombination, and inactivation of ExoN results in decreased recombination frequency and altered recombination products. The viral genome, notably DNA viruses, had shown that exonucleases are essential for recombination. To test this hypothesis, Gribble et al.<sup>31</sup> infected Murine DBT cells with wild-type (Exon<sup>+</sup>) or ExoN-deleted (ExoN<sup>-</sup>) murine hepatitis virus (MHV). Indeed, infected cells (MHV-ExoN<sup>-</sup>) had significantly decreased unique recombination junctions compared to wild-type (MHV-Exon<sup>+</sup>). Thus, relative to (MHV-Exon<sup>+</sup>), (MHV-ExoN<sup>-</sup>) had reduced recombination junction frequency and number of specific junctions, suggesting that lack of operation of nsp14-ExoN resulted in considerably less recombination throughout infection. Further research requires to clarify the function of SARS-CoV-2 nsp14-ExoN interaction in RNA recombination. The role of nsp14-ExoN in recombination, combined with the several essential integrated functions, make it vulnerable, and highly selective target for antiviral treatment inhibition and viral attenuation.

## 9 | NSP14-EXON AS A TARGET FOR COMBINATION COV INHIBITORS

Since the beginning of the COVID-19 trials, the class of antivirals that are RNA-dependent RNA polymerase (RdRp) inhibitors such as Favipiravir, Remdesivir, Ribavirin, and Galidesivir have been of high importance. These drugs, which are nucleoside analogs, work either by introducing viral RNA mutations or terminating the chain during replication. These medications had a strong effect on viruses that do not have proofreading enzymes.<sup>47</sup> However, in SARS-CoV-2, Due to its 3'–5' exonuclease proofreading operation, ExoN is capable of excising integrated nucleoside analogs. This results in negating these medicines' function to varying extents, depending on the kind of nucleoside analog chemistry (Ribavirin, 5FU, and Remdesivir). Compounds with affinity to the catalytic subunit nsp14-ExoN or those with allosteric effects at critical sites with retained residues cause the entire viral RNA repair to alter conformationally. Therefore, finding an inhibitor for this viral nuclease should be the researchers' priority.<sup>48</sup> ExoN inactivation was shown to create a "mutator phenotype," which was apparent from a 15- to 21-fold rise in the incidence of mutation during replication and passage in cell culture



**FIGURE 3** Schematic describing ExoN proofreading activity and mode of action of inhibitors. (Left Panel) Replication in viruses like Hepatitis C Virus (HCV), with no proofreading mechanism. (Mid panel) Replication in viruses like SARS-CoV with proofreading Exoribonuclease. (Right panel) It is the same as a mid-panel but in an ExoN inhibitor<sup>51</sup>

compared to the wild-type control.<sup>29</sup> There is evidence that Remdesivir has 4.5 times more efficacy in the ExoN mutant background, and Ribavirin has 200 times higher efficacy relative to the wild-type ExoN viral genome.<sup>4,38</sup> This provides a clear justification for a mixture of Favirpiravir/Remdesivir/Ribavirin/Galidesivir and SARS-CoV-2 ExoN inhibitor combinations to be used. There is currently no available drug for inhibiting nsp14. To identify small molecules/peptides/natural molecules that have the potential to inhibit nsp14 ExoN, comprehensive molecular docking studies are urgently needed.<sup>48</sup> The crystal structure of SARS-CoV nsp14-nsp10 offers the possibility of molecular docking of the nsp14 ExoN domain to various available drugs. Three known antivirals, Conivaptan, Hesperidin, and Glycyrrhizic acid, show promise based on docking results and their known inhibitory effects on in vitro beta coronaviruses<sup>49</sup> and in patients.<sup>50</sup> As potential proofreading inhibitors of ExoN (nsp14), dexamethasone metasulfobenzoate, conivaptan, hesperidin, and glycyrrhizic acid, and their use in combination with RdRp inhibitors could lead to potentially high levels of antiviral activity and promising COVID-19 therapy<sup>51</sup> (Figure 3). According to Smith et al.,<sup>5</sup> while inhibitors of ExoN alone might be useful, Combining it with an RNA mutagen would increase the ExoN inhibition's intrinsic fidelity defect and drive high-level mutagenesis. A possible benefit of such an approach would be to quickly drive the virus to extinction while limiting or preventing the virus's ability to resolve reversion inhibition.

## 10 | CONCLUSION

There have been many outbreaks of major zoonotic CoV diseases in the 21st century: SARS-CoV in 2002/3, the ongoing MERS-CoV that began in 2012, and the current worldwide SARS-CoV-2 pandemic. However, the SARS-CoV-2 outbreak has emerged when postgenomics technologies are increasingly evolving in molecular biology in terms of sophistication and target specificity. CoVs have an nsp14, bifunctional enzyme with an amino-terminal domain coding for proofreading exoribonuclease and N7-methyltransferase domain to methylate the viral RNA cap. The exoribonuclease function of ExoN 3'-5' is a central player in a variety of essential life-cycle processes for coronaviruses and has made many previously active antiviral compounds ineffective against CoVs. ExoN is a logical therapeutic target for new technologies in genomics.

## ACKNOWLEDGMENTS

Thanks to Dr. Nawroz O. Kareem (Keele University, UK) and Dr. Anwar O. Mohammad (University of Sulaimani, Iraq) for their critical reviews of the manuscript.

## CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

## ORCID

Mohammed Tahir  <http://orcid.org/0000-0003-1588-469X>

## REFERENCES

1. Snijder EJ, Bredenbeek PJ, Dobbe JC, et al. Unique and conserved features of genome and proteome of SARS-coronavirus, an early split-off from the coronavirus group 2 lineage. *J Mol Biol.* 2003; 331(5):991-1004.
2. King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ. *Virus Taxonomy: Ninth Report of the International Committee on Taxonomy of Viruses.* Cambridge, UK: Elsevier; 2012.
3. Hu B, Guo H, Zhou P, Shi ZL. Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol.* 2020;19:141-154.
4. Shannon A, Le NTT, Selisko B, et al. Remdesivir and SARS-CoV-2: structural requirements at both nsp12 RdRp and nsp14 exonuclease active-sites. *Antiviral Res.* 2020;178:104793.
5. Smith EC, Blanc H, Vignuzzi M, Denison MR. Coronaviruses lacking exoribonuclease activity are susceptible to lethal mutagenesis: evidence for proofreading and potential therapeutics. *PLoS Pathog.* 2013;9(8):e1003565.
6. Jauregui AR, Savalia D, Lowry VK, Farrell CM, Wathelet MG. Identification of residues of SARS-CoV nsp1 that differentially affect inhibition of gene expression and antiviral signaling. *PLoS One.* 2013;8(4):e62416.
7. Narayanan K, Ramirez SI, Lokugamage KG, Makino S. Coronavirus nonstructural protein 1: common and distinct functions in the regulation of host and viral gene expression. *Virus Res.* 2015;202: 89-100.
8. Cornillez-Ty CT, Liao L, Yates JR, Kuhn P, Buchmeier MJ. Severe acute respiratory syndrome coronavirus nonstructural protein 2 interacts with a host protein complex involved in mitochondrial biogenesis and intracellular signaling. *J Virol.* 2009;83(19): 10314-10318.
9. Lu J-H, Zhang DM, Wang GL, et al. Sequence analysis and structural prediction of the severe acute respiratory syndrome coronavirus nsp5. *Acta Biochim Biophys Sin (Shanghai).* 2005;37(7):473-479.
10. Anand K. Coronavirus main proteinase (3CLpro) structure: basis for design of anti-SARS drugs. *Science.* 2003;300(5626): 1763-1767.
11. Stobart CC, Sexton NR, Munjal H, et al. Chimeric exchange of coronavirus nsp5 proteases (3CLpro) identifies common and divergent regulatory determinants of protease activity. *J Virol.* 2013;87(23): 12611-12618.
12. Cottam EM, Whelband MC, Wileman T. Coronavirus NSP6 restricts autophagosome expansion. *Autophagy.* 2014;10(8):1426-1441.
13. Benvenuto D, Angeletti S, Giovanetti M, et al. Evolutionary analysis of SARS-CoV-2: how mutation of non-structural protein 6 (NSP6) could affect viral autophagy. *J Infect.* 2020;81(1):e24-e27.
14. Oostra M, te Lintelo EG, Deijs M, Verheije MH, Rottier PJM, de Haan CAM. Localization and membrane topology of coronavirus nonstructural protein 4: involvement of the early secretory pathway in replication. *J Virol.* 2007;81(22):12323-12336.
15. Mielech AM, Chen Y, Mesecar AD, Baker SC. Nidovirus papain-like proteases: multifunctional enzymes with protease, deubiquitinating and deISGylating activities. *Virus Res.* 2014;194:184-190.
16. Xiao Y, Ma Q, Restle T, et al. Nonstructural proteins 7 and 8 of feline coronavirus form a 2:1 heterotrimer that exhibits primer-independent RNA polymerase activity. *J Virol.* 2012;86(8): 4444-4454.
17. Tan YW, Fung TS, Shen H, Huang M, Liu DX. Coronavirus infectious bronchitis virus non-structural proteins 8 and 12 form stable complex independent of the non-translated regions of viral RNA and other viral proteins. *Virology.* 2018;513:75-84.
18. Bouvet M, Lugari A, Posthuma CC, et al. Coronavirus Nsp10, a critical co-factor for activation of multiple replicative enzymes. *J Biol Chem.* 2014;289(37):25783-25796.
19. Bouvet M, Imbert I, Subissi L, Gluais L, Canard B, Decroly E. RNA 3'-end mismatch excision by the severe acute respiratory syndrome

- coronavirus nonstructural protein nsp10/nsp14 exoribonuclease complex. *Proc Natl Acad Sci USA*. 2012;109(24):9372-9377.
20. Deming DJ, Graham RL, Denison MR, Baric RS. Processing of open reading frame 1a replicase proteins nsp7 to nsp10 in murine hepatitis virus strain A59 replication. *J Virol*. 2007;81(19):10280-10291.
  21. Fang SG, Shen H, Wang J, Tay FPL, Liu DX. Proteolytic processing of polyproteins 1a and 1ab between non-structural proteins 10 and 11/12 of Coronavirus infectious bronchitis virus is dispensable for viral replication in cultured cells. *Virology*. 2008;379(2):175-180.
  22. te Velthuis AJW, Arnold JJ, Cameron CE, van den Worm SHE, Snijder EJ. The RNA polymerase activity of SARS-coronavirus nsp12 is primer dependent. *Nucleic Acids Res*. 2010;38(1):203-214.
  23. Minskaia E, Hertzog T, Gorbalenya AE, et al. Discovery of an RNA virus 3'→5' exoribonuclease that is critically involved in coronavirus RNA synthesis. *Proc Natl Acad Sci USA*. 2006;103(13):5108-5113.
  24. Chen P, Jiang M, Hu T, et al. Biochemical characterization of exoribonuclease encoded by SARS coronavirus. *J Biochem Mol Biol*. 2007;40(5):649-655.
  25. Bhardwaj K, Guarino L, Kao CC. The severe acute respiratory syndrome coronavirus Nsp15 protein is an endoribonuclease that prefers manganese as a cofactor. *J Virol*. 2004;78(22):12218-12224.
  26. Chen Y, Cai H, Pan J, et al. Functional screen reveals SARS coronavirus nonstructural protein nsp14 as a novel cap N7 methyltransferase. *Proc Natl Acad Sci USA*. 2009;106(9):3484-3489.
  27. Perales C, Domingo E. Antiviral strategies based on lethal mutagenesis and error threshold. *Curr Top Microbiol Immunol*. 2015:392-339.
  28. Ma Y., Wu L, Shaw N, et al. Structural basis and functional analysis of the SARS coronavirus nsp14–nsp10 complex. *Proc Natl Acad Sci USA*. 2015;112(30):9436.
  29. Eckerle LD, Becker MM, Halpin RA, et al. Infidelity of SARS-CoV Nsp14-exonuclease mutant virus replication is revealed by complete genome sequencing. *PLoS Pathog*. 2010;6(5):e1000896.
  30. Ogando NS, Zevenhoven-Dobbe JC, van der Meer Y, Bredenbeek PJ, Posthuma CC, Snijder EJ. The enzymatic activity of the nsp14 exoribonuclease is critical for replication of MERS-CoV and SARS-CoV-2. *J Virol*. 2020;94(23):e01246-20.
  31. Gribble J, Stevens LJ, Agostini ML, et al. The coronavirus proofreading exoribonuclease mediates extensive viral recombination. *PLoS Pathog*. 2021;17(1):e1009226.
  32. Saramago M, Bárria C, Costa VG, et al. New targets for drug design: importance of nsp14/nsp10 complex formation for the 3'-5' exoribonucleolytic activity on SARS-CoV-2. *FEBS J*. 2021. <https://doi.org/10.1111/febs.15815>
  33. Case JB, Ashbrook AW, Dermody TS, Denison MR. Mutagenesis of S-adenosyl-L-methionine-binding residues in coronavirus nsp14 N7-methyltransferase demonstrates differing requirements for genome translation and resistance to innate immunity. *J Virol*. 2016;90(16):7248-7256.
  34. Furuichi Y, Shatkin AJ. Viral and Cellular mRNA Capping: Past and Prospects, in *Advances in Virus Research*. Cambridge, MA: Academic Press; 2000:135-184.
  35. Nallagatla SR, Toroney R, Beliacqua PC. A brilliant disguise for self RNA: 5'-end and internal modifications of primary transcripts suppress elements of innate immunity. *RNA Biol*. 2008;5(3):140-144.
  36. Liu H, Kiledjian M. Decapping the message: a beginning or an end. *Biochem Soc Trans*. 2006;34(1):35-38.
  37. Denison MR, Graham RL, Donaldson EF, Eckerle LD, Baric RS. Coronaviruses. *RNA Biol*. 2011;8(2):270-279.
  38. Ferron F, Subissi L, Silveira De Morais AT, et al. Structural and molecular basis of mismatch correction and ribavirin excision from coronavirus RNA. *Proc Natl Acad Sci USA*. 2018;115(2):E162.
  39. Chen P, Jiang M, Hu T, Liu Q, Chen XS, Guo D. Biochemical characterization of exoribonuclease encoded by SARS coronavirus. *J Biochem Mol Biol*. 2007;40(5):649-655.
  40. Robson F, Khan KS, Le TK, et al. Coronavirus RNA proofreading: molecular basis and therapeutic targeting. *Mol Cell*. 2020;79(5):710-727.
  41. Pan J, Peng X, Gao Y, et al. Genome-wide analysis of protein-protein interactions and involvement of viral proteins in SARS-CoV replication. *PLoS One*. 2008;3(10):e3299.
  42. Subissi L, Imbert I, Ferron F, et al. SARS-CoV ORF1b-encoded nonstructural proteins 12–16: replicative enzymes as antiviral targets. *Antiviral Res*. 2014;101:122-130.
  43. Becares M, Pascual-Iglesias A, Nogales A, Sola I, Enjuanes L, Zúñiga S. Mutagenesis of Coronavirus nsp14 reveals its potential role in modulation of the innate immune response. *J Virol*. 2016;90(11):5399-5414.
  44. Patiño-Galindo JÁ, Filip I, AlQuraishi M, Rabadan R. Recombination and lineage-specific mutations led to the emergence of SARS-CoV-2. *bioRxiv*. 2020:2020. <https://doi.org/10.1101/2020.02.10.942748>
  45. Chen N, Li S, Zhou R, et al. Two novel porcine epidemic diarrhoea virus (PEDV) recombinants from a natural recombinant and distinct subtypes of PEDV variants. *Virus Res*. 2017;242:90-95.
  46. Sola I, Almazán A, Zúñiga S, Enjuanes L. Continuous and discontinuous RNA synthesis in coronaviruses. *Annu Rev Virol*. 2015;2(1):265-288.
  47. Warren TK, Jordan R, Lo MK, et al. Therapeutic efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys. *Nature*. 2016;531(7594):381-385.
  48. Senanayake SL. Drug repurposing strategies for COVID-19. *Future Drug Discov*. 2020;2(2):fdd-2020-0010.
  49. De Clercq E. Potential antivirals and antiviral strategies against SARS coronavirus infections. *Expert Rev Anti Infect Ther*. 2006;4(2):291-302.
  50. Hoever G, Baltina L, Michaelis M, et al. Antiviral activity of glycyrrhizic acid derivatives against SARS–coronavirus. *J Med Chem*. 2005;48(4):1256-1259.
  51. Khater S, Dasgupta N, Das G. Combining SARS-CoV-2 proofreading exonuclease and RNA-dependent RNA polymerase inhibitors as a strategy to combat COVID-19: a high-throughput in silico screen. *OSF Preprints*. 2020.

**How to cite this article:** Tahir M. Coronavirus genomic nsp14-ExoN, structure, role, mechanism, and potential application as a drug target. *J Med Virol*. 2021;93:4258–4264. <https://doi.org/10.1002/jmv.27009>