

Original article:

NEUROPROTECTIVE MECHANISM OF LOW-DOSE SODIUM NITRITE IN OXYGEN-GLUCOSE DEPRIVATION MODEL OF CEREBRAL ISCHEMIC STROKE IN PC12 CELLS

Nader Shakib¹, Mohammad Hassan Khadem Ansari^{1*}, Poursan Karimi^{2*}, Yousef Rasmi¹

¹ Department of Biochemistry, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran

² Neurosciences Research Center (NSRC), Tabriz University of Medical Sciences, Tabriz, Iran

* Corresponding authors: Dr. Mohammad Hassan Khadem Ansari, Department of Biochemistry, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran; E-mail: ansari_mh@umsu.ac.ir;
Dr. Poursan Karimi, Neurosciences Research Center (NSRC), Tabriz University of Medical Sciences, Tabriz, Iran; E-mail: karimip@tbzmed.ac.ir

<http://dx.doi.org/10.17179/excli2018-1947>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

ABSTRACT

The purpose of this study was to clarify the mechanisms of the protective effects of low-dose sodium nitrite (SN) on oxygen and glucose deprivation (OGD)-induced endoplasmic reticulum (ER) stress in PC12 cells. The PC12 cells were exposed to 4 h of OGD and treated with 100 μ mol SN. The expression and activity of ER stress markers, including PKR-like endoplasmic reticulum kinase (PERK), transcription factor 6 (ATF6), CCAAT/enhancer binding protein homologous protein (CHOP), as well as caspase-12 and -3, were detected by immunoblotting assay. Fluorescence staining was used to detect the levels of reactive oxygen species (ROS) and Ca^{2+} release from the ER. Cell viability was also evaluated by MTT assay. It was found that SN significantly inhibited ROS production and Ca^{2+} release from the ER in OGD-injured PC12 cells. Moreover, ER stress marker expression and cleaved fragments of caspase-3 and -12 in OGD-injured PC12 cells were decreased after SN treatment. These findings were accompanied by a significant increase in cell viability. It seems that SN exerts a neuroprotective effect at least partially through reduction of ROS-mediated ER stress caused by OGD insult.

Keywords: Sodium nitrite, oxygen and glucose deprivation, PC12 cells, endoplasmic reticulum stress, Ca^{2+} release

INTRODUCTION

According to World Health Organization statistics, stroke is the second leading cause of death worldwide after myocardial infarction (Moskowitz et al., 2010). Recent developments have shown that endoplasmic reticulum (ER) stress is an important cause of neuronal injury following cerebral ischemia

(Anelli and Sitia, 2008; Ma and Hendershot, 2004; Nakka et al., 2010). Neurons are vulnerable to ischemia and disturbance in the central nervous system leads to severe effects such as paresis, memory disruption and even neuronal death (Anderson and Arciniegas, 2010). Hypoxia and hypoglycemia, two common features in ischemic stroke, increase the

generation of reactive oxygen species (ROS), cellular membrane depolarization, and glutamate release from neurons. Moreover, in neuronal cells, ischemia increases the release of Ca^{2+} from the ER to the cytosol, promoting excitotoxicity and apoptosis (Ankarcona et al., 1995; Lipton and Rosenberg, 1994; Pellegrini-Giampietro et al., 1997).

The ER is the main organelle for secretory pathways in all eukaryotic cells. Neuronal cells in particular have a highly developed ER. This organelle contributes to protein folding, biosynthesis, translocation, and post-translational modifications including glycosylation, disulfide bond formation and chaperone-mediated protein folding processes, which are essential for effective functioning. ER also serves as a cellular Ca^{2+} reservoir (Chang et al., 2006; Hampton, 2002; Schroder and Kaufman, 2005). During brain ischemia, neuronal cells endure lack of the oxygen and glucose required for ATP synthesis, which results in a disturbance in energy consuming protein-folding and the accumulation of unfolded proteins (UPs) in the ER (DeGracia et al., 2002). Accumulated UPs increase the size of ER membranes, accelerate degradation of UPs, increase the translation of folding chaperones, and inhibit synthesis of other proteins, which as a whole is denoted as ER stress (Herrmann et al., 2013; Kaufman, 1999; Kaufman et al., 2002; Oyadomari et al., 2002).

In response to ER stress, three important sensor and transducer proteins on the ER membrane, PKR-like endoplasmic reticulum kinase (PERK), transcription factor 6 (ATF6), and inositol-requiring enzyme 1 (IRE1), are activated (Harding et al., 2000; Haze et al., 1999; Wang et al., 1998). When cellular stress is very serious or prolonged, these components can promote apoptosis signaling pathways through the activation of the CCAAT/enhancer binding protein homologous protein (CHOP), c-Jun N-terminal kinase (JNK) and caspase-12 proteins (Ferri and Kroemer, 2001; Paschen and Mengesdorf, 2005; Szegezdi et al., 2006). The induction of CHOP mRNA by brain ischemia or hypoxia

has been demonstrated in previous studies to be one of the most significant pathways leading to neuronal apoptosis (Carmeliet et al., 1998; Doutheil et al., 1997; Jin et al., 2001; Paschen et al., 1998), CHOP can also reduce the expression of B-cell lymphoma-2 (Bcl-2) leading to ER stress and an increase in oxidative damage (Ferri and Kroemer, 2001; Mori, 2000).

Caspase-12 is a member of the interleukin-1 β converting enzyme (ICE) subfamily of caspases which normally exist in an inactive pro-enzyme state attached to the ER membrane (Badiola et al., 2011; Nakagawa et al., 2000). Active caspase-12 triggers caspase-9 and caspase-3 activation, resulting in DNA fragmentation. Caspase-12 activation can be induced by oxygen and glucose deprivation (OGD) in glial cells (Martinez et al., 2010; Rao et al., 2001).

It has been reported that the overproduction of ROS is a substantial cause of cerebral injury following ischemia/reperfusion (Qi et al., 2016). Mitochondria which are damaged during prolonged ischemic insult are the predominant source of ROS. ROS generation initiates several intracellular signaling cascades and concomitantly induces mitochondrial dysfunction and ER stress, which activates apoptosis (Hildeman et al., 2003; Ray et al., 2012; Tajiri et al., 2004).

Nitrite, a physiological reservoir of nitric oxide (NO), is a powerful mediator of cytoprotection following ischemia/reperfusion which acts by reducing mitochondrial ROS generation. Moreover, NO-derived from NO donors (NODs) has a scavenging effect on ROS and directly reacts with ROS or regulates mitochondrial ROS levels by mitochondrial complex I modification (Pluta et al., 2001; Raat et al., 2009; Steiner et al., 2002). Nitrite is reduced to NO by the reductase system under ischemia/hypoxia conditions (Calvert and Lefer, 2010).

In the present study, the effect of sodium nitrite (SN) was investigated on ROS production, Ca^{2+} release and ER stress-related markers (phospho-PERK, ATF6, CHOP, caspase-

12, and caspase-3) in OGD-exposed PC12 cells.

MATERIALS AND METHODS

Chemicals and antibodies

The following antibodies (Santa Cruz Biologics; USA) were used in Western blotting: polyclonal rabbit antibodies against caspase-3 (1:500), caspase-12 (1:500), CHOP (1:500), p90ATF6 (1:500) and phospho-PERK (1:500). Nitrite sodium (Sigma; 563218) was a gift from Dr. Amiri. Fetal bovine serum (FBS), Dulbecco's Modified Eagle Medium (DMEM) and trypsin were obtained from Hyclone (USA). Sigma-Aldrich Fluorometric Intracellular ROS kits and green fluorescence (MAK143) were used for ROS staining and Goryo Chemical red fluorescent Ca^{2+} probe CaTM-3 AM (GC 501) was used to determine Ca^{2+} release.

Cell culture

The pheochromocytoma derived-PC12 cell line was purchased from Pasteur Institute (Iran), cultured in DMEM enriched with 10 % horse serum and incubated under normoxic conditions (95 % air and 5 % CO_2) at 37 °C. The four study groups were: normal control (NC), normoxic/normoglycemic (NO/NG), OGD for 4 h (OGD), 100 μM SN added to the culture medium concomitantly with OGD onset (OGD+SN) and 100 μM SN added to NO/NG (NC+ SN).

For induction of the *in vitro* OGD model, PC12 cells were cultured in six-well plates in the NO/NG condition. At 80 % confluence, the culture medium was replaced by glucose-free Hank's Balanced Salt Solution (HBSS) and incubated in a homemade hypoxia chamber (95 % N_2 , 5 % CO_2) at humidity 60 % for 4 h. In the treatment groups, PC12 cells were cultured in the aforementioned media and 100 μM SN was applied to the medium in the presence or absence of OGD.

ROS level detection

ROS or its intermediates were produced by the uncompleted reduction of oxygen. Or-

ganisms living under aerobic conditions generate various kinds of ROS molecules, such as superoxide ($\bullet\text{O}_2^-$), hydrogen peroxide (H_2O_2), hydroxyl radicals (OH^-) and singlet oxygen. ROS are highly reactive molecules and are exceedingly unstable, so detection of ROS depends on measurement of the end products formed when they react with specific substances. ROS such as superoxide and hydrogen peroxide can be detected conventionally by staining techniques.

In the present study, 2',7'-dichlorodihydrofluorescein diacetate (DCFH-DA; Sigma; USA) was used to measure ROS levels. Briefly, PC12 cells were cultured in growth medium to reach 1×10^6 cells per well in a 96-well microplate and 100 μL /well DCFH-DA working solution was added directly to the medium and incubated at 37 °C for 15 min. The cells were then washed with phosphate buffered saline (PBS) once and kept on ice for immediate DCF detection. Fluorescent images were obtained under a fluorescence microscope (Olympus; Japan). Intracellular ROS generation was monitored by measuring the fluorescent intensity of the cells using a BioTek Synergy plate reader and the results were expressed as the percentage of control. The wavelength used for excitation was set at 490 nm and for emission was set at 525 nm.

Immunoblotting assay

PC12 cells were homogenated in ice-cold RIPA lysis buffer containing protease inhibitors and centrifuged at 12000 g for 15 min at 4 °C. The supernatant was taken for quantification of protein concentration using the Bradford method. The samples then were mixed 1:1 with sample loading buffer 2X (Sigma; USA) and boiled for 10 min before loading onto 10 % SDS-polyacrylamide gel for electrophoresis. After transfer of the proteins onto the methanol pre-activated polyvinylidene fluoride (PVDF) membrane, the membranes were incubated with blocking solution for 2 h at room temperature. The membrane then was incubated overnight with polyclonal anti-ATF6, anti-CHOP, anti-caspase-3 and -12 and anti- β -actin antibodies at 4 °C.

After washing three times with PBS, the membrane was incubated with an appropriate horseradish peroxidase conjugated anti-rabbit secondary antibody containing 0.1 % Tween-20 for 2 h at room temperature. The blots were detected using a chemiluminescence detection kit (Amersham Biosciences; USA) and radiographic film (Fuji; Japan). The density of the target bands was measured by Image J software. β -actin was used as a loading control (Farajdokht et al., 2018).

Ca²⁺ release

Goryo Chemical CaTM-2 was used to determine the rate of Ca²⁺ release. After seeding the cells in six-well plates, the treated and control cells were washed twice in Krebs buffer (pH 7.4) composed of 132 mM NaCl, 4 mM KCl, 1.4 mM MgCl₂, 6 mM glucose, 10 mM HEPES, 10 mM NaHCO₃ and 1 mM CaCl₂. Then, 50 μ g CaTM-2 was dissolved in 41 μ L of DMSO to a concentration of 1 mM. To improve the induction efficiency and inhibit localization of the probe, Pluronic F-127 was added. Next, an aliquot of the stock solution was diluted to a final concentration of 1-10 μ M in loading medium. The culture medium was removed from the wells and washed with loading medium. Stain solution then was added to the wells and incubated for 10-60 min at 37 °C and 5 % CO₂. After staining, the stain solution was discarded and the wells were washed three times. The changes in intracellular fluorescence intensity were measured using a fluorimeter. The excitation wavelength was set at 597 nm and the maximum peak of the fluorescent wavelength was detected at 609 nm.

MTT assay

PC12 cells were seeded in 96-well plates at an initial density of 10⁴ cells/100 μ l/well. After 48 h, when cell count reached 10⁵, cell viability was determined using the 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide (MTT) assay. At the end of the incu-

bation period (4 h), the supernatant was removed and 200 μ l of DMSO was added to each well. Absorbance was read at 570 nm using a microplate reader (Stat fax; Awareness; USA) and the results were expressed as percentage of control.

Statistical analysis

The data are expressed as mean \pm SEM. The comparisons of means were done by one-way analysis of variance (ANOVA) followed by Tukey's post-hoc test (SPSS, version 20; USA). A p-value of <0.05 was considered statistically significant. The statistical power was 0.8 at α = 0.05 and β = 0.2.

RESULTS

Effect of SN on ER stress indicators in OGD-injured PC12 cells

The phosphorylation of PERK (p-PERK) and expressions of ATF6 and CHOP as the common indicators of ER stress were determined by immunoblotting technique under four experimental conditions. The results showed that OGD significantly increased immunoreactivity of p-PERK (p<0.001; Figure 1B) and CHOP (p<0.001; Figure 1D) as compared to the NC, which decreased after the SN treatment. Moreover, OGD considerably decreased ATF6 expression (p<0.001; Figure 1C) as compared to the NC condition. However, parallel SN treatment with OGD onset significantly increased ATF6 expression (p<0.001) in PC12 cells.

SN decreased cleavage of Caspase-3 and Caspase-12 in OGD-treated PC12 cells

The results also demonstrate that OGD markedly increased the ratio of cleaved caspase-12/procaspase-12 (p<0.001; Figure 2B) and cleaved caspase-3/procaspase-3 (p<0.001; Figure 2C) as compared to the NC condition. On the other hand, cleaved caspase-3/procaspase-3 ratio was significantly decreased in the OGD-SN condition (p<0.001).




Figure 1: SN up-regulated ATF6 and down-regulated CHOP and PERK in OGD-injured PC12 cells: **(A)** immunoblotting images of expression of p-PERK, PERK, ATF6, CHOP and β -actin proteins under different experimental conditions; **(B)**, **(C)**, and **(D)**, respectively, quantitative densitometric analysis of the p-PERK, ATF6 and CHOP protein bands. Data are shown as means \pm SEM (n=3): ***p<0.001 vs. NC; ###p<0.01 vs. OGD [NC: normal control; OGD: oxygen and glucose deprivation (4 h); OGD-SN: oxygen and glucose deprivation (4 h) co-treated with SN (100 μ M); SN: sodium nitrite (100 μ M) treated].

SN decreased ROS production in OGD-treated PC12 cells

Figure 3 shows that the percentage of DCF fluorescence intensity as an index of the rate of ROS production increased significantly ($p<0.001$) following 4 h of OGD in PC12 cells. Nevertheless, in the OGD+SN condition, co-treatment with SN (100 μ M) significantly decreased ROS generation levels ($p<0.01$) as compared to the OGD condition (Figure 3).

SN decreased intracellular Ca^{2+} release from ER

Figure 4 shows that the lowest release of Ca^{2+} was observed in the NC and SN conditions. In the OGD condition the release of Ca^{2+} increased considerably ($p<0.001$). Co-treatment of the PC12 cells with 100 μ M SN in the OGD+SN condition significantly decreased Ca^{2+} release ($p<0.01$).




Figure 2: SN down-regulated cleavage of caspase-12 and caspase-3 in OGD injured PC12 cells: **(A)** immunoblotting images of cleaved caspase-12, procaspase-12, cleaved caspase-3, and procaspase-3 in different experimental conditions; **(B)** and **(C)**, respectively, quantitative densitometric analysis of cleaved caspase-12/procaspase-12 and cleaved caspase-3/procaspase-3 ratios. Data are shown as means \pm SEM (n=3): ***p<0.001 vs. NC; ###p<0.01 vs. OGD [NC: normal control; OGD: oxygen and glucose deprivation (4 h); OGD-SN: oxygen and glucose deprivation (4 h) co-treated with SN (100 μ m); SN: sodium nitrite (100 μ m) treated].




Figure 3: SN attenuated ROS levels in OGD-injured PC12 cells: **(A)** fluorescence emission spectrum of DCF probe in PC12 cells exposed to 4 h OGD and SN 100 μ m treatment under different experimental conditions; **(B)** percentage of relative intensity of DCF fluorescence per mg protein (n=3): ***p<0.001 vs. NC; ##p<0.01 vs. OGD [NC: normal control; OGD: oxygen and glucose deprivation (4 h); OGD-SN: oxygen and glucose deprivation (4 h) co-treated with SN (100 μ m); SN: sodium nitrite (100 μ m) treated].




Figure 4: SN decreased Ca²⁺ release from ER in OGD-exposed PC12 cells: **(A)** intracellular Ca²⁺ levels stained by the Fluo4-AM dye; **(B)** relative fluorescence intensity analysis of intracellular Ca²⁺ under different experimental conditions (n=3): ***p<0.001 vs. N; ##p<0.01 vs. OGD [NC: normal control; OGD: oxygen and glucose deprivation (4 h); OGD-SN: oxygen and glucose deprivation (4 h) co-treated with SN (100 μM); SN: sodium nitrite (100 μM) treated].

SN increased cell viability in OGD-treated PC12 cells

The results of MTT showed that cell viability of OGD-exposed PC12 cells decreased markedly as compared with normal cultured cells (p<0.001; Figure 5). Administration of SN (100 μM) concomitantly with OGD onset dramatically increased cell viability (p<0.05).

DISCUSSION

The findings revealed that 4 h OGD increased ROS production and Ca²⁺ release from the ER, increased the expression of ER stress markers as well as caspases-3 and -12, while decreased cell viability. However, SN treatment reversed all changes induced by OGD insult. Recently, we showed that SN has a neuroprotective effect against OGD insult in PC12 cells through down-regulation of mitochondrial-mediated pro-apoptotic markers and up-regulation of anti-apoptotic markers (Ansari et al., 2018). In the current study, we aimed to investigate whether the effect of SN-induced neuroprotective operates through modulation of ER stress markers (PERK,

ATF6 and CHOP) and/or through ROS and Ca²⁺ signaling.




Figure 5: SN increased cell viability in PC12 cells-exposed to 4 h OGD. Data are presented as the percentage of control (OGD 0, normoxic normoglycemic) (n=3): ***p<0.001 vs. NC; #p<0.05 vs. OGD [NC: normal control; OGD: oxygen and glucose deprivation (4 h); OGD-SN: oxygen and glucose deprivation (4 h) co-treated with SN (100 μM); SN: sodium nitrite (100 μM) treated].

Cumulative evidence has confirmed that ischemia induces ROS overproduction. ROS production initiates several intracellular signaling cascades and concomitantly induces

mitochondrial dysfunction and ER stress, leading to apoptotic cell death (Hildeman et al., 2003; Ray et al., 2012; Tajiri et al., 2004). Similar to previous reports, the current results showed that OGD resulted in excessive ROS production, although SN administration attenuated ROS levels in PC12 cells exposed to OGD for 4 h and was associated with increased cell viability. Similar results have been obtained in several studies that have shown a decrease in ROS and scavenging by NO in cerebral cells, myocardial cells and gastric mucosal cells (Kwiecien et al., 2008; Pluta et al., 2001; Wink et al., 1993; Yannopoulos et al., 2011). It is believed that the neuroprotective effect of nitrite is mediated through modulation of ROS generation following ischemia and reperfusion (Ansari et al., 2018; Raat et al., 2009).

Prolonged ischemia disrupts the ER homeostasis that triggers cellular stress responses, including the UP response as well as the increase in p-PERK and expression of ATF6 and CHOP proteins (Feldman et al., 2005; Ohoka et al., 2005). Several studies have demonstrated that ischemia induces a UP response and subsequently activates PERK through phosphorylation (Badiola et al., 2011; Bodalia et al., 2013; Cazanave et al., 2010; Hu et al., 2017; Kumar et al., 2001; Montie et al., 2005; Vavilis et al., 2016). Similarly, we found that 4 h OGD insult considerably increased ROS levels in the PC12 cells and was accompanied by an increase in ER stress markers. PERK is a type-I ER transmembrane protein with serine/threonine kinase activity in its C-terminal cytosolic domain (Bertolotti et al., 2000). Once ER stress is prolonged or strong, activated PERK phosphorylates eIF2 α , which in turn increases ATF6 expression (Harding et al., 1999, 2000). Subsequently, ATF6 activates the transcription of genes involved in functional UPR and ER stress-induced apoptosis, such as CHOP (Ameri and Harris, 2008; Harding et al., 2003; Lange et al., 2008).

ATF6 (90 kDa) is a type-II ER transmembrane protein (Haze et al., 2001). In response to the accumulation of misfolded proteins in

the ER, immunoglobulin-binding protein (BiP) detaches from p90ATF6, leading to the interaction of p90ATF6 with misfolded proteins (Shen et al., 2005) which ultimately results in translocation of p90ATF6 from the ER membrane to the Golgi (Chen et al., 2002) and produces p50ATF6. As a result, p50ATF6 translocates to the nucleus and acts as a stimulus to increase the expression of a number of genes having protein products that participate in protein folding and protein secretion, thereby supporting the cell's effort to manage ER stress and accumulation of misfolded/unfolded proteins (Adachi et al., 2008; Dorner et al., 1990; Gething and Sambrook, 1992; Haze et al., 1999; Healy et al., 2009; Li et al., 2000; Parmar and Schroder, 2012; Yamamoto et al., 2007; Yoshida et al., 2000, 2001). In the present study, a strong decrease in ATF6 expression was observed. We used p90ATF6 antibody for determining the activity of this marker by Western blotting. The results showed that, in the OGD condition, the quantity of p90ATF6 decreased, suggesting that OGD increases the cleavage of p90ATF6 to p50ATF6. Nevertheless, in our study, SN reversed these changes, suggesting that SN attenuated the protein entry from the ER to the Golgi apparatus and restored neuronal homeostasis during OGD-induced ER stress.

CHOP is a downstream product of the PERK-eIF2 α -ATF6 pathway which induces apoptosis under ER stress through down-regulation of anti-apoptotic factor such as Bcl-2, up-regulation of pro-apoptotic factors and ROS production (Oida et al., 2008; Tajiri et al., 2004; Zhao et al., 2005). It has been shown that CHOP suppression protects astrocytes from OGD injury, whereas CHOP overexpression results in astrocyte apoptosis (Osada et al., 2010). Our results demonstrate that 4 h OGD insult up-regulated the expression of CHOP, possibly through the PERK-eIF2 α -ATF6 pathways. Moreover, overexpression of CHOP was associated with an increase in the cleavage of caspase-12 and caspase-3. According to previous studies, the caspase-4 dependent pathway and mitochondrial-dependent apoptotic pathways also are

activated by ER stress, which leads to activation of caspases-12 and caspase-9 to activate caspase-3, an initiator of the apoptosis (Arduino et al., 2009; Oyadomari and Mori, 2004).

SN treatment significantly decreased the expression of CHOP in PC12 cells. Our findings suggest that the decrease in phosphorylation of PERK, as well as down-regulation of CHOP by SN treatment, may change the balance in favor of cell survival. SN treatment through down-regulation of the extrinsic apoptotic pathway, caspase-12 and -3 increased resistance to OGD-induced apoptosis. Several investigations also have confirmed that NO directly and indirectly attenuated caspase activity in cells (Ansari et al., 2018; Kim et al., 1997, 2002).

Nitrite, a NO donor, can be reduced to NO under low oxygen and pH conditions. Studies have shown that, after OGD, the fall in mitochondrial pH provides appropriate conditions for the reduction of nitrite to NO (Butler and Ridd, 2004). Nitrite-derived NO can prevent neuronal nuclear translocation of NF- κ B, an apoptosis-related transcription factor, resulting in the down-regulation of NF- κ B target genes that are generally involved in inflammatory response and apoptotic pathways (Miller and Megson, 2007; Mo et al., 2016).

ER, the major internal Ca^{2+} storage organelle, acts as a Ca^{2+} buffer and controls Ca^{2+} -dependent signaling in the cytosol (Morishima et al., 2002). Activation of the UP response and overproduction of ROS resulting from OGD disrupted the intracellular Ca^{2+} homeostasis, leading to activation of calpains. Calpains, in turn, activates the caspases, particularly pro-caspase-12, promoting cell death (Nakagawa and Yuan, 2000). The results also demonstrate that OGD-induced ROS production induced ER stress markers as well as disrupting the ER membrane and triggered Ca^{2+} release from the ER in PC12 injured cells, causing cleavage of procaspase-12 and procaspase-3 and cell death, as confirmed by the low cell viability. It should be noted that co-treatment of SN with OGD insult decreased Ca^{2+} release. The inhibitory effects of NO on

ER Ca^{2+} release have been demonstrated in skeletal muscle fibers, myocytes and small mesenteric arteries. NO can also modulate agonist-evoked intracellular Ca^{2+} release in neurosecretory PC12 cells (Clementi et al., 1995; Pouvreau and Jacquemond, 2005; Pucovsky et al., 2002).

Our findings reveal that treatment with 100 μmol SN attenuated OGD-induced ROS production, Ca^{2+} release and subsequently decreased ER stress-related markers such as p-PERK, ATF6, and CHOP expression as well as caspase-12 and caspase-3 activation. These results were confirmed by the increased cell viability as indicated by MTT assay. Some studies have reported contradictory results. For instance, it has been reported that a 1 mM dose of sodium nitroprusside induced ER stress and increased expression of CHOP as well as apoptosis in the chondrocytes (Takada et al., 2013). Oyadomari et al. (2001) found that 0.5 mM S-nitroso-N-acetyl-D,L-penicillamine (SNAP) induces ER stress and CHOP expression in pancreatic β cells. Uchiyama et al. (2002) also reported that a high concentration (1 mM) of diethylenetriamine NO or SNAP provoked both apoptosis and necrosis through activation of caspase-3 in the cardiomyocytes of neonatal rats. Furthermore, NO increases the activity of ryanodine receptors 1 and 2 (RyR1 and RyR2), a class of intracellular Ca^{2+} channels, in excitable tissues such as muscles and neurons through S-nitrosylation (Xu et al., 1998). These discrepancies in the protective or toxic effects of NO donors are possibly due to differences in NO donor type, treatment concentration or doses, time of release, cell type and treatment conditions, as many researchers have reported that the dose of the NO donor plays an important role in its cytoprotective effect (Khan et al., 2005, 2006; Takuma et al., 2002; Wedgwood and Black, 2003).

Overall, our findings demonstrated that OGD decreased cell viability and induced cell death in PC12 cells which were mediated by activation of ROS generation and induction of ER stress. However, SN as a NO donor at a low concentration (100 μM), by reducing

ROS generation, decreased expression of ER stress markers and increased cell viability in OGD injured PC12 (Figure 6). Our findings may provide a rationale for further research and the development of the use of SN as a NO donor against ischemic insult.




Figure 6: Oxygen and glucose deprivation induced ROS overproduction, leading to ER stress and cell death

Conflict of interest

All authors declare no conflict of interest.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

REFERENCES

- Adachi Y, Yamamoto K, Okada T, Yoshida H, Harada A, Mori K. ATF6 is a transcription factor specializing in the regulation of quality control proteins in the endoplasmic reticulum. *Cell Struct Funct.* 2008;33:75-89.
- Ameri K, Harris AL. Activating transcription factor 4. *Int J Biochem Cell Biol.* 2008;40:14-21.
- Anderson CA, Arciniegas DB. Cognitive sequelae of hypoxic-ischemic brain injury: a review. *NeuroRehabilitation.* 2010;26:47-63.
- Anelli T, Sitia R. Protein quality control in the early secretory pathway. *EMBO J.* 2008;27:315-27.
- Ankarcrona M, Dypbukt JM, Bonfoco E, Zhivotovsky B, Orrenius S, Lipton SA, et al. Glutamate-induced neuronal death: a succession of necrosis or apoptosis depending on mitochondrial function. *Neuron.* 1995;15:961-73.
- Ansari MHK, Karimi P, Shakib N, Beyrami SM. The neuroprotective effect of sodium nitrite on ischemic stroke-induced mitochondrial dysfunction via down-regulation of intrinsic apoptosis pathway. *Crescent J Med Biol Sci.* 2018;5:50-6.
- Arduino DM, Esteves AR, Cardoso SM, Oliveira CR. Endoplasmic reticulum and mitochondria interplay mediates apoptotic cell death: Relevance to Parkinson's disease. *Neurochem Int.* 2009;55:341-8.
- Badiola N, Penas C, Minano-Molina A, Barneda-Zahonero B, Fado R, Sanchez-Opazo G, et al. Induction of ER stress in response to oxygen-glucose deprivation of cortical cultures involves the activation of the PERK and IRE-1 pathways and of caspase-12. *Cell Death Dis.* 2011;2:e149.
- Bertolotti A, Zhang Y, Hendershot LM, Harding HP, Ron D. Dynamic interaction of BiP and ER stress transducers in the unfolded-protein response. *Nat Cell Biol.* 2000;2:326-32.
- Bodalia A, Li H, Jackson MF. Loss of endoplasmic reticulum Ca²⁺ homeostasis: contribution to neuronal cell death during cerebral ischemia. *Acta Pharmacol Sin.* 2013;34:49-59.
- Butler AR, Ridd JH. Formation of nitric oxide from nitrous acid in ischemic tissue and skin. *Nitric Oxide.* 2004;10:20-4.
- Calvert JW, Lefer DJ. Clinical translation of nitrite therapy for cardiovascular diseases. *Nitric Oxide.* 2010;22:91-7.
- Carmeliet P, Dor Y, Herbert JM, Fukumura D, Brusselmans K, Dewerchin M, et al. Role of HIF-1 α in hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis. *Nature.* 1998;394:485-90.
- Cazanave SC, Elmi NA, Akazawa Y, Bronk SF, Mott JL, Gores GJ. CHOP and AP-1 cooperatively mediate PUMA expression during lipoapoptosis. *Am J Physiol Gastrointest Liver Physiol.* 2010;299:G236-43.
- Chang TY, Chang CC, Ohgami N, Yamauchi Y. Cholesterol sensing, trafficking, and esterification. *Annu Rev Cell Dev Biol.* 2006;22:129-57.
- Chen X, Shen J, Prywes R. The luminal domain of ATF6 senses endoplasmic reticulum (ER) stress and causes translocation of ATF6 from the ER to the Golgi. *J Biol Chem.* 2002;277:13045-52.

- Clementi E, Vecchio I, Sciorati C, Nistico G. Nitric oxide modulation of agonist-evoked intracellular Ca²⁺ release in neurosecretory PC-12 cells: inhibition of phospholipase C activity via cyclic GMP-dependent protein kinase I. *Mol Pharmacol.* 1995;47:517-24.
- DeGracia DJ, Kumar R, Owen CR, Krause GS, White BC. Molecular pathways of protein synthesis inhibition during brain reperfusion: implications for neuronal survival or death. *J Cereb Blood Flow Metab.* 2002;22:127-41.
- Dorner AJ, Wasley LC, Raney P, Haugejorden S, Green M, Kaufman RJ. The stress response in Chinese hamster ovary cells. Regulation of ERp72 and protein disulfide isomerase expression and secretion. *J Biol Chem.* 1990;265:22029-34.
- Doutheil J, Gissel C, Oschlies U, Hossmann KA, Paschen W. Relation of neuronal endoplasmic reticulum calcium homeostasis to ribosomal aggregation and protein synthesis: implications for stress-induced suppression of protein synthesis. *Brain Res.* 1997;775:43-51.
- Farajdokht F, Mohaddes G, Karimi-Sales E, Kafshdooz T, Mahmoudi J, Aberoumandi SM, et al. Inhibition of PTEN protects PC12 cells against oxygen-glucose deprivation induced cell death through mitoprotection. *Brain Res.* 2018;1692:100-9.
- Feldman DE, Chauhan V, Koong AC. The unfolded protein response: a novel component of the hypoxic stress response in tumors. *Mol Cancer Res.* 2005;3:597-605.
- Ferri KF, Kroemer G. Organelle-specific initiation of cell death pathways. *Nat Cell Biol.* 2001;3:E255-63.
- Gething M-J, Sambrook J. Protein folding in the cell. *Nature.* 1992;355:33-45.
- Hampton RY. ER-associated degradation in protein quality control and cellular regulation. *Curr Opin Cell Biol.* 2002;14:476-82.
- Harding HP, Zhang Y, Ron D. Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase. *Nature.* 1999;397:271-4.
- Harding HP, Zhang Y, Bertolotti A, Zeng H, Ron D. Perk is essential for translational regulation and cell survival during the unfolded protein response. *Mol Cell.* 2000;5:897-904.
- Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, et al. An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell.* 2003;11:619-33.
- Haze K, Yoshida H, Yanagi H, Yura T, Mori K. Mammalian transcription factor ATF6 is synthesized as a transmembrane protein and activated by proteolysis in response to endoplasmic reticulum stress. *Mol Biol Cell.* 1999;10:3787-99.
- Haze K, Okada T, Yoshida H, Yanagi H, Yura T, Negishi M, et al. Identification of the G13 (cAMP-response-element-binding protein-related protein) gene product related to activating transcription factor 6 as a transcriptional activator of the mammalian unfolded protein response. *Biochem J.* 2001;355:19-28.
- Healy SJ, Gorman AM, Mousavi-Shafaei P, Gupta S, Samali A. Targeting the endoplasmic reticulum-stress response as an anticancer strategy. *Eur J Pharmacol.* 2009;625:234-46.
- Herrmann AG, Deighton RF, Le Bihan T, McCulloch MC, Searcy JL, Kerr LE, et al. Adaptive changes in the neuronal proteome: mitochondrial energy production, endoplasmic reticulum stress, and ribosomal dysfunction in the cellular response to metabolic stress. *J Cereb Blood Flow Metab.* 2013;33:673-83.
- Hildeman DA, Mitchell T, Kappler J, Marrack P. T cell apoptosis and reactive oxygen species. *J Clin Invest.* 2003;111:575-81.
- Hu YQ, Chen W, Yan MH, Lai JJ, Tang N, Wu L. Ischemic preconditioning protects brain from ischemia/reperfusion injury by attenuating endoplasmic reticulum stress-induced apoptosis through PERK pathway. *Eur Rev Med Pharmacol Sci.* 2017;21:5736-44.
- Jin K, Mao XO, Eshoo MW, Nagayama T, Minami M, Simon RP, et al. Microarray analysis of hippocampal gene expression in global cerebral ischemia. *Ann Neurol.* 2001;50:93-103.
- Kaufman RJ. Stress signaling from the lumen of the endoplasmic reticulum: coordination of gene transcriptional and translational controls. *Genes Dev.* 1999;13:1211-33.
- Kaufman RJ, Scheuner D, Schroder M, Shen X, Lee K, Liu CY, et al. The unfolded protein response in nutrient sensing and differentiation. *Nat Rev Mol Cell Biol.* 2002;3:411-21.
- Khan M, Sekhon B, Giri S, Jatana M, Gilg AG, Ayasolla K, et al. S-Nitrosoglutathione reduces inflammation and protects brain against focal cerebral ischemia in a rat model of experimental stroke. *J Cereb Blood Flow Metab.* 2005;25:177-92.
- Khan M, Jatana M, Elango C, Singh Paintlia A, Singh AK, Singh I. Cerebrovascular protection by various nitric oxide donors in rats after experimental stroke. *Nitric Oxide.* 2006;15:114-24.

- Kim PK, Kwon YG, Chung HT, Kim YM. Regulation of caspases by nitric oxide. *Ann N Y Acad Sci.* 2002; 962:42-52.
- Kim YM, Talanian RV, Billiar TR. Nitric oxide inhibits apoptosis by preventing increases in caspase-3-like activity via two distinct mechanisms. *J Biol Chem.* 1997;272:31138-48.
- Kumar R, Azam S, Sullivan JM, Owen C, Cavener DR, Zhang P, et al. Brain ischemia and reperfusion activates the eukaryotic initiation factor 2alpha kinase, PERK. *J Neurochem.* 2001;77:1418-21.
- Kwiecien S, Pawlik MW, Brzozowski T, Konturek PC, Sliwowski Z, Pawlik WW, et al. Nitric oxide (NO)-releasing aspirin and (NO) donors in protection of gastric mucosa against stress. *J Physiol Pharmacol* 2008;59 (Suppl 2):103-15.
- Lange PS, Chavez JC, Pinto JT, Coppola G, Sun C-W, Townes TM, et al. ATF4 is an oxidative stress-inducible, prodeath transcription factor in neurons in vitro and in vivo. *J Exp Med.* 2008;205:1227-42.
- Li M, Baumeister P, Roy B, Phan T, Foti D, Luo S, et al. ATF6 as a transcription activator of the endoplasmic reticulum stress element: thapsigargin stress-induced changes and synergistic interactions with NF-Y and YY1. *Mol Cell Biol.* 2000;20:5096-106.
- Lipton SA, Rosenberg PA. Excitatory amino acids as a final common pathway for neurologic disorders. *N Engl J Med.* 1994;330:613-22.
- Ma Y, Hendershot LM. ER chaperone functions during normal and stress conditions. *J Chem Neuroanat.* 2004; 28:51-65.
- Martinez JA, Zhang Z, Svetlov SI, Hayes RL, Wang KK, Larner SF. Calpain and caspase processing of caspase-12 contribute to the ER stress-induced cell death pathway in differentiated PC12 cells. *Apoptosis.* 2010;15:1480-93.
- Miller M, Megson I. Recent developments in nitric oxide donor drugs. *Br J Pharmacol.* 2007;151:305-21.
- Mo Z-t, Li W-n, Zhai Y-r, Gong Q-h. Icaritin attenuates OGD/R-induced autophagy via Bcl-2-dependent cross talk between apoptosis and autophagy in PC12 cells. *Evid Based Complement Alternat Med.* 2016;2016: 4343084.
- Montie HL, Kayali F, Haezebrouck AJ, Rossi NF, Degracia DJ. Renal ischemia and reperfusion activates the eIF 2 alpha kinase PERK. *Biochim Biophys Acta.* 2005;1741:314-24.
- Mori K. Tripartite management of unfolded proteins in the endoplasmic reticulum. *Cell.* 2000;101:451-4.
- Morishima N, Nakanishi K, Takenouchi H, Shibata T, Yasuhiko Y. An endoplasmic reticulum stress-specific caspase cascade in apoptosis. Cytochrome c-independent activation of caspase-9 by caspase-12. *J Biol Chem.* 2002;277:34287-94.
- Moskowitz MA, Lo EH, Iadecola C. The science of stroke: mechanisms in search of treatments. *Neuron.* 2010;67:181-98.
- Nakagawa T, Yuan J. Cross-talk between two cysteine protease families: activation of caspase-12 by calpain in apoptosis. *J Cell Biol.* 2000;150:887-94.
- Nakagawa T, Zhu H, Morishima N, Li E, Xu J, Yankner BA, et al. Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature.* 2000;403:98-103.
- Nakka VP, Gusain A, Raghurir R. Endoplasmic reticulum stress plays critical role in brain damage after cerebral ischemia/reperfusion in rats. *Neurotox Res.* 2010; 17:189-202.
- Ohoka N, Yoshii S, Hattori T, Onozaki K, Hayashi H. TRB3, a novel ER stress-inducible gene, is induced via ATF4-CHOP pathway and is involved in cell death. *EMBO J.* 2005;24:1243-55.
- Oida Y, Shimazawa M, Imaizumi K, Hara H. Involvement of endoplasmic reticulum stress in the neuronal death induced by transient forebrain ischemia in gerbil. *Neuroscience.* 2008;151:111-9.
- Osada N, Kosuge Y, Ishige K, Ito Y. Characterization of neuronal and astroglial responses to ER stress in the hippocampal CA1 area in mice following transient forebrain ischemia. *Neurochem Int.* 2010;57:1-7.
- Oyadomari S, Mori M. Roles of CHOP/GADD153 in endoplasmic reticulum stress. *Cell Death Differ.* 2004; 11:381-9.
- Oyadomari S, Takeda K, Takiguchi M, Gotoh T, Matsumoto M, Wada I, et al. Nitric oxide-induced apoptosis in pancreatic β cells is mediated by the endoplasmic reticulum stress pathway. *Proc Natl Acad Sci U S A.* 2001;98:10845-50.
- Oyadomari S, Araki E, Mori M. Endoplasmic reticulum stress-mediated apoptosis in pancreatic beta-cells. *Apoptosis.* 2002;7 335-45.
- Parmar VM, Schroder M. Sensing endoplasmic reticulum stress. *Adv Exp Med Biol.* 2012;738:153-68.
- Paschen W, Mengesdorf T. Endoplasmic reticulum stress response and neurodegeneration. *Cell Calcium.* 2005;38:409-15.

- Paschen W, Gissel C, Linden T, Althausen S, Doutheil J. Activation of gadd153 expression through transient cerebral ischemia: evidence that ischemia causes endoplasmic reticulum dysfunction. *Mol Brain Res.* 1998;60:115-22.
- Pellegrini-Giampietro DE, Gorter JA, Bennett MV, Zukin RS. The GluR2 (GluR-B) hypothesis: Ca(2+)-permeable AMPA receptors in neurological disorders. *Trends Neurosci.* 1997;20:464-70.
- Pluta RM, Rak R, Wink DA, Woodward JJ, Khaldi A, Oldfield EH, et al. Effects of nitric oxide on reactive oxygen species production and infarction size after brain reperfusion injury. *Neurosurgery.* 2001;48:884-92; discussion 892-3.
- Pouvreau S, Jacquemond V. Nitric oxide synthase inhibition affects sarcoplasmic reticulum Ca²⁺ release in skeletal muscle fibres from mouse. *J Physiol.* 2005;567:815-28.
- Pucovsky V, Gordienko DV, Bolton TB. Effect of nitric oxide donors and noradrenaline on Ca²⁺ release sites and global intracellular Ca²⁺ in myocytes from guinea-pig small mesenteric arteries. *J Physiol.* 2002;539:25-39.
- Qi X, Zhou R, Liu Y, Wang J, Zhang W-N, Tan H-R, et al. Trans-cinnamaldehyde protected PC12 cells against oxygen and glucose deprivation/reperfusion (OGD/R)-induced injury via anti-apoptosis and anti-oxidative stress. *Mol Cell Biochem.* 2016;421:67-74.
- Raat NJH, Shiva S, Gladwin MT. Effects of nitrite on modulating ROS generation following ischemia and reperfusion. *Adv Drug Deliv Rev.* 2009;61:339-50.
- Rao RV, Hermel E, Castro-Obregon S, del Rio G, Ellerby LM, Ellerby HM, et al. Coupling endoplasmic reticulum stress to the cell death program. Mechanism of caspase activation. *J Biol Chem.* 2001;276:33869-74.
- Ray PD, Huang BW, Tsuji Y. Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cell Signal.* 2012;24:981-90.
- Schroder M, Kaufman RJ. The mammalian unfolded protein response. *Annu Rev Biochem.* 2005;74:739-89.
- Shen J, Snapp EL, Lippincott-Schwartz J, Prywes R. Stable binding of ATF6 to BiP in the endoplasmic reticulum stress response. *Mol Cell Biol.* 2005;25:921-32.
- Steiner DR, Gonzalez NC, Wood JG. Interaction between reactive oxygen species and nitric oxide in the microvascular response to systemic hypoxia. *J Appl Physiol.* 2002;93:1411-8.
- Szegezdi E, Logue SE, Gorman AM, Samali A. Mediators of endoplasmic reticulum stress-induced apoptosis. *EMBO Rep.* 2006;7:880-5.
- Tajiri S, Oyadomari S, Yano S, Morioka M, Gotoh T, Hamada JJ, et al. Ischemia-induced neuronal cell death is mediated by the endoplasmic reticulum stress pathway involving CHOP. *Cell Death Differ.* 2004;11:403-15.
- Takada K, Hirose JUN, Yamabe S, Uehara Y, Mizuta H. Endoplasmic reticulum stress mediates nitric oxide-induced chondrocyte apoptosis. *Biomed Rep.* 2013;1:315-9.
- Takuma K, Phuapthong P, Lee E, Enomoto R, Mori K, Baba A, et al. The nitric oxide donor NOC12 protects cultured astrocytes against apoptosis via a cGMP-dependent mechanism. *Jpn J Pharmacol.* 2002;89:64-71.
- Uchiyama T, Otani H, Okada T, Ninomiya H, Kido M, Imamura H, et al. Nitric oxide induces caspase-dependent apoptosis and necrosis in neonatal rat cardiomyocytes. *J Mol Cell Cardiol.* 2002;34:1049-61.
- Vavilis T, Delivanoglou N, Aggelidou E, Stamoula E, Mellidis K, Kaidoglou A, et al. Oxygen-Glucose Deprivation (OGD) modulates the Unfolded Protein Response (UPR) and inflicts autophagy in a PC12 hypoxia cell line model. *Cell Mol Neurobiol.* 2016;36:701-12.
- Wang XZ, Harding HP, Zhang Y, Jolicoeur EM, Kuroda M, Ron D. Cloning of mammalian Ire1 reveals diversity in the ER stress responses. *EMBO J.* 1998;17:5708-17.
- Wedgwood S, Black SM. Molecular mechanisms of nitric oxide-induced growth arrest and apoptosis in fetal pulmonary arterial smooth muscle cells. *Nitric Oxide.* 2003;9:201-10.
- Wink DA, Hanbauer I, Krishna MC, DeGraff W, Gamson J, Mitchell JB. Nitric oxide protects against cellular damage and cytotoxicity from reactive oxygen species. *Proc Natl Acad Sci U S A.* 1993;90:9813-7.
- Xu L, Eu JP, Meissner G, Stamler JS. Activation of the cardiac calcium release channel (ryanodine receptor) by poly-S-nitrosylation. *Science.* 1998;279:234-7.
- Yamamoto K, Sato T, Matsui T, Sato M, Okada T, Yoshida H, et al. Transcriptional induction of mammalian ER quality control proteins is mediated by single or combined action of ATF6alpha and XBP1. *Dev Cell.* 2007;13:365-76.

Yannopoulos D, Matsuura T, Schultz J, Rudser K, Halperin HR, Lurie KG. Sodium nitroprusside enhanced cardiopulmonary resuscitation improves survival with good neurological function in a porcine model of prolonged cardiac arrest. *Crit Care Med.* 2011;39:1269-74.

Yoshida H, Okada T, Haze K, Yanagi H, Yura T, Negishi M, et al. ATF6 activated by proteolysis binds in the presence of NF-Y (CBF) directly to the cis-acting element responsible for the mammalian unfolded protein response. *Mol Cell Biol.* 2000;20:6755-67.

Yoshida H, Matsui T, Yamamoto A, Okada T, Mori K. XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell.* 2001;107:881-91.

Zhao H, Yenari MA, Cheng D, Sapolsky RM, Steinberg GK. Biphasic cytochrome c release after transient global ischemia and its inhibition by hypothermia. *J Cereb Blood Flow Metab.* 2005;25:1119-29.