

CORRECTION

## Correction: Serum- and Glucocorticoid-Inducible Kinase-1 (SGK-1) Plays a Role in Membrane Trafficking in Caenorhabditis elegans

Ming Zhu, Gang Wu, Yu-Xin Li, Julia Kathrin Stevens, Chao-Xuan Fan, Anne Spang, Meng-Qiu Dong

Fig 6B appears incorrectly as a duplicate of Fig 6C in the published article. Please see the correct Fig 6 and its legend here.



## G OPEN ACCESS

Citation: Zhu M, Wu G, Li Y-X, Stevens JK, Fan C-X, Spang A, et al. (2016) Correction: Serum- and Glucocorticoid-Inducible Kinase-1 (SGK-1) Plays a Role in Membrane Trafficking in Caenorhabditis elegans. PLoS ONE 11(9): e0163398. doi:10.1371/journal.pone.0163398

Published: September 15, 2016

Copyright: © 2016 Zhu et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.






Fig 6. sgk-1 mutant worms treated with cgt-1/3 double RNAi developed very slowly. Images of wild type (A, B, C) and sgk-1(ok538) mutant animals (D, E, F) treated with control RNAi (A, D) and cgt-1/3 double RNAi from Ahringer library (B, E) and RCE1181 library (C, F). The experiments started with 5 gravid adult worms on each plate and the plates were imaged 5 days later.

doi:10.1371/journal.pone.0163398.g001

## Reference

 Zhu M, Wu G, Li Y-X, Stevens JK, Fan C-X, Spang A, et al. (2015) Serum- and Glucocorticoid-Inducible Kinase-1 (SGK-1) Plays a Role in Membrane Trafficking in *Caenorhabditis elegans*. PLoS ONE 10(6): e0130778. doi: 10.1371/journal.pone.0130778 PMID: 26115433