

MEETING ABSTRACT

Open Access

# The role of CB<sub>2</sub> receptor ligands in human eosinophil function

Robert Frei, Eva Sturm, Ákos Heinemann\*

From 18th Scientific Symposium of the Austrian Pharmacological Society (APHAR). Joint meeting with the Croatian, Serbian and Slovenian Pharmacological Societies. Graz, Austria. 20-21 September 2012

## Background

Eosinophils play a key role in allergic diseases such as bronchial asthma and atopic dermatitis. A prominent feature of these diseases is the accumulation of eosinophils in inflamed tissue induced by several chemoattractants like prostaglandin (PG) D<sub>2</sub> or eotaxins. After the discovery of the endocannabinoid system and investigation of several endogenous and synthetic ligands, evidence has accumulated that cannabinoids, especially CB<sub>2</sub> receptor ligands, may play a major role in mediating inflammatory responses. Elevated levels of 2-arachidonoylglycerol (2-AG; a CB<sub>1</sub>/CB<sub>2</sub> agonist) were found in tissues of mouse models of allergic inflammation, suggesting a possible involvement in leukocyte recruitment.

## Methods

Blood was sampled from healthy volunteers, erythrocytes were removed by dextran sedimentation and polymorphonuclear leukocytes were obtained via Histopaque gradients. For all assays eosinophils were further purified by negative magnetic isolation. Shape change was recorded immediately after stimulation on a FACSCalibur flow cytometer. For chemotaxis assays an AP48 microBoyden chamber was used and migration time was 1h at 37°C. Intracellular Ca<sup>2+</sup> levels were analyzed by flow cytometry after treating eosinophils with Fluo-3-AM for 60min at room temperature.

## Results

We found that CB<sub>2</sub> receptor agonists like the endocannabinoid 2-AG or the synthetic selective agonist JWH-133 significantly increased eosinophil responses in shape change assays induced by PGD<sub>2</sub> or eotaxin-1. The

observed effects could be abolished by pretreatment with the selective CB<sub>2</sub> receptor antagonists SR144528 or AM630. As cytoskeletal changes are required for firm arrest and leukocyte diapedesis, transmigration assays were conducted subsequently, confirming the shape change data as eosinophil migration induced by PGD<sub>2</sub> was increased by pretreatment with JWH-133. Calcium flux assays showed Ca<sup>2+</sup> mobilization induced by JWH-133 and the tendency to increase PGD<sub>2</sub>-induced Ca<sup>2+</sup> release.

## Conclusions

We could show that JWH-133 can influence human eosinophil activation/migration via activation of CB<sub>2</sub> receptors as chemoattractant effects were significantly modulated, suggesting a possible pro-inflammatory role in allergic inflammation which may further lead to cannabinoid-based treatment options in allergic inflammatory diseases.

## Acknowledgements

This study was supported by the Jubiläumsfonds of the Austrian National Bank (OeNB, grants 14446 and 14263) and the Austrian Science Fund (FWF, grant P22521).

Published: 17 September 2012

doi:10.1186/2050-6511-13-S1-A13

Cite this article as: Frei et al.: The role of CB<sub>2</sub> receptor ligands in human eosinophil function. *BMC Pharmacology and Toxicology* 2012 **13**(Suppl 1):A13.

\* Correspondence: akos.heinemann@medunigraz.at  
Institute of Experimental and Clinical Pharmacology, Medical University of Graz, 8010 Graz, Austria