



REVIEW

Recent advances in understanding synaptic abnormalities in Rett syndrome [version 1; referees: 2 approved]

Michael Johnston¹⁻⁴, Mary E. Blue^{1,2,5}, Sakkubai Naidu^{2,6}

¹Developmental Neuroscience Laboratory, Kennedy Krieger Institute, Johns Hopkins University School of Medicine, Baltimore, MD, USA

²Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD, USA

³Department of Pediatrics, Johns Hopkins University School of Medicine, Baltimore, USA

⁴Department of Physical Medicine and Rehabilitation, Johns Hopkins University School of Medicine, Baltimore, MD, USA

⁵Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD, USA

⁶Department of Neurogenetics, Kennedy Krieger Institute, Johns Hopkins University School of Medicine, Baltimore, MD, USA

v1 First published: 22 Dec 2015, 4(F1000 Faculty Rev):1490 (doi: 10.12688/f1000research.6987.1)

Latest published: 22 Dec 2015, 4(F1000 Faculty Rev):1490 (doi: 10.12688/f1000research.6987.1)

Abstract

Rett syndrome is an extremely disabling X-linked nervous system disorder that mainly affects girls in early childhood and causes autism-like behavior, severe intellectual disability, seizures, sleep disturbances, autonomic instability, and other disorders due to mutations in the MeCP2 (methyl CpG-binding protein 2) transcription factor. The disorder targets synapses and synaptic plasticity and has been shown to disrupt the balance between glutamate excitatory synapses and GABAergic inhibitory synapses. In fact, it can be argued that Rett syndrome is primarily a disorder of synaptic plasticity and that agents that can correct this imbalance may have beneficial effects on brain development. This review briefly summarizes the link between disrupted synaptic plasticity mechanisms and Rett syndrome and early clinical trials that aim to target these abnormalities to improve the outcome for these severely disabled children.



This article is included in the **F1000 Faculty Reviews** channel.

Open Peer Review

Referee Status:

	Invited Referees	
	1	2
version 1 published 22 Dec 2015	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>

F1000 Faculty Reviews are commissioned from members of the prestigious F1000 Faculty. In order to make these reviews as comprehensive and accessible as possible, peer review takes place before publication; the referees are listed below, but their reports are not formally published.

- Zhi-Qi Xiong**, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences China
- Thierry Bienvenu**, AP-HP, Laboratoire de Biochimie et Génétique Moléculaire, GH Cochin-Broca-Hôtel Dieu, Paris, France and Université Paris Descartes Paris, Institut Cochin, INSERM U1016 France

Discuss this article

Comments (0)

Corresponding author: Michael Johnston (Johnston@kennedykrieger.org)

How to cite this article: Johnston M, Blue ME and Naidu S. **Recent advances in understanding synaptic abnormalities in Rett syndrome [version 1; referees: 2 approved]** *F1000Research* 2015, 4(F1000 Faculty Rev):1490 (doi: [10.12688/f1000research.6987.1](https://doi.org/10.12688/f1000research.6987.1))

Copyright: © 2015 Johnston M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Grant information: This work was supported by FDA Grant 5RO1FD004247 and NIH-NICHD U54 HD079123.

Competing interests: The authors declare that they have no competing interests.

First published: 22 Dec 2015, 4(F1000 Faculty Rev):1490 (doi: [10.12688/f1000research.6987.1](https://doi.org/10.12688/f1000research.6987.1))

Introduction

Rett syndrome (RTT) is characterized by severe neurodevelopmental delay in psychomotor development that occurs predominantly in girls and is due to mutations in the gene on the X chromosome that codes for the transcription factor methyl CpG-binding protein 2 (MECP2)^{1,2}. MECP2 is a transcription factor and pleiotropic protein that regulates the expression of numerous genes, including brain-derived neurotrophic factor (BDNF)³, and can mediate transcriptional activation, mRNA splicing, and post-translational processing of microRNAs^{4,5}. A phenotype similar to RTT has also been described in children with mutations in the CDKL5 and FOXP1 genes^{6,7}. Girls with RTT often present with features similar to autism with little or no speech development, poor contact with others, and stereotyped movements, including characteristic repetitive hand-wringing movements⁸. Girls with RTT typically undergo a postnatal regression phase during the first 3 years after birth in which hand wringing, seizures, loss of hand use, and loss of expressive language emerge^{9,10}. Clinicians who have seen patients with RTT are usually able to identify others because of their unique combination of physical and neurologic features. In addition to autistic features and hand-wringing movements, the characteristic behavioral abnormalities in RTT include severely erratic breathing while awake with periods of alternating hyperventilation, cyanosis and apnea, severe seizures and autonomic instability in heart rate and blood pressure, and changes in the color of the extremities consistent with sympathetic instability¹¹. The incidence of sudden death is increased in girls with RTT¹². Developing synapses appear to be a special target of the disorder, and head circumference is usually normal at birth but then falls behind over the first year of life so that it falls into the microcephalic range^{9,13–16}. Neuropathologic examination of brain tissue from older girls with RTT who have died has revealed that neurons are typically too close together, suggesting a stunting of axonodendritic development^{17,18}. This hypothesis is supported by studies of nasal epithelium from girls with RTT compared with controls, which showed that the neurogenesis of olfactory receptor neurons is normal but their maturation is blocked at the point of formation of synapses with the olfactory bulb¹⁹.

Role of excitatory synapse abnormalities in Rett syndrome

Early studies of patients with RTT, including some who had not been tested for mutations in the MECP2 gene, suggested an abnormality of glutamate excitatory synapses¹⁸. For example, Hamberger *et al.*²⁰ and Lappalainen and Riikonen²¹ in the 1990s reported that cerebrospinal fluid from girls with RTT had concentrations of glutamate but not other amino acids that were more than twice as high as those of children without RTT. Blue *et al.* in 1999 reported the first autoradiographic study of glutamate and GABA receptors in postmortem brain tissue from nine girls with RTT and 10 female controls of different ages²². The authors reported that densities of N-methyl-D-aspartate (NMDA) were elevated in the frontal lobe cortex in girls younger than 8 years of age but were significantly lower in girls older than 8 years without RTT²³. Similar age-related changes in glutamate receptors were observed in the basal ganglia. Horská *et al.*²⁴ used magnetic resonance spectroscopy to demonstrate that glutamate was elevated in the cerebral cortex of younger girls (less than 8 years of age) with RTT, especially those with seizures, but reduced in older girls. These changes in levels of glutamate

between younger and older girls correlate with the bi-phasic changes in the clinical condition of girls with RTT who have more clinical signs of excitatory activity, including seizures early in early childhood, but have reduced signs of excessive central nervous system activity later in childhood⁹. More recently, Blue *et al.* demonstrated similar age-related changes in the NMDA1 subtype in the Bird model of RTT in mice²⁵. Taken together, the elevations in glutamate as well as in glutamate receptors in younger girls with RTT suggest that there is an abnormality of the normal homeostatic synaptic scaling by which post-synaptic receptors adjust upwards or downwards to maintain a stable level of neurotransmission²⁶. This abnormality in synaptic function probably contributes to deficits in excitatory synaptic plasticity and enhanced excitability observed in mouse models of RTT^{27–30}.

The pathophysiology of elevated glutamate in the face of elevated NMDA glutamate receptors in RTT is not clear. The major regulator of glutamate levels in and outside the synapse is the activity of glial glutamate transporters, which normally are activated when glutamate is released into the synapse³¹. Intra-synaptic levels of glutamate are thought to provide trophic influences on post-synaptic neurons, but glutamate flooding outside the synapse can activate excitotoxicity mediated by extra-synaptic NMDA receptors³². Maezawa and Jin³³ reported that conditioned media obtained from *Mecp2*-null microglia from *Mecp2*^{tm1.1Bird/+} mice³⁴ contain toxic levels of glutamate that is damaging to neural dendrites *in vitro* and this effect can be blocked by inhibiting glutaminase or glutamate receptors. Jin *et al.*, using the same model of *Mecp2* deficiency, reported that abnormal function of the SNAT1 glutamine transporter in microglia is associated with NMDA receptor-mediated neurotoxicity, mitochondrial dysfunction, and decreased viability of microglia³⁵. Okabe *et al.*³⁶ reported that astroglia from *Mecp2*-null mice (*Mecp2*^{tm1.1Bird/+}) had downregulation of glutamate transporters that could increase extracellular glutamate to toxic levels. Liyo *et al.*³⁷ also reported that the restoration of wild-type MeCP2 in astrocytes of MeCP2-deficient mice (*Mecp2*^{Stop}) restored normal dendritic morphology and increased levels of the excitatory glutamate transporter. We also found that cerebellar granule cells from *Mecp2*-null mice (*Mecp2*^{tm1.1Bird/+}) are more sensitive to cell death from hypoxia-ischemia and glutamate excitotoxicity compared with neurons from wild-type mice, suggesting that they also have intrinsic vulnerability to cell death³⁸. These findings are consistent with the above-cited evidence of abnormal homeostatic synaptic scaling, which is expected to degrade normal excitatory neurotransmission and activity-dependent neuronal plasticity needed for learning and memory in RTT.

Abnormal diurnal regulation of sleep and brain glutamate levels in MeCP2 deficiency

Recently, the circadian sleep cycle in rodents has been associated with alterations in brain glutamate levels, with higher glutamate levels associated with wakefulness and sleep associated with a reduction in glutamate³⁹. We applied the relatively new technology that can measure 24-hour continuous electroencephalogram (EEG) markers of sleep cycles and correlated them with biosensor measurement of *in vivo* glutamate concentrations⁴⁰. We found that mice with a *Mecp2* knockout mutation (*Mecp2*^{tm1.1Bird/+}) had abnormal behaviors and remarkably abnormal sleep cycles with long periods

of sleeplessness associated with very high relative concentrations of brain glutamate. These *Mecp2*-deficient mice also had elevated baseline levels of cortical glutamate measured with a separate colorimetric method in postmortem tissue. These data suggest that periods of sleep disturbance that are common in girls with RTT could be associated with enhanced glutamate neurotoxicity.

Role of GABA-containing interneurons in Rett syndrome

GABA (gamma-amino butyric acid) is the predominant inhibitory neurotransmitter in the brain and is used as a transmitter at approximately 15% of synapses, and there is ample evidence of its altered metabolism in the mice with *Mecp2* deficiency. Chao *et al.*⁴¹ reported that the selective deletion of MeCP2 in GABA interneurons (*Viaat-Mecp2^{2/2}* mice) led to a reduction in the pre-synaptic release of GABA as well as autism-like stereotypies and RTT phenotypes in mice. Kang *et al.*⁴² reported that there is a significant location-specific downregulation of synaptic GABA transporters in *Mecp2* knockout mice (*Mecp2^{tm1.1Bird/+}*), and El-Khoury *et al.*⁴³ reported age- and region-specific reductions in function in GABAergic pathways in MeCP2-deficient mice (B6.129P2(c)-*Mecp2^{tm1-1Bird}*). These observations along with those from the glutamate system indicate that there is a shift in the balance between excitation and inhibition in favor of excitation, as has been suggested to occur in other autism spectrum disorders^{8,44}.

One of the most interesting recent observations on the role of altered development in GABA and glutamate-containing neurons in mice with MeCP2 mutations relates to the link between MeCP2 and visual cortical plasticity⁴⁵. Visual cortical plasticity associated with the occlusion of vision in one eye can be measured precisely in the laboratory in immature mice, and GABAergic interneurons play a critical role in this process⁴⁶. Durand *et al.* reported that MeCP2-null mice (*Mecp2^{tm1.1Bird/+}*) have normal visual function early in the postnatal period but that visual acuity regresses after postnatal day 35–40 and the cortex fell silent by postnatal day 55–60⁴⁷. Remarkably, this effect could be prevented by genetic deletion of the NMDA glutamate receptor subunit NR2A. He *et al.* also reported that the genetic conditional deletion of MeCP2 in GABAergic parvalbumin-expressing neurons prevented local circuit inhibitory functions required for experience-dependent visual cortical plasticity⁴⁸. Krishnan *et al.*⁴⁵ reported that the MeCP2 regulates the timing of the critical period of plasticity in the primary visual cortex. Kron *et al.*⁴⁹ have also examined brain activity maps by using activation of the immediate early gene Fos and electrophysiology and found evidence of synaptic hyperexcitability in the cortical default mode network that was reduced by the administration of the NMDA channel blocking anesthetic ketamine. The results of these studies support the themes raised earlier in this discussion of RTT, namely that MeCP2 deficiency leads to excessive excitation versus inhibition and defective activity-dependent synaptic plasticity. Although ketamine can cause damage to developing neurons by excessively blocking the activity of NMDA receptors⁵⁰, its use in RTT is intended to reduce excessive activity of NMDA receptors found in RTT to normal levels. Both too little and too much activity at NMDA receptors can damage the developing brain⁵¹.

Potential new therapies based on advances in understanding Rett syndrome

Several clinical studies are examining potential therapies for girls with RTT on the basis of the basic science insights from experiments on mice with MeCP2 deficiency. Naidu *et al.* launched a randomized open-label study of oral dextromethorphan, a competitive NMDA antagonist, for seizures and cognitive function in girls with RTT from 2008 to 2014 (ClinicalTrials.gov: NCT00593957). Results of this study suggest a significant improvement in a secondary outcome of receptive speech by using the Mullen Scales of Early Learning over the course of 6 months but no change in seizures. Based on these promising preliminary data on cognition, Naidu *et al.* are carrying out a randomized double-blind study of dextromethorphan for girls with RTT which was started in January 2012 (ClinicalTrials.gov: NCT01520363) with support from the US Food and Drug Administration, and this study is still seeking subjects for enrollment. Other blockers of the NMDA receptor, including memantine and ketamine as mentioned above, are under discussion but have not been registered in ClinicalTrials.gov.

Published data are available on the administration of an active peptide of insulin-like growth factor 1 (IGF-1), which has been reported to be deficient in mice with *Mecp2* deficiency⁵². IGF-1 activates tropomyosin receptor kinase B (TrkB) receptors and stimulates downstream signaling through the PI3K, AKT, and mTOR (mechanistic target of rapamycin) signaling pathway to stimulate protein synthesis⁵³. Recent evidence also indicates that the positive effect of *Mecp2* on mTOR is mediated by post-transcriptional processing of microRNAs that enhance mTOR activity so that *Mecp2* depletion leads to a reduction in mTOR activity⁴.

IGF-1 administration has been reported to improve behavioral functional recovery in *Mecp2*-deficient mice (*Mecp2^{lox/+}* females⁵⁴) as well as spine density, synaptic amplitude, and PSD-95, a major constituent of the synaptic post-synaptic density in excitatory synapses^{55,56}. Khwaja *et al.*⁵² recently reported a preliminary assessment of the use of recombinant human IGF-1 (mecaserman) in girls with RTT. Some behavioral parameters, including anxiety and mood and an EEG parameter, seemed to improve, and the medication appeared to be safe and well tolerated⁵². Recent preclinical data suggest that fingolimod, a sphingosine-1 phosphate receptor modulator that can increase levels of BDNF, can also reduce signs of disease in a mouse model of RTT⁵⁷. Valproic acid has also recently been reported to have a positive behavioral effect on mice with Rett mutations and phenotype⁵⁸.

Conclusions

Mice with *Mecp2* deficiency display a variety of aberrant behaviors and synaptic abnormalities that correlate very well with those in human RTT. Girls with RTT and mice with *Mecp2* deficiency display elevations in both glutamate and NMDA glutamate receptors and this is likely to produce abnormal homeostatic synaptic scaling and probably contributes to impaired activity-dependent synaptic plasticity. RTT appears to involve an imbalance between elevated glutamate levels and reduced GABA levels, similar to models of autism spectrum disorders. Chaotic sleep in mice with *Mecp2*

deficiency has been shown to correlate with elevated baseline glutamate levels in the brain and very high elevations in glutamate associated with prolonged waking. Several clinical studies are under way to attempt to normalize synaptic abnormalities in RTT, including the studies by Naidu *et al.* with dextromethorphan, a clinically approved competitive NMDA receptor blocker. A randomized un-blinded trial of dextromethorphan showed improvement in receptive language in girls with RTT, and a randomized trial of this drug is under way. Preclinical studies suggest that low-dose ketamine, a non-competitive NMDA blocker, might be useful for improving the connectivity of brain circuits affected in RTT and improving function. Human recombinant IGF-1 has also shown

benefit in mice with *Mecp2* deficiency, and a preliminary study in girls with RTT showed that it was well tolerated and may have some benefits.

Competing interests

The authors declare that they have no competing interests.

Grant information

This work was supported by FDA Grant 5RO1FD004247 and NIH-NICHD U54 HD079123.

References

- Chen L, Chen K, Lavery LA, *et al.*: **MeCP2 binds to non-CG methylated DNA as neurons mature, influencing transcription and the timing of onset for Rett syndrome.** *Proc Natl Acad Sci U S A.* 2015; **112**(17): 5509–14.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Naidu S, Johnston MV: **Neurodevelopmental disorders: Clinical criteria for Rett syndrome.** *Nat Rev Neurol.* 2011; **7**(6): 312–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chapleau CA, Larimore JL, Theibert A, *et al.*: **Modulation of dendritic spine development and plasticity by BDNF and vesicular trafficking: fundamental roles in neurodevelopmental disorders associated with mental retardation and autism.** *J Neurodev Disord.* 2009; **1**(3): 185–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Tsujimura K, Irie K, Nakashima H, *et al.*: **miR-199a Links MeCP2 with mTOR Signaling and Its Dysregulation Leads to Rett Syndrome Phenotypes.** *Cell Rep.* 2015; **12**(11): 1887–901.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nectoux J, Fichou Y, Cagnard N, *et al.*: **Cell cloning-based transcriptome analysis in cyclin-dependent kinase-like 5 mutation patients with severe epileptic encephalopathy.** *J Mol Med (Berl).* 2011; **89**(2): 193–202.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bahi-Buisson N, Nectoux J, Girard B, *et al.*: **Revisiting the phenotype associated with FOXP1 mutations: two novel cases of congenital Rett variant.** *Neurogenetics.* 2010; **11**(2): 241–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Fichou Y, Bieth E, Bahi-Buisson N, *et al.*: **Re: CDKL5 mutations in boys with severe encephalopathy and early-onset intractable epilepsy.** *Neurology.* 2009; **73**(1): 77–8; author reply 78.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Canitano R: **New experimental treatments for core social domain in autism spectrum disorders.** *Front Pediatr.* 2014; **2**: 61.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Naidu S: **Rett syndrome: a disorder affecting early brain growth.** *Ann Neurol.* 1997; **42**(1): 3–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Glaze DG: **Rett syndrome: of girls and mice—lessons for regression in autism.** *Ment Retard Dev Disabil Res Rev.* 2004; **10**(2): 154–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cooper RA, Kerr AM, Amos PM: **Rett syndrome: critical examination of clinical features, serial EEG and video-monitoring in understanding and management.** *Eur J Paediatr Neurol.* 1998; **2**(3): 127–35.
[PubMed Abstract](#) | [Publisher Full Text](#)
- de Felice C, Maffei S, Signorini C, *et al.*: **Subclinical myocardial dysfunction in Rett syndrome.** *Eur Heart J Cardiovasc Imaging.* 2012; **13**(4): 339–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Jentarra GM, Olfers SL, Rice SG, *et al.*: **Abnormalities of cell packing density and dendritic complexity in the MeCP2 A140V mouse model of Rett syndrome/X-linked mental retardation.** *BMC Neurosci.* 2010; **11**: 19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Nguyen MV, Du F, Felice CA, *et al.*: **MeCP2 is critical for maintaining mature neuronal networks and global brain anatomy during late stages of postnatal brain development and in the mature adult brain.** *J Neurosci.* 2012; **32**(29): 10021–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Landi S, Putignano E, Boggio EM, *et al.*: **The short-time structural plasticity of dendritic spines is altered in a model of Rett syndrome.** *Sci Rep.* 2011; **1**: 45.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Chapleau CA, Calfa GD, Lane MC, *et al.*: **Dendritic spine pathologies in hippocampal pyramidal neurons from Rett syndrome brain and after expression of Rett-associated MECP2 mutations.** *Neurobiol Dis.* 2009; **35**(2): 219–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Armstrong DD: **Neuropathology of Rett syndrome.** *J Child Neurol.* 2005; **20**(9): 747–53.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Johnston MV, Jeon OH, Pevsner J, *et al.*: **Neurobiology of Rett syndrome: a genetic disorder of synapse development.** *Brain Dev.* 2001; **23**(Suppl 1): S206–13.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ronnett GV, Leopold D, Cai X, *et al.*: **Olfactory biopsies demonstrate a defect in neuronal development in Rett's syndrome.** *Ann Neurol.* 2003; **54**(2): 206–18.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hamberger A, Gillberg C, Palm A, *et al.*: **Elevated CSF glutamate in Rett syndrome.** *Neuropediatrics.* 1992; **23**(4): 212–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lappalainen R, Riikonen RS: **High levels of cerebrospinal fluid glutamate in Rett syndrome.** *Pediatr Neurol.* 1996; **15**(3): 213–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Blue ME, Naidu S, Johnston MV: **Altered development of glutamate and GABA receptors in the basal ganglia of girls with Rett syndrome.** *Exp Neurol.* 1999; **156**(2): 345–52.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Blue ME, Naidu S, Johnston MV: **Development of amino acid receptors in frontal cortex from girls with Rett syndrome.** *Ann Neurol.* 1999; **45**(4): 541–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Horská A, Farage L, Bibat G, *et al.*: **Brain metabolism in Rett syndrome: age, clinical, and genotype correlations.** *Ann Neurol.* 2009; **65**(1): 90–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Blue ME, Kaufmann WE, Bressler J, *et al.*: **Temporal and regional alterations in NMDA receptor expression in *Mecp2*-null mice.** *Anat Rec (Hoboken).* 2011; **294**(10): 1624–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Blackman MP, Djukic B, Nelson SB, *et al.*: **A critical and cell-autonomous role for MeCP2 in synaptic scaling up.** *J Neurosci.* 2012; **32**(39): 13529–36.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Asaka Y, Jugloff DG, Zhang L, *et al.*: **Hippocampal synaptic plasticity is impaired in the *Mecp2*-null mouse model of Rett syndrome.** *Neurobiol Dis.* 2006; **21**(1): 217–27.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Dani VS, Chang Q, Maffei A, *et al.*: **Reduced cortical activity due to a shift in the balance between excitation and inhibition in a mouse model of Rett syndrome.** *Proc Natl Acad Sci U S A.* 2005; **102**(35): 12560–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zhang L, He J, Jugloff DG, *et al.*: **The MeCP2-null mouse hippocampus displays altered basal inhibitory rhythms and is prone to hyperexcitability.** *Hippocampus.* 2008; **18**(3): 294–309.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Maliszewska-Cyna E, Bawa D, Eubanks JH: **Diminished prevalence but preserved synaptic distribution of N-methyl-D-aspartate receptor subunits in the methyl CpG binding protein 2 (MeCP2)-null mouse brain.** *Neuroscience.* 2010; **168**(3): 624–32.
[PubMed Abstract](#) | [Publisher Full Text](#)

31. Magistretti PJ, Pellerin L: **Cellular mechanisms of brain energy metabolism and their relevance to functional brain imaging.** *Philos Trans R Soc Lond B Biol Sci.* 1999; **354**(1387): 1155–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
32. Zhou X, Hollern D, Liao J, *et al.*: **NMDA receptor-mediated excitotoxicity depends on the coactivation of synaptic and extrasynaptic receptors.** *Cell Death Dis.* 2013; **4**(3): e560.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Maezawa I, Jin LW: **Rett syndrome microglia damage dendrites and synapses by the elevated release of glutamate.** *J Neurosci.* 2010; **30**(15): 5346–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
34. Guy J, Hendrich B, Holmes M, *et al.*: **A mouse *Mecp2*-null mutation causes neurological symptoms that mimic Rett syndrome.** *Nat Genet.* 2001; **27**(3): 322–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
35. Jin LW, Horiuchi M, Wulff H, *et al.*: **Dysregulation of glutamine transporter SNAT1 in Rett syndrome microglia: a mechanism for mitochondrial dysfunction and neurotoxicity.** *J Neurosci.* 2015; **35**(6): 2516–29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Okabe Y, Takahashi T, Mitsumasu C, *et al.*: **Alterations of gene expression and glutamate clearance in astrocytes derived from an *MeCP2*-null mouse model of Rett syndrome.** *PLoS One.* 2012; **7**(4): e35354.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. Lioy DT, Garg SK, Monaghan CE, *et al.*: **A role for glia in the progression of Rett's syndrome.** *Nature.* 2011; **475**(7357): 497–500.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
38. Russell JC, Blue ME, Johnston MV, *et al.*: **Enhanced cell death in *MeCP2* null cerebellar granule neurons exposed to excitotoxicity and hypoxia.** *Neuroscience.* 2007; **150**(3): 563–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Azuma S, Kodama T, Honda K, *et al.*: **State-dependent changes of extracellular glutamate in the medial preoptic area in freely behaving rats.** *Neurosci Lett.* 1996; **214**(2–3): 179–82.
[PubMed Abstract](#) | [Publisher Full Text](#)
40. Johnston MV, Ammanuel S, O'Driscoll C, *et al.*: **Twenty-four hour quantitative-EEG and *in-vivo* glutamate biosensor detects activity and circadian rhythm dependent biomarkers of pathogenesis in *Mecp2* null mice.** *Front Syst Neurosci.* 2014; **8**: 118.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
41. Chao HT, Chen H, Samaco RC, *et al.*: **Dysfunction in GABA signalling mediates autism-like stereotypies and Rett syndrome phenotypes.** *Nature.* 2010; **468**(7321): 263–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Kang SK, Kim ST, Johnston MV, *et al.*: **Temporal- and Location-Specific Alterations of the GABA Recycling System in *Mecp2* KO Mouse Brains.** *J Cent Nerv Syst Dis.* 2014; **6**: 21–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. El-Khoury R, Panayotis N, Matagne V, *et al.*: **GABA and glutamate pathways are spatially and developmentally affected in the brain of *Mecp2*-deficient mice.** *PLoS One.* 2014; **9**(3): e92169.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
44. Nelson SB, Valakh V: **Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders.** *Neuron.* 2015; **87**(4): 684–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
45. Krishnan K, Wang BS, Lu J, *et al.*: ***MeCP2* regulates the timing of critical period plasticity that shapes functional connectivity in primary visual cortex.** *Proc Natl Acad Sci U S A.* 2015; **112**(34): E4782–91.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
46. Fagiolini M, Fritschy JM, Löw K, *et al.*: **Specific GABAA circuits for visual cortical plasticity.** *Science.* 2004; **303**(5664): 1681–3.
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Durand S, Patrizi A, Quast KB, *et al.*: **NMDA receptor regulation prevents regression of visual cortical function in the absence of *Mecp2*.** *Neuron.* 2012; **76**(6): 1078–90.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
48. He LJ, Liu N, Cheng TL, *et al.*: **Conditional deletion of *Mecp2* in parvalbumin-expressing GABAergic cells results in the absence of critical period plasticity.** *Nat Commun.* 2014; **5**: 5036.
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Kron M, Howell CJ, Adams IT, *et al.*: **Brain activity mapping in *Mecp2* mutant mice reveals functional deficits in forebrain circuits, including key nodes in the default mode network, that are reversed with ketamine treatment.** *J Neurosci.* 2012; **32**(40): 13860–72.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Slikker W Jr, Liu F, Rainosek SW, *et al.*: **Ketamine-Induced Toxicity in Neurons Differentiated from Neural Stem Cells.** *Mol Neurobiol.* 2015; **52**(2): 959–69.
[PubMed Abstract](#) | [Publisher Full Text](#)
51. McDonald JW, Johnston MV: **Physiological and pathophysiological roles of excitatory amino acids during central nervous system development.** *Brain Res Brain Res Rev.* 1990; **15**(1): 41–70.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Khwaja OS, Ho E, Barnes KV, *et al.*: **Safety, pharmacokinetics, and preliminary assessment of efficacy of mecasermin (recombinant human IGF-1) for the treatment of Rett syndrome.** *Proc Natl Acad Sci U S A.* 2014; **111**(12): 4596–601.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Banerjee A, Castro J, Sur M: **Rett syndrome: genes, synapses, circuits, and therapeutics.** *Front Psychiatry.* 2012; **3**: 34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. Chen RZ, Akbarian S, Tudor M, *et al.*: **Deficiency of methyl-CpG binding protein-2 in CNS neurons results in a Rett-like phenotype in mice.** *Nat Genet.* 2001; **27**(3): 327–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
55. Tropea D, Giacometti E, Wilson NR, *et al.*: **Partial reversal of Rett Syndrome-like symptoms in *MeCP2* mutant mice.** *Proc Natl Acad Sci U S A.* 2009; **106**(6): 2029–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Castro J, Garcia RI, Kwok S, *et al.*: **Functional recovery with recombinant human IGF1 treatment in a mouse model of Rett Syndrome.** *Proc Natl Acad Sci U S A.* 2014; **111**(27): 9941–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
57. Deogracias R, Yazdani M, Dekkers MP, *et al.*: **Fingolimod, a sphingosine-1 phosphate receptor modulator, increases BDNF levels and improves symptoms of a mouse model of Rett syndrome.** *Proc Natl Acad Sci U S A.* 2012; **109**(35): 14230–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
58. Guo W, Tsujimura K, Otsuka IM, *et al.*: **VPA alleviates neurological deficits and restores gene expression in a mouse model of Rett syndrome.** *PLoS One.* 2014; **9**(6): e100215.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Referee Status:



Editorial Note on the Review Process

F1000 Faculty Reviews are commissioned from members of the prestigious **F1000 Faculty** and are edited as a service to readers. In order to make these reviews as comprehensive and accessible as possible, the referees provide input before publication and only the final, revised version is published. The referees who approved the final version are listed with their names and affiliations but without their reports on earlier versions (any comments will already have been addressed in the published version).

The referees who approved this article are:

Version 1

- 1 Thierry Bienvenu**, AP-HP, Laboratoire de Biochimie et Génétique Moléculaire, AP-HP, Laboratoire de Biochimie et Génétique Moléculaire, GH Cochin-Broca-Hôtel Dieu, Paris, France and Université Paris Descartes Paris, Institut Cochin, INSERM U1016, Paris, France
Competing Interests: No competing interests were disclosed.
- 2 Zhi-Qi Xiong**, Institute of Neuroscience, State Key Laboratory of Neuroscience, CAS Center for Excellence in Brain Science, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai, China
Competing Interests: No competing interests were disclosed.