

# Nontraumatic spontaneous intracerebral hemorrhage: Baseline characteristics and early outcomes

Mohamed Al-Khaled  | Samer Awwad | Toralf Brüning 

Department of Neurology, University of Lübeck, Lübeck, Germany

## Correspondence

Mohamed Al-Khaled, Department of Neurology, University of Lübeck, Ratzeburger Allee 160, 23538 Lübeck, Germany.

Email: Mohamed.al-khaled@neuro.uni-luebeck.de

## Abstract

**Background and Purpose:** Hemorrhagic stroke, particularly nontraumatic spontaneous intracerebral hemorrhage (SICH), is a cerebrovascular condition with unfavorable outcomes. The aims of the present study were to evaluate patients who suffered from SICH and investigate the early outcomes in a single-center study.

**Methods:** During a study -period of 6 years (2008–2014), 613 consecutive patients (mean age,  $72 \pm 12.7$  years; 51.1% female), who suffered from nontraumatic SICH and were treated at the Department of Neurology at the University Hospital of Schleswig-Holstein, Campus Lübeck, Germany, were included and prospectively analyzed.

**Results:** During a mean hospitalization time of 12 days, 148 patients (24.1%) died, 47% of those within the first 2 days and 79% within the first week. The patients who died stayed at the hospital for a shorter time (3) than those who survived ( $p < .001$ ). In the multivariate logistic regression, following parameters were found to be associated with the in-hospital mortality: female sex (OR, 2.0; 95%-CI, 1.2–3.4;  $p = .009$ ), a NIHSS score  $> 10$  (OR, 10.5; 95%-CI, 5.6–19.5;  $p < .001$ ), history of hypertension (OR, 0.35; 95%-CI, 0.19–0.64;  $p = .001$ ), previous oral anticoagulation (OR, 2; 95%-CI, 1.0–3.8;  $p = .032$ ), and intraventricular extension of hemorrhage (OR, 2.8; 95%-CI, 1.7–4.7;  $p = .001$ ). At discharge, 192 patients (41.2%) showed favorable outcomes ( $mRS \leq 2$ ) whereas the median mRS of patients who survived was 3 (IQR 2–4). The good functional outcome at discharge from the acute hospital was decreased by an age  $> 70$  years (OR, 0.56; 95%-CI, 0.35–0.9;  $p = .017$ ), NIHSS score  $> 10$  at admission (OR, 0.07; 95%-CI, 0.04–0.13;  $p < .001$ ), and development of pneumonia during hospitalization (OR, 0.35; 95%-CI, 0.2–0.6;  $p < .001$ ).

**Conclusion:** The present study showed that SICH is a serious disease causing high mortality and disability, particularly in the early period after event.

## KEYWORDS

hemorrhage, nontraumatic spontaneous intracerebral hemorrhage, prognosis

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Brain and Behavior* published by Wiley Periodicals, Inc.

## 1 | INTRODUCTION

The hemorrhagic stroke, nontraumatic spontaneous intracerebral hemorrhage (SICH), is a heterogeneous cerebrovascular condition that leads to disability and rapid death (Al-Khaled, Eggers, & Qug, 2014; Broderick et al., 2007; Weimar et al., 2003). It represents the second form of stroke with a number of patients affected worldwide each year as high as 4 million and a median case fatality of one month of 40% (Asch et al., 2010; Ferro, 2006; Kolominsky-Rabas et al., 1998; Sacco et al., 1998; Sadamasa, Yoshida, Narumi, & Chin, 2011). In such a way, SICH is characterized as the most killing disease in the early phase after onset (Hemphill, Farrant, & Neill, 2009). Many survivors remain severely disabled with only one in four patients having a good outcome (Feigin, Lawes, Bennett, & Anderson, 2003).

In SICH, the bleeding results from cerebrovascular vessels most likely due to first diagnosed or preexisting blood hypertension and as well as a result of amyloid angiopathy (Butcher and Laidlaw, 2003). The brain regions basal ganglia (34%) followed by lobar regions (25%) are the most frequently affected sites in SICH (Hemphill, Bonovich, Besmertis, Manley, and Johnston, 2001). The SICH occurs typically in the basal region by hypertension (Figure 1), whereas it may atypically occur in other brain regions (Figure 2). The cerebral amyloid angiopathy causes SICH as well as asymptomatic microbleeds in the brain that can be detected by brain MRI (Figure 3).

Whereas an absolute benefit from surgery could have been revealed recently in patients with traumatic intracerebral hemorrhage (Gregson et al., 2015), the potential benefit of surgical compared to conservative treatment of SICH still remains controversial, a significant minor advantage was found in cases of SICH without

intraventricular extension (Kim, Lee, Koh, & Choi, 2015; Mendelow et al., 2013).

## 2 | METHODS

### 2.1 | Study design

Between 2008 and 2014, 613 patients (mean age,  $72 \pm 12.7$  years; female, 51.1%), who were suffering from SICH and were treated at the University Hospital of Schleswig-Holstein, Campus Lübeck, Germany, were included in the present prospective study. We used standardized radiological analysis in characterizing SICH.

The diagnosis of SICH was made by using cranial computed tomography (CCT) and magnetic resonance imaging of brain (cMRI).

Baseline parameters, diagnostic, and therapeutic procedures were retrieved from the medical patient's files (Table 1). The assessment of outcome was performed by the modified Rankin Scale (mRS), with a range from mRS = 0 points [no symptoms] to mRS = 6 points [death].


All deaths were assumed as neurological mortality. Favorable outcomes were set at a mRS  $\leq 2$ . The patients were evaluated at the time of admission to the discharge from the hospital.

All included patients were admitted in the semi-intensive care (stroke unit) or in the intensive care unit in cases of altered consciousness and/or when assistance of respirations was required. All patients were treated by physicians with stroke experience as well as stroke neurologists.

The present study was part of the Stroke Registry at the Department of Neurology. The study (Stroke Registry) was approved by the local ethics committee.




**FIGURE 1** Brain CT scan showed a SICH by a 45-year-old patient, firstly diagnosed with blood hypertension. The patient was presented with a weakness on the left side and hypertensive crisis



**FIGURE 2** Atypically localized sICB with ventricular extension showed by brain CT scan. A 64-year-old woman who were presented with a fast-progredient altered consciousness

**FIGURE 3** MRI scan showed cerebral amyloid angiopathy, several microbleeds in the infra- as well extratentorial region. A 67 years patients presented with increasing cognitive impairment in the last year



The patients, who were entered in this study, were exclusively treated conservatively. This conservative care was carried out in accordance with the recommendations and guidelines of the German Stroke Society and German Society of Neurology (European Stroke Initiative Writing C, 2006; Thieme Verlag, 2012). Briefly, the management of SICH was conservative and included controlling of blood pressure and other vital parameters, if necessary an intra-arterial measurement was used. The monitoring of vital parameters was essential in the care of hemorrhage patients (Qureshi et al., 2010). Furthermore, patients with respiratory insufficiency due to altered consciousness received an intubation and ventilation for airway protection, those patients were treated initially on intensive care, after weaning the treatment was continued on the stroke unit.

## 2.2 | Statistical analysis

The SPSS software (version 23; IBM) was used to analyze the data. The values of the continuous variables were reported with the mean and standard deviations, whereas the scores with the median and interquartile range (IQR). Absolute numbers and percentages were used to describe the categorical and nominal variables. A chi-square test was used to determine the correlation between categorical variables and a Mann-Whitney test between scores as well as a *t* test between continuous variables was performed. The multivariate analysis was performed to estimate the odds ratio (ORs), and all parameters with a significant association in the univariate analysis were entered in the logistic regression. A *p*-value below .05 was considered significant.

## 3 | RESULTS

In the present study, a total of 613 patients (mean age  $72 \pm 12.7$  year; 51.1% female; median NIHSS score 10 [IQR 4–18]) were included conservatively for nontraumatic SICH.

All patients have undergone at least one CCT scan, one at admission, and control CT scan during hospitalization or/and MRI scan.

During hospitalization with a mean overall duration of hospital stay of  $12 \pm 8$  days, 148 patients (24.1%) died, 47% of those died

within the first 2 days and 79% within the first week. The mean hospitalization of patients who did not survive was remarkably shorter (3 days) than that of those who survived ( $p < .001$ ).

The baseline parameters and a comparison between patients who survived vs. those who did not are shown in Table 1. The rate of early death was found to be higher in the elderly, female gender, history of hypertension, and premedication with oral anticoagulants. Furthermore, a significant association between mortality and localization as well as etiology of bleeding has been established (Table 1).

In the multivariate logistic regression, following parameters were found to be associated with the in-hospital mortality: female sex (OR, 2.0; 95%-CI, 1.2–3.4;  $p = .009$ ), a NIHSS score  $> 10$  (OR, 10.5; 95%-CI, 5.6–19.5;  $p < .001$ ), previous oral anticoagulation (OR, 2; 95%-CI, 1.0–3.8;  $p = .032$ ), and intraventricular extension of hemorrhage (OR, 2.8; 95%-CI, 1.7–4.7;  $p = .001$ ) that were associated with increased death, contrarywise a known hypertension was linked to decreased in-hospital mortality (OR, 0.35; 95%-CI, 0.19–0.64;  $p = .001$ ) in logistic regression analysis.

At discharge, 192 patients (41.2%) showed favorable outcomes ( $mRS \leq 2$ ) whereas the median *mRS* of patients who survived was 3 (IQR 2–4). Patients who showed favorable outcomes seem to be younger, less affected at admission, had hypercholesterolemia and had suffered the bleeding in the lobar region (Table 2).

In the multivariate logistic regression analysis, a good functional outcome at discharge from the acute hospital was decreased by an age  $> 70$  years (OR, 0.56; 95%-CI, 0.35–0.9;  $p = .017$ ), NIHSS score  $> 10$  at admission (OR, 0.07; 95%-CI, 0.04–0.13;  $p < .001$ ), and development of pneumonia during hospitalization (OR, 0.35; 95%-CI, 0.2–0.6;  $p < .001$ ).

## 4 | DISCUSSION

Intracerebral hemorrhage is a cerebrovascular disease with poor outcomes. It is associated with a considerably high rate of disability and mortality, especially in the short time following the event, regardless of conservative management or interventional care (Kim et al., 2015; Mendelow et al., 2013; Qureshi et al., 2010; Qureshi & Qureshi, 2018; Trabert & Steiner, 2012).

Comparable to previous study results (Hemphill et al., 2001; Mayer & Rincon, 2005), the intracerebral hemorrhage seems to be

Baseline Characteristics	All (N = 613)	Surviving (n = 465)	Died (n = 148)	p-value
Age (years), mean (SD)	72 (12.7)	70 (13)	77 (11.3)	<.001
Female sex	313 (51.1%)	221 (47.5)	91 (62)	.003
NIHSS, median (IQR)	10 (4–18)	7 (3–13)	20 (16–30)	<.001
Medical history				
Hypertension	474 (77.3)	374 (84)	99 (71.2)	.002
History of diabetes	114 (18.6)	86 (19)	27 (19.7)	.8
Hypercholesterolemia	103 (16.8)	86 (19)	17 (12.4)	.06
Atrial fibrillation	127 (20.7)	92 (20.5)	35 (25.7)	.2
Alcohol use	43 (7.0)	37 (8)	6 (4.2)	.1
History of epilepsy	29 (4.7)	22 (4.9)	7 (5.0)	1.0
Prior medication				
ASA	161 (26.3)	128 (29)	32 (30.5)	.7
Clopidogrel	28 (4.6)	21 (4.8)	6 (6.1)	.4
Oral anticoagulants	86 (14.0)	60 (13.8)	26 (25.2)	.004
Statins	115 (18.8)	96 (22)	19 (18.8)	.5
Hospital stay (days), median (IQR)	11 (6–16)	12 (10–17)	3 (1–6)	<.001
Bleeding localization				
Lobar region	282 (46.0)	192 (41)	86 (58.5) <sup>a</sup>	<.001
Basal ganglia	193 (31.5)	159 (34)	34 (23.1) <sup>a</sup>	
Pons	18 (2.9)	10 (2)	8 (5.4) <sup>a</sup>	
Thalamus	53 (8.6)	48 (10)	5 (3.4) <sup>a</sup>	
Cerebellum	38 (6.2)	32 (7)	6 (4.1)	
Multifocal	18 (2.9)	12 (3)	6 (4.1)	
Ventricle	9 (1.5)	6 (1.3)	3 (2.0)	
Ventricular extension	197 (32)	115 (25)	82 (56)	<.001
Bleeding etiology				
Uncertain	110 (18.0)	69 (15)	42 (28.6) <sup>a</sup>	<.001
Hypertensive	288 (47.1)	237 (51)	51 (34.7) <sup>a</sup>	
Oral anticoagulation	66 (10.8)	46 (10)	20 (13.6)	
Amyloid angiopathy	67 (10.9)	58 (13)	9 (6.1) <sup>a</sup>	
Other causes <sup>b</sup>	81 (13.2)	56 (12)	25 (17.0)	

Note: Values are in absolute number and percentage unless other indicated.

Abbreviations: ASA, acetylsalicylic acid; SD, standard deviation.

<sup>a</sup>Change's by Bonferroni-corrections are significant.

<sup>b</sup>Indicates rare known causes (e.g., small vessel malformation, brain tumors, sinus venous thrombosis etc).

**TABLE 1** Baseline characteristics and comparison between survival and death

very lethal in the early short period after onset. We noticed in the group that did not survive the hemorrhage that four out of five patients died within the first week. However, it is important to note that patients who died in the hospital were older and much more severely neurologically impaired by the time of admission to the hospital than those who survived. In Addition, the mortality was associated with gender, consciousness level, history of hypertension, and anticoagulation as well as the localization of bleeding. Similar findings have been shown in survival with good outcomes.

Conservative treatment focuses on regulating and controlling blood pressure and the treatment of hypertension. Reduction in systolic blood pressure to values between 130 and 150 mm Hg, particularly in the early phase of symptom onset could play a role in reducing the expansion of intracerebral bleeding (Qureshi et al., 2010; Qureshi & Qureshi, 2018).

Contrarywise, we found that the previous hypertension was associated with survive of bleeding in the logistic regression analysis. This could be related to the effectiveness of the preexisting

**TABLE 2** Baseline characteristics, risk factors, bleeding localization, and complications of patients with spontaneous intracerebral hemorrhage

Baseline characteristics	Functional outcome at discharge (N = 445)		p-value
	Good outcome; mRS = ≤2 (n = 193)	Poor outcome; mRS = 3–5 (n = 252)	
Age (years), mean (SD)	68 (12)	72 (13)	.002
Female gender	87 (45.3%)	125 (50)	.3
NIHSS, median (IQR)	3 (2–6)	12 (7–16)	<.001
<b>Medical history</b>			
Arterial hypertension	156 (82)	213 (85)	.2
Alcohol abuse	11 (5.7)	25 (10)	.1
Diabetes mellitus	35 (18.3)	51 (20)	.5
Hypercholesterolemia	47 (24.6)	39 (15.6)	.019
Atrial fibrillation	34 (17.7)	57 (23)	.2
<b>Prior medication</b>			
ASS	58 (30.5)	68 (28)	.6
Clopidogrel	8 (4)	13 (5)	.5
Anticoagulants	26 (14)	34 (14)	.9
Statin	47 (25)	49 (20)	.2
Hospitalization length (days), median (IQR)	11 (8–13)	15 (11–20)	<.001
<b>Bleeding localization</b>			
Lobar	91 (47.4)	93 (37) <sup>a</sup>	.058
Basal ganglia	56 (29.2)	94 (37.5)	
Pons	5 (2.6)	5 (2.6)	
Thalamus	15 (7.8)	30 (12)	
Cerebellum	15 (7.8)	16 (6.4)	
Multifocal	2 (1.0)	10 (4)	
Ventricular	5 (2.6)	1 (0.4)	
Hypophysis	1 (0.5)	0	
Intraventricular extension	31 (16)	80 (32)	<.001
<b>Bleeding etiology</b>			
Uncertain	32 (16.7)	32 (12.7)	.8
Hypertensive	89 (46.4)	139 (55)	
Oral anticoagulation	21 (10.9)	25 (10)	
Amyloid angiopathy	25 (13.0)	29 (11.6)	
Other causes <sup>b</sup>	25 (13.0)	23 (9)	

Note: Values in numbers and percentage unless otherwise indicated.

Abbreviations: ASS, acetylsalicylic acid; IQR, interquartile range; SD, standard deviation.

<sup>a</sup>Indicates that changes in the Bonferroni methods are significant.

<sup>b</sup>Indicates rare known causes (e.g., small vessel malformation, brain tumors, sinus venous thrombosis etc).

hypertensive drugs that predict hypertensive blood pressure above 180mm Hg and bleeding expansion.

Recent study (Bernardo, Rebordao, Machado, Salgado, & Pinto, 2019) showed that in an age between 18 and 65 years, the hospital mortality was found remarkably lower (14.9%) than that in our study with mean age of 70 and 77 years among patients who died. However, the age is uncontrollable predictor for the prognosis compared to other factors, for example, pneumonia that leads to worsening of prognosis (Lindner et al., 2019).

Studies have shown that up to 20% of hemorrhage survivors return to independency (mRS ≤ 2) at 6 months, and these rates fairly have changed over the past two decades (Flaherty et al., 2006; Grysiewicz, Thomas, & Pandey, 2008). Whereas the prognosis is related to the size and localization of hemorrhage, for example, hemorrhage lacunar stroke that accounts for 7.4% of cerebral hemorrhages is associated with a more favorable prognosis as compared to the remaining group of intracerebral hemorrhages, usually with survival and deficit-free at discharge from the hospital in 22.8% of

cases (Arboix et al., 2002; Arboix, Carica-Eroles, Massons, Oliveres, & Targa, 2000).

In our study, the functional good outcome was significantly decreased in patients who were older (>70 years) and severely affected at admission (NIHSS score > 10) as well as in cases of development of pneumonia during hospitalization.

Interestingly, the hypercholesterolemia was found to be positively associated with the prognosis, in particular favorable functional outcomes, whereas the preexisting of statin use showed no association with the mortality and disability in patients with SICH. Similar findings were found in a large cohort study that showed low cholesterol level was associated with poor functional and death at 3 months after event (Chen et al., 2017).

Our study has a limitation that it does not include long-term follow-up. Despite all efforts, the prognosis of SICH is still devastating and is associated with poor outcomes compared to an ischemic stroke.

In summary, SICH is a lethal disease apart from the management and maximal care. The survival is related to factors that are in whole or in part uncontrollable.

## DISCLOSURE

The data were presented in the European Stroke Organization Conference 2018. The abstract was published in the European Stroke Journal 2018.

## CONFLICT OF INTEREST

On behalf of all authors, the corresponding author declares that no potential conflict of interest with respect to the research, authorship, and/or publication of this article is present.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## ORCID

Mohamed Al-Khaled  <https://orcid.org/0000-0001-8406-5044>

Toralf Brüning  <https://orcid.org/0000-0002-7897-2905>

## REFERENCES

- Al-Khaled, M., Eggers, J., & Qug, S. S. S. (2014). Prognosis of intracerebral hemorrhage after conservative treatment. *Journal of Stroke and Cerebrovascular Diseases: The Official Journal of National Stroke Association*, 23, 230–234. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2012.12.018>
- Arboix, A., Carica-Eroles, L., Massons, J., Oliveres, M., & Targa, C. (2000). Hemorrhagic lacunar stroke. *Cerebrovascular Disease*, 10, 229–234. <https://doi.org/10.1159/000016061>
- Arboix, A., Comes, A., Carica-Eroles, L., Massons, J., Oliveres, M., Balcells, M., & Targa, C. (2002). Site of bleeding and early outcome in primary intracerebral hemorrhage. *Acta Neurologica Scandinavica*, 105, 282–288.
- Bernardo, F., Rebordao, L., Machado, S., Salgado, V., & Pinto, A. N. (2019). In-hospital and long-term prognosis after spontaneous intracerebral hemorrhage among young adults aged 18–65 years. *Journal of Stroke and Cerebrovascular Diseases: The Official Journal of National Stroke Association*, 3, 104350. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2019.104350>
- Broderick, J., Connolly, S., Feldmann, E., Hanley, D., Kase, C., Krieger, D., et al (2007). Guidelines for the management of spontaneous intracerebral hemorrhage in adults: 2007 update: A guideline from the American Heart Association/American Stroke Association Stroke Council, High Blood Pressure Research Council, and the Quality of Care and Outcomes in Research Interdisciplinary Working Group. *Circulation*, 116, e391–413.
- Butcher, K., & Laidlaw, J. (2003). Current intracerebral haemorrhage management. *Journal of Clinical Neuroscience: Official Journal of the Neurosurgical Society of Australasia*, 10, 158–167. [https://doi.org/10.1016/S0967-5868\(02\)00324-7](https://doi.org/10.1016/S0967-5868(02)00324-7)
- Chen, Y.-W., Li, C.-H., Yang, C.-D., Liu, C.-H., Chen, C.-H., Sheu, J.-J., ... Hsu, C. Y. (2017). Low cholesterol level associated with severity and outcome of spontaneous intracerebral hemorrhage: Results from Taiwan Stroke Registry. *PLoS ONE*, 12(4), e0171379. <https://doi.org/10.1371>
- European Stroke Initiative Writing C, Writing Committee for the EEC, Steiner, T., Kaste, M., Forsting, M., & Mendelow, D. ... Hacke, W. (2006). Recommendations for the management of intracranial haemorrhage - part i: Spontaneous intracerebral haemorrhage. The European stroke initiative writing committee and the writing committee for the eusi executive committee. *Cerebrovascular Diseases*, 22, 294–316.
- Feigin, V. L., Lawes, C. M., Bennett, D. A., & Anderson, C. S. (2003). Stroke epidemiology: A review of population-based studies of incidence, prevalence, and case-fatality in the late 20th century. *Lancet Neurology*, 2, 43–53. [https://doi.org/10.1016/S1474-4422\(03\)00266-7](https://doi.org/10.1016/S1474-4422(03)00266-7)
- Ferro, J. M. (2006). Update on intracerebral haemorrhage. *Journal of Neurology*, 253, 985–999. <https://doi.org/10.1007/s00415-006-0201-4>
- Flaherty, M. L., Haverbusch, M., Sekar, P., Kissela, B., Kleindorfer, D., Moomaw, C. J., ... Woo, D. (2006). Long-term mortality after intracerebral hemorrhage. *Neurology*, 66, 1182–1186. <https://doi.org/10.1212/01.wnl.0000208400.08722.7c>
- Gregson, B. A., Rowan, E. N., Francis, R., McNamee, P., Boyers, D., Mitchell, P., ... Mendelow, A. D. (2015). Surgical trial in traumatic intracerebral haemorrhage (stitch): A randomised controlled trial of early surgery compared with initial conservative treatment. *Health Technology Assessment*, 19, 1–138. <https://doi.org/10.3310/hta19700>
- Grysiwicz, R. A., Thomas, K., & Pandey, D. K. (2008). Epidemiology of ischemic and hemorrhagic stroke: Incidence, prevalence, mortality, and risk factors. *Neurologic Clinics*, 26(871–895), vii. <https://doi.org/10.1016/j.ncl.2008.07.003>
- Hemphill, J. C. 3rd, Bonovich, D. C., Besmertis, L., Manley, G. T., & Johnston, S. C. (2001). The ich score: A simple, reliable grading scale for intracerebral hemorrhage. *Stroke; A Journal of Cerebral Circulation*, 32, 891–897. <https://doi.org/10.1161/01.STR.32.4.891>
- Hemphill, J. C. 3rd, Farrant, M., & Neill, T. A. Jr (2009). Prospective validation of the ich score for 12-month functional outcome. *Neurology*, 73, 1088–1094. <https://doi.org/10.1212/WNL.0b013e3181b8b332>
- Kim, H. T., Lee, J. M., Koh, E. J., & Choi, H. Y. (2015). Surgery versus conservative treatment for spontaneous supratentorial intracerebral hemorrhage in spot sign positive patients. *Journal of Korean Neurosurgical Society*, 58, 309–315. <https://doi.org/10.3340/jkns.2015.58.4.309>
- Kolominsky-Rabas, P. L., Sarti, C., Heuschmann, P. U., Graf, C., Siemonsen, S., Neundoerfer, B., ... von Stockert, T. R. (1998). A prospective community-based study of stroke in Germany—the Erlangen stroke project (espro): Incidence and case fatality at 1, 3, and 12 months. *Stroke; A Journal of Cerebral Circulation*, 29, 2501–2506. <https://doi.org/10.1161/01.STR.29.12.2501>

- Lindner, A., Kofler, M., Rass, V., Ianosi, B., Gaasch, M., Schiefecker, A. J., ... Helbok, R. (2019). Early predictors for infectious complications in patients with spontaneous intracerebral hemorrhage and their impact on outcomes. *Front Neurol*, 10, 817. <https://doi.org/10.3389/fneur.2019.00817>
- Mayer, S. A., & Rincon, F. (2005). Treatment of intracerebral haemorrhage. *The Lancet Neurology*, 4, 662–672. [https://doi.org/10.1016/S1474-4422\(05\)70195-2](https://doi.org/10.1016/S1474-4422(05)70195-2)
- Mendelow, A. D., Gregson, B. A., Rowan, E. N., Murray, G. D., Gholkar, A., Mitchell, P. M. et al (2013). Early surgery versus initial conservative treatment in patients with spontaneous supratentorial lobar intracerebral haematomas (stich ii): A randomised trial. *Lancet*, 382, 397–408. [https://doi.org/10.1016/S0140-6736\(13\)60986-1](https://doi.org/10.1016/S0140-6736(13)60986-1)
- Qureshi, A. I., Palesch, Y. Y., Martin, R., Novitzke, J., Cruz-Flores, S., Ehtisham, A. ... Antihypertensive Treatment of Acute Cerebral Hemorrhage Study Investigators (2010). Effect of systolic blood pressure reduction on hematoma expansion, perihematomal edema, and 3-month outcome among patients with intracerebral hemorrhage: Results from the antihypertensive treatment of acute cerebral hemorrhage study. *Archives of Neurology*, 67, 570–576. <https://doi.org/10.1001/archneurol.2010.61>
- Qureshi, A. I., & Qureshi, M. H. (2018). Acute hypertensive response in patients with intracerebral hemorrhage pathophysiology and treatment. *Journal of Cerebral Blood Flow and Metabolism*, 38, 1551–1563.
- Sacco, R. L., Boden-Albala, B., Gan, R., Chen, X., Kargman, D. E., Shea, S., ... Hauser, W. A. (1998). Stroke incidence among white, black, and hispanic residents of an urban community: The northern manhattan stroke study. *American Journal of Epidemiology*, 147, 259–268. <https://doi.org/10.1093/oxfordjournals.aje.a009445>
- Sadamasa, N., Yoshida, K., Narumi, O., Chin, M., & Yamagata, S.. (2011). [in-hospital mortality in patient with acute ischemic and hemorrhagic stroke]. *Brain Nerve*, 63, 995–999.
- Thieme Verlag (2012). *Leitlinien für diagnostik und therapie in der neurologie*. Stuttgart: Thieme Verlag.
- Trabert, J., & Steiner, T. (2012). Medical versus surgical management of intracerebral hematomas. *Current Atherosclerosis Reports*, 14, 366–372. <https://doi.org/10.1007/s11883-012-0259-7>
- van Asch, C. J., Luitse, M. J., Rinkel, G. J., van der Tweel, I., Algra, A., & Klijn, C. J. (2010). Incidence, case fatality, and functional outcome of intracerebral haemorrhage over time, according to age, sex, and ethnic origin: A systematic review and meta-analysis. *Lancet Neurology*, 9, 167–176. [https://doi.org/10.1016/S1474-4422\(09\)70340-0](https://doi.org/10.1016/S1474-4422(09)70340-0)
- Weimar, C., Weber, C., Wagner, M., Busse, O., Haberl, R. L., Lauterbach, K. W., & Diener, H. C. (2003). Management patterns and health care use after intracerebral hemorrhage. A cost-of-illness study from a societal perspective in Germany. *Cerebrovascular Diseases*, 15, 29–36. <https://doi.org/10.1159/000067119>

**How to cite this article:** Al-Khaled M, Awwad S, Brüning T. Nontraumatic spontaneous intracerebral hemorrhage: Baseline characteristics and early outcomes. *Brain Behav*. 2020;10:e01512. <https://doi.org/10.1002/brb3.1512>