

REVIEW

Open Access

HIV-associated neurocognitive disorders

Montserrat Sanmarti^{1,2*}, Laura Ibáñez², Sonia Huertas³, Dolors Badenes³, David Dalmau^{1,2}, Mark Slevin⁴, Jerzy Krupinski^{2,3,4}, Aurel Popa-Wagner⁵ and Angeles Jaen²

Abstract

Currently, neuropsychological impairment among HIV+ patients on antiretroviral therapy leads to a reduction in the quality of life and it is an important challenge due to the high prevalence of HIV-associated neurocognitive disorders and its concomitant consequences in relation to morbidity and mortality- including those HIV+ patients with adequate immunological and virological status. The fact that the virus is established in CNS in the early stages and its persistence within the CNS can help us to understand HIV-related brain injury even when highly active antiretroviral therapy is effective. The rising interest in HIV associated neurocognitive disorders has led to development new diagnostic tools, improvement of the neuropsychological tests, and the use of new biomarkers and new neuroimaging techniques that can help the diagnosis. Standardization and homogenization of neurocognitive tests as well as normalizing and simplification of easily accessible tools that can identify patients with increased risk of cognitive impairment represent an urgent requirement. Future efforts should also focus on diagnostic keys and searching for useful strategies in order to decrease HIV neurotoxicity within the CNS.

Keywords: HIV, Neurocognitive disorders, HAND

Review

Disruption of neurocognitive functioning is one of the most frequent complications in patients infected with HIV nowadays. This is a common reason for consultation of HIV patients and it negatively affects their quality of life, treatment adherence and life span. Since the introduction of Highly Active Antiretroviral Therapy (HAART) the spectrum of HIV-associated neurocognitive disorders (HAND) has been radically changed with a significant reduction in dementia but with a high prevalence of asymptomatic and mild neurocognitive impairments. Currently, in clinical practice, patients with HIV are still admitted to the clinic for memory, concentration or planification problems even when virology is under control. These usually begin with subtle changes but it can lead to more severe forms of neurocognitive impairment. The aim of this review is to describe the different types of neurocognitive disorders, possible mechanisms of development, epidemiology and risk factors in HIV patients, as well as the clinical approach and current treatment of HAND.

Classification

Before 1991, there was only one kind of neurocognitive disorder defined, the HIV-associated dementia (HAD), which was known as the complex AIDS-Dementia. It affected patients with severe immune-depression causing severe impairment of cognition, frequently accompanied by motor and behavioural alterations. More recently, the American Academy of Neurology proposed a new classification by defining two levels of neurological impairment in patients with HIV: the classical HAD and the minor cognitive motor disorder (MCMD) representing patients that did not meet dementia criteria but complained of slight impairments that interfered with their daily life [1].

In 2007, a further revised classification system of HAND was introduced which is thought to be more precise and sensitive (Frascati Criteria). It describes, beside HAD, other two neurocognitive disorders: Mild Neurocognitive disorder (MND) and Asymptomatic Neurocognitive impairment (ANI). MND is defined as mild to moderate impairment within at least two cognitive areas with at least mild impairment of daily function. ANI is defined as any degree of neuropsychological testing impairment in at least two cognitive domains but without causing an observable functional impairment [2] (See the "Classification of HAND" Section). Finally, cognitive neuropsychology

* Correspondence: msanmarti@mutuaterrassa.cat

¹Servei de Medicina Interna, Unitat VIH/Sida, Hospital Universitari MútuaTerrassa, Pl.Dr.Robert, 5 088221, Terrassa, Barcelona, Spain

²Fundació Docència i Recerca MútuaTerrassa, Terrassa, Barcelona, Spain
Full list of author information is available at the end of the article

aims to elucidate the component processes of HAND across the domains of executive functions, motor skills, speeded information processing, episodic memory, attention/working memory, language, and visuoception [3].

Classification of HIV-associated neurocognitive disorders

Asymptomatic neurocognitive impairment (ANI)

1. No evidence of preexisting cause. Cognitive impairment must be attributable to HIV and no other etiology (e.g. dementia, delirium).
2. The cognitive impairment does not interfere with activities of daily living.
3. Involves at least two cognitive areas (memory, attention, language, processing speed, sensory-perceptual, motor skills) documented by performance of > 1 standard deviation below the mean of standardized neuropsychological testing.

Mild neurocognitive disorder (MND)

1. No evidence of preexisting cause. Cognitive impairment must be attributable to HIV and no other etiology (e.g. dementia, delirium).
2. At least mild interference in > 1 activities of daily living including mental acuity, inefficiency at work, homemaking or social functioning.

HIV-Associated dementia (HAD)

1. No evidence of another preexisting cause for dementia (i.e. CNS infections, CNS neoplasm, cerebrovascular disease).
2. Marked interference in activities of daily living.
3. Marked cognitive impairment involving at least two cognitive domains by performance of > two standard deviation below the mean of standardized neuropsychological tests, especially in learning of new information, slowed information processing and defective attention or concentration.

Neuropathogenesis

The reason for neurocognitive disorders in HIV-patients is still unclear. It is well known that the central nervous system (CNS) is one of the target organs where HIV can be detected soon after primary infection. Early neuroinvasion is characterized by measurable markers of CSF inflammation (eg, neopterin level) and by brain parenchymal inflammation detected by magnetic resonance spectroscopy (MRS), although changes in neurocognitive functioning are seen more clearly in advanced stages [2,4]. HIV enters the brain carried within migrating monocytes and lymphocytes that cross the blood brain barrier (BBB) and possibly CD4+ T lymphocytes that use the brain as a reservoir for viral replication. After crossing the BBB, HIV infected monocytes become active perivascular macrophages being able to produce HIV within the CNS, release free virions and facilitate infection of microglial cells [5].

Both cells types allow HIV replication concomitant with expression of neurotoxic molecules- thought to be the ones involved in the pathogenesis of HAND [6] (eg. soluble immune mediators). Astrocytes may also harbour HIV and contribute to HIV-related brain disease through mechanisms of astrogliosis induced by local chemokines and cytokines leading to increased BBB permeability and consequently, monocyte and lymphocyte migration. Neurons are not productively infected.

Autopsy studies of AIDS patients with HAD shows characteristic white matter changes and demyelination, microglial nodules, multinucleated giant cells and perivascular infiltrates [7]. A variety of proteins encoded by the viral genome were identified as neurotoxic agents, including gp120, the virus's envelope protein, and transactivator of transcription (Tat). Gp120 is necessary for infectivity, but also interacts with host cellular receptors to alter glutamate pathway signaling and induce cytokine production that can injure neurons and affect the activation state of microglia and astrocytes.

Tat protein which is produced by infected astrocytes [8] has both neurotoxic and proinflammatory features. In several studies where Tat-expressing astrocytes were injected into the rat dentate gyrus it caused dendritic loss and neuronal cell death [9]. Furthermore, it is known that the virus is not evenly distributed in the CNS. Different areas of the brain are more susceptible to HIV due to the presence of individual viral genetic conditions and characteristics of the host [10]. The regions in which HIV is most prevalent are basal ganglia and the hippocampus, and to a lesser extent in the cortex and cerebellar grey matter of the mid-frontal cortex. These findings correspond with neuropsychological impairment of the fronto-subcortical-region and dysfunction, corresponding to frontal and subcortical areas [11]. Patients infected with HIV frequently have co-morbid conditions that may contribute to or amplify the pathogenicity of HIV, thus having an important role in the development of neurocognitive impairment (e.g. drug abuse and alcohol or infection with viral co-pathogens such as hepatitis C virus) [12].

Other factors that may contribute to HAND are the prolonged survival of HIV-infected patients, thereby extending the brain's exposure to HIV virions and proteins, the use of increasingly toxic combinations of poorly penetrating drugs in highly antiretroviral-experienced AIDS patients, and selection of more virulent HIV strains with higher replication rates and greater virulence in neural tissues. Eden and colleagues found that a substantial number of patients still show signs of macrophage/microglia activation even after 4 years of viral suppression with HAART [13]. All these factors converge in producing damage to dendrites and synapses leading to the disruption of the highly integrated functioning of neural systems that is required to process information, leading to HIV-associated neurocognitive disorders.

Epidemiology

Several studies demonstrated that neurocognitive deficits are more common in HIV+ patients than the HIV- population, and these deficits were independent of either ART, or disease state [14,15]. However, one recent study demonstrated similar rates of cognitive impairment between a subgroup of HIV+ patients, with a high CD4 level and HIV viral suppression as compared with HIV- controls [16]. A high prevalence of HIV-associated neurocognitive disorders (HAND) is associated with a change in the pattern of HAND and with an increased prevalence of mild disease vs dementia [17,18]. Before the introduction of HAART, the HIV-associated neurocognitive impairment was categorized as AIDS-dementia complex, HIV encephalitis or encephalopathy, with an estimated prevalence of 16% in patients with a diagnosis of AIDS [19]. However, in the post-HAART era, the prevalence of dementia is estimated to be less than 5% [17,20]. In contrast, it appears that mild cognitive disorders are very common despite treatment [17].

The prevalence of HAND is estimated in approximately 40-50% of all cases. Mild neurocognitive disorder (MND) and Asymptomatic Neurocognitive Impairment (ANI) are now more common than HAD. In a cross-sectional study conducted in Switzerland, there was a prevalence of HAND in 85% of patients with evidence of cognitive decline, and 64% among those who did not demonstrate any deterioration. In the first group, 24% had ANI, 52% MND and 8% HAD, while in the second group 60% had ANI, 4% MND and 0% HAD. Subclinical alterations that do not affect the functionality of the patients or alter their daily lives are the most frequent in both groups [17]. In a cross-sectional study conducted on 1555 HIV infected patients the CNS HIV Anti-Retroviral Therapy Effects Research (CHARTER) found that 52% had HAND, of which 33% had ANI, 12% MND and 2% HAD. The prevalence of impairments correlated with the number of patient comorbidities [20]. Another study showed a 48% rate of occurrence of cognitive decline in HIV-infected patients [21], whilst the Aquitaine cohort found a prevalence of 58.5% TNAVs, 20.8% ANI, 31% MND, and 6.7% HAD [18]. In Spain, a very recent multicenter study conducted by questionnaire, found that 50% of the HIV+ population complained of cognitive impairments. A high proportion of them (72.1%) believed that their cognitive impairments interfered with daily life activity [22].

Progression and implication

The real clinical relevance of ANI is not clear. It does not interfere with daily life and in addition, it is not proven that ANI patients are at increased risk of progression to more advanced stages of deterioration. Moreover, there are no satisfactory biomarkers or clinical prognostic/diagnostic indicators associated with this subgroup of patients

that can determine evolution and end-point. The CHARTER study reported preliminary results of monitoring at 18 and 42 months. The proportion of patients with worsening neurocognitive function according to their basal HAND classification was different in each group (ANI-23%, MND-30%, not TNAV-13%), however there was no statistically significant difference in mean decline in neurocognitive function between ANI and MND groups [23]. From a prospective cohort, ACTG Longitudinal Linked Randomized Trials (ALLRT) of 1160 patients from clinical trials, 921 were examined using neuropsychological tests, and were subsequently followed for 48 weeks. The prevalence of HAND was 39% at inclusion, of them 44% improved after one year. In contrast, of those patients without HAND, 21% developed worse cognitive function [24].

Risk factors

The risk factors associated with neurocognitive disorders are not well established in the post HAART era. They may include host factors (e.g. genetic predisposition, metabolic disorders, cardiovascular risk factor or aging), HIV-related factors (AIDS, immune activation, drug resistance) and comorbidities such as hepatitis C virus co-infection or depression. Poor immunological status, reflected by lower nadir CD4 cell count, has been associated with neurocognitive impairment before and after HAART [20,24,25]. However, other studies did not find this association with the CD4 levels [17,18,26]. The correlation with virological status is even less clear in the HAART era [17,18,26]. Older age was associated with neurocognitive impairment [18,21,25] and HAD [27]. Valcour and collaborators [27] found about 3-fold increased risk of HAD in patient >50 years compared with those between 20 to 39 years old, independently of other factors. Harezlak and co-workers found an association between age and markers of brain injury in HIV infected patients as compared with those being HIV negative [21]. However, in other studies there was not a higher age-related impairment of neuropsychological function between HIV+ and HIV- populations [28]. Comorbidities, such as coinfection with hepatitis C virus (HCV), were found to be a risk factor for the presence of HAND [25,29] that could be explained by an independent neurotoxic effect of HCV. Other comorbidities as obesity and diabetes were associated with neurocognitive impairment, especially in older patients [30].

Data on neuropsychiatric symptoms in coinfecting persons are inconclusive at this time. With regard of depression was the most common disorder by far, reaching rates of 71% and 62% for past depression in coinfecting and HIV-infected persons, respectively, and 42% for current rates in both groups. Past dysthymia, past posttraumatic stress disorder, and childhood conduct disorder were the next most prevalent, with rates for each ranging between 16% and 19% in both groups [31]. Other cardiovascular

risk factors and diseases were associated with lower cognitive performance [18,32]. Host genetic factors such as apolipoprotein E4 have been suggested as possible risk factors for neurocognitive impairment [33]. Whilst in the general population, sub-groups with the ApoE4 allele have a worse prognosis for neurodegeneration [34], this risk is less clear in the HIV+ population [35,36].

Clinical features

The landmark of neuropsychological impairment in patients infected with HIV has changed in recent years particularly after the introduction of HAART. Currently, most patients with alterations in neurocognitive tests have no evident symptoms or dysfunction in their daily lives (ANI). It has been demonstrated that the neurocognitive impairment in early stages of infection shows a subcortical pattern and in more advanced stages implicates more cortical areas. The main cognitive complaints of the patients with mild cognitive dysfunction involve subtle changes in working memory and attention, speed of informational processing, executive functioning (including organizing, planning and problem-solving) or difficulties in verbal fluency. Motor symptoms are less common but can include leg weakness, tremor or unsteady gait. HAD, however, is often characterized by a progressing subcortical dementia with classically severe memory loss and altered executive function. More advanced stages also show aphasia, agnosia and apraxia, that are more typical of cortical dementias such as Alzheimer's disease. Other neurobehavioral disturbances resulting from HIV-mediated neural damage include emotional and other behavioural disturbances (e.g. depression, anxiety, sleep disorders, mania and psychosis) [37].

Diagnosis

Any individual with HIV infection could be at risk of developing HAND [38]. Therefore, the need for a systematic study of cognitive HIV infected individuals seems increasingly evident. Although many different studies assessing neurocognitive impairment in HIV infection have been published, a standard approach to optimally assessing neuropsychological deficits does not currently exist. Multiple clinical tests are available for use in order to evaluate neurocognitive function although there is currently no single and universal neuropsychological battery to assess specifically people with HIV. Evaluation of the different cognitive functions requires a concerted effort (2–4 hours) with the need for expert staff specialized in clinical neuropsychology who has to evaluate at least five of the following cognitive abilities: verbal/language, attention/working memory, abstraction/executive functioning, learning/recall, speed of informational processing, and sensory-perceptual and motor skills. This has led to the need for screening instruments that, although maybe useful in the initial

screening, are insufficient to make a good neuropsychological diagnosis within clinics. The classic tests-minimal state examination (MMSE) or Montreal cognitive assessment (MoCA) are not sufficiently sensitive to screen for HAND unless the patients are symptomatic [39].


There are specific screening instruments for assessing the HIV Dementia Scale HAND (HDS) [40], *The International HIV Dementia Scale* (IHDS) [41] or the Montreal cognitive assessment (MoCA) [42]. All these tests can be performed in less than ten minutes. Other tests often used in conjunction with these include the *Brief Neurocognitive Screen* (BNC) [43], screening battery of HNRC [44], computed COGSTATE [45] or Neu Screening [46]. The diagnosis of HAND is confirmed by the clinical detection of cognitive impairment through neuropsychological testing and the evidence of neurocognitive impairment. Cognitive impairment must be attributable to HIV after excluding other causes that can justify the disorder (Figure 1).

Differential diagnosis

The differential diagnosis of neurocognitive disorders associated with HIV infections presents with two difficulties: 1) to discard all the pathologies that can lead to neurocognitive disorders in the general population, associated or not with HIV; 2) to determine the degree of contribution of certain comorbidities associated with HIV infection, before attributing neurological damage to HIV itself. This group may act as true comorbidity confounders while attempting to diagnose a neurocognitive disorder in patients associated with HIV infection [47]. When the presence of neurocognitive impairment is suspected, an exhaustive study has to be done in order to detect other potential causes of neurocognitive impairment, including laboratory testing of vitamin B12, folate level, TSH, syphilis, hepatitis C, glucose and vitamin B1 as well as screening for dyslipidaemia. Sometimes, it is necessary to examine CFS especially when there are other signs of CNS infection, fever, CD4 cell counts below 200 cells/mm³ or positive serology for syphilis.

Other testing should include screening for major depression and anxiety, prescription of psycho-pharmacologic treatments and identification of self-abuse of other substances. Frequently, patients present other comorbidities such as psychiatric illness, vascular cerebral disease, opportunistic infections, alcohol and drugs abuse. All these factors can cause neurocognitive disorders or contribute to HAND progression [2]. It is a requirement to know the real impact of comorbidities in order to find new prevention and treatment strategies.

There is a growing interest in finding biomarkers that are sensitive and specific for HAND. One of the most reliable biomarker currently used for monitoring disease progression and HAART effectiveness is CD4 cell count. Counts



above 200 cells/mL are rarely associated with dementia [48,49]. Detection of HIV RNA concentration in the CSF can also be determined. The presence of HIV RNA in CSF, even at high concentrations, is not diagnostic for HAND and cognitively asymptomatic subjects may have high Viral Load (VL). In the CHARTER study, patients with CSF VL < 2.5 copies/mL had a higher degree of neurocognitive impairment compared to the ones with a VL of 2.5-40 copies/mL [50]. Some studies have shown slight hyper-proteinorraquia and pleocytosis with increased mononuclear counts [51]. This increased cellularity has been positively correlated with HIV viral load in CSF and can be reversed with proper antiretroviral treatment. However, these changes can also be found in subjects without cognitive impairment, so are not specific for diagnosis of HAND. In HIV infected patients, different biomarkers have been tested for their effectiveness, including amyloid peptide B, Tau and phospho-Tau [52]. HIV population studies show conflicting results. Some authors confirm that CSF abnormalities in HIV patients with HAD are similar to those from Alzheimer's disease patients (increased tau and phospho-tau and decreased beta-amyloid peptide) [53] but still others do not support the idea [54]. A recent study,

however, by Steinbrink et al., found a positive correlation between tau levels and the degree of HAND [55].

The existence of reliable biomarkers would help to diagnose patients and would allow better understanding of the pathophysiology of cognitive impairment in HIV patients, as well as designing therapeutic strategies aimed at treating complications. Although there are no pathognomonic neuroimaging alterations in HAND, it is pertinent to recommend neuroimaging for investigating other neurological disorders (opportunistic infections, tumours and other structural lesions or ischemia/infarction). RM is recommended and it is preferred to TAC because of its sensitivity [47]. Other techniques are being investigated to improve the diagnosis of HAND such as cerebral volumetry, PET, MRI and spectroscopy. They all are able to identify structural, metabolic and functional disorders which are related to the presence of cognitive disorder.

Treatment

Since the main cause of cognitive impairment in HIV-infected patients is the virus infection itself, the treatment of choice (accepted treatment strategy) remains the use of HAART defined as a combination of three or more drugs. Clear improvement in neurological outcomes in the era of

HAART has also been achieved resulting in a notable reduction of HAND. However, HAART benefits for HIV cognitive disorders differ substantially between individuals [56] and several studies have shown in recent years the persistence of asymptomatic or mild/moderate forms of HAND [57] leading to speculations that systemically effective HAART might fail to control CNS-HIV infection, and also that the type of antiretroviral drug and its penetration may have a role in the development of HAND. CSF HIV RNA is generally lower than in plasma and normally decays in parallel to plasma viral load when HAART is initiated. Therefore, the majority of patients with undetectable plasma VL have also undetectable CSF VL. However, in some cases CSF decay is delayed in response to HAART and approximately 10% of patients treated with HAART (for at least 6 months) have VL detectable in LCR despite suppression of plasma human HIV RNA [58]. It has been demonstrated that a greater viral suppression within the CNS is associated with a better neurocognitive function [59] and may influence survival [60]. Eden et al., found that patients with neurological disease had a significantly higher CSF viral load both at baseline and at follow-up compared with neurologically asymptomatic patients [61].

There is a considerable heterogeneity among different ARV agents with respect to their CNS penetration. A number of drug characteristics influence CNS penetration across the BBB. The latter is composed of a complex of endothelial cells, transport proteins and basement membrane with fenestrated capillaries that together prevent passage of drugs and other substances into the brain and have the capacity to express molecular drug pumps or transports such as P-glycoprotein that can limit the amount of drug that passes into the brain. Letendre et al., created a CNS penetration effectiveness ranking system of different ARV agents based on chemical properties of the drugs (molecular weight, liposolubility, ion charge and its union with plasma proteins), data on CSF concentrations, and HAART effectiveness [62]. It was demonstrated that combination antiretroviral treatments with higher CNS penetration-effectiveness (CPE) achieved better CSF viral suppression [63] and could be associated with better cognition. There are no conclusive studies on the efficacy of HAART regimens regarding improvement of cognitive performance or which is the best combination to prevent or reduce cognitive impairment. In general, antiretroviral drugs have low molecular weight that favors crossing the BBB. The nucleoside analogue reverse transcriptase inhibitors (nRTIs) have a low molecular weight, followed by non nucleoside reverse transcriptase inhibitors (NNRTIs). The protease inhibitors (IPs) have clearly higher molecular weights and therefore show a low BBB penetration. In addition, their high affinity for P-glycoprotein makes it difficult to achieve high concentrations in the CNS. Ritonavir, a potent inhibitor of P-gp, improves CNS levels of other

PIs when used as a booster [64]. Drugs with less affinity for P-gp penetrate the CNS more easily. PIs and NNRTIs exhibit greater fat solubility than nRTIs. Recently, new ARVs drugs have been approved and are also used in combination with other antiretroviral agents for treating HIV adult patients: Raltegravir, an integrase inhibitor which prevents viral DNA insertion into the human cell genome and Maraviroc, an antagonist of the CCR5 chemokine receptor, which also serves as an HIV-1 co-receptor. They have a lower plasma protein binding level than other ARVs and seem to be a substrate of P-gp transporter. Enfuvirtide is a fusion inhibitor which is indicated for HIV patients who have failed prior ART regimens. There are limited data on the CNS pharmacokinetics of this inhibitor. Finally, treatment regimens should be designed to optimize CNS penetration without sacrificing viral suppression or immunological improvement. Therefore a treatment regimen with good CNS penetration could be critical in the case of patients with disorders associated with neurocognitive function [65].

Other pharmacological treatments

Despite the introduction of HAART, the persistence of HAND has led to development of new strategy adjuvants to HAART. Multiple drugs with neuroprotective properties have been evaluated. Minocycline, a tetracyclin with neuroprotective properties and good penetration through the BBB is being evaluated as a possible adjunct to HAART. However, clinical trials performed on a cohort of 107 individuals have not demonstrated improvement in cognitive function [66]. Memantine, a non-competitive antagonist of N-methyl-D-aspartate, approved for treatment of Alzheimer's disease has also been tested in a randomized double-blind, placebo-controlled, multicenter trial within the Adult AIDS Clinical Trials Group. No significant improvement in patients with mild to severe cognitive impairment after 16 weeks of treatment was noted. However, the magnetic resonance spectroscopy data suggested that memantine may ameliorate neuronal metabolism, an important step to stabilizing or preventing neuronal injury [67]. Other drugs have been investigated, including lithium, selegiline and valproate. At the present time, there is no data supporting the use of any adjuvant therapy to HAART for the treatment of neurocognitive disorders associated with HIV infection although some drugs such as memantine have been shown to improve markers of neuronal damage and inflammation within the CNS.

Non pharmacologic interventions

There is a little information regarding pharmacological interventions in HIV populations which can be extrapolated to the general population. It is essential to modify lifestyle (diet, physical activity, stress) and to give up tobacco and alcohol. Currently, we can prevent cardiovascular risk factors

(e.g. HTA, DM, DLP). Adherence to treatment remains the central issue in HIV patients in order to keep control of VL. Neurocognitive impairment is strongly related to poor adherence [68]. The later can lead to drug resistance, increase morbidity/mortality and development of cognitive impairment, especially in the elderly [69]. Neuropsychological intervention programs proved to be useful in multiple pathologies like schizophrenia, acquired brain injury or Alzheimer's disease. However, few studies had been performed in the HIV field. Some of the programs showed some positive effects [70] although proper studies with new strategies presently do not exist.

Prognosis

Before the introduction of HAART, HIV associated neurocognitive impairment was considered a significant independent risk factor for AIDS related death. Although the median survival of patients with HAD has decreased considerably [71] it remains a negative prognostic factor associated with mortality [25,72]. Other studies have shown that mild cognitive impairment is also associated with poorer adherence to treatment [73], higher rates of virological failure [74], difficulties in the development of everyday function [75], loss of employment [76], impairments in the quality of life [77] and progression to dementia [78]. In the post-HAART era, the course of dementia progression has become more heterogeneous and appears to be related to factors such as antiretroviral therapy adherence, aging, substance abuse and psychosocial factors. Evolution is unknown in patients who have cognitive impairment. Few studies have been conducted to assess the incidence and persistence of HAND despite HAART. A study of cohorts which evaluated a group of 94 patients with HIV-related neurocognitive impairment, found that to 63% of them, had persistent neurocognitive deficits within 63 months of successful treatment. Also, 37% had permanent deficits without alterations in subsequent visits. They concluded that the severity of neurocognitive impairment at HAART initiation seems to be the strongest predictor of persistent neuropsychological deficits [26]. In another cohort of 1561 HIV+ patients the presence of HAND was associated with a 3-fold increase in mortality [79].

In recent years the concept of "brain reserve" (CR) has emerged to describe some individuals having an increased "baseline adaptive neuroplasticity", providing greater dynamic capacity for adjusting and remodeling cortical circuits to various stressors [80]. Another neuropsychiatric symptom that results from CNS viral damage is apathy is significantly associated with cognitive impairment in individuals with HIV; Previous literature has demonstrated CR's capacity to protect against cognitive decline and apathy in individuals in HIV-infected individuals across the age span, particularly after a certain threshold of disease severity [81].

Conclusions

Currently, neuropsychological impairment among HIV+ patients on ARV therapy leads to a reduction in the quality of life and it is an important challenge due to the high prevalence of HAND and its concomitant consequences in relation to morbidity and mortality- including those HIV+ patients with adequate immunological and virological status. The fact that the virus is established in CNS in the early stages and its persistence within the CNS can help us to understand HIV-related brain injury even when HAART is effective. Longitudinal studies focussing on neuropsychological function are needed, as there is a few data on progression and clinical significance of ANI in these patients. In addition, we have little information about risk factors or markers of progression of HAND. The most important risk factors associated with neurocognitive impairment have been described in cross-sectional studies or non-representative sample, and very often with contradictory results. It is essential to address comorbidities in patients with cardiovascular risk factors or depression. The rising interest in HIV associated neurocognitive disorders has led to development new diagnostic tools, improvement of the neuropsychological tests, and the use of new biomarkers and new neuroimaging techniques that can help the diagnosis. Standardization and homogenization of neurocognitive tests as well as normalizing and simplification of easily accessible tools to screen for patients with increased risk of cognitive impairment and prioritize full neurocognitive study is an urgent requirement. The short screening instruments would be more important in places where it is not possible to complete full study. For the moment, there is not a universal recommendation how to proceed in clinical practise. It is strongly recommended to ask patients routinely for cognitive symptoms as detection of neurocognitive impairment can justify starting ARV therapy. It is also important to make a correct monitoring of cognitive complaints on every visit.

Finally, standard treatment for HIV-associated neurocognitive disorders is an optimal antiretroviral treatment regimen, along with control and treatment of other comorbidities or psychiatric problems. For naïve HIV+ patients with evidence of cognitive deficits, it is recommended to start ART with an optimized central nervous system penetration effectiveness rank. However, pharmacodynamic studies of HAART are needed in order to determine which drugs can be more effective in preserving cognitive functioning and when it is necessary to change the treatment if cognitive impairment appears, even when virological control has been achieved. Future efforts should also focus on diagnostic keys and searching for useful strategies in order to decrease HIV neurotoxicity within the CNS.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed to the design and analysis of the review. MS, AJ, APW and JK drafted the manuscript. MS, AJ, APW, JK, MAS provide administrative support. All authors critically revised the manuscript. All authors read and approved the final manuscript.

Author details

¹Servei de Medicina Interna, Unitat VIH/Sida, Hospital Universitari MútuaTerrassa, Pl.Dr.Robert, 5 088221, Terrassa, Barcelona, Spain. ²Fundació Docència i Recerca MútuaTerrassa, Terrassa, Barcelona, Spain. ³Servei de Neurologia, Hospital Universitari MútuaTerrassa, Barcelona, Spain. ⁴Department of Healthcare Science, Manchester Metropolitan University, Manchester, UK. ⁵Clinic for Psychiatry and Psychotherapy, University of Medicine Rostock, Rostock, Germany.

Received: 3 January 2014 Accepted: 25 February 2014

Published: 4 March 2014

References

1. Report of a Working Group of the American Academy of Neurology AIDS Task force: **Nomenclature and research case definitions for neurologic manifestations of human immunodeficiency virus-type 1 infection.** *Neurology* 1991, **41**:778–785.
2. Antinori A, Arendt G, Becker JT, Brew BJ, Byrd DA, Cherner M, Clifford DB, Cinque P, Epstein LG, Goodkin K, Gisslen M, Grant I, Heaton RK, Joseph J, Marder K, Marra CM, McArthur JC, Nunn M, Price RW, Pulliam L, Robertson KR, Sacktor N, Valcour V, Wojna VE: **Updated research nosology for HIV-associated neurocognitive disorders.** *Neurology* 2007, **69**:1789–1799.
3. Woods SP, More DJ, Weber E, Grant I: **Cognitive neuropsychology of HIV-associated neurocognitive disorders.** *Neuropsychol Rev* 2009, **19**:152–168.
4. Valcour V, Chalermchai T, Sailasuta N, Marovich M, Lerdlum S, Suttichom D, Suwanwala NC, Jagodzinski L, Michael N, Spudich S, Griensven FV, de Souza M, Kim J, Ananworanich J: **Central nervous system viral invasion and inflammation during acute HIV infection.** *J Infect Dis* 2012, **206**:275–282.
5. Spudich S, Gonzalez-Scarano F: **HIV-1 related central nervous system disease: current issues in pathogenesis, diagnosis and treatment.** *Cold Spring Harb Perspect Med* 2012, **2**:a007120.
6. Gonzalez-Scarano F, Martin-Garcia J: **The neuropathogenesis of AIDs.** *Nat Rev Immunol* 2005, **5**:69.
7. Gray F, Chretien F, Vallat-Decouvaere AV: **The changing pattern of HIV neuropathology in the HAART era.** *J Neuropathol Exp Neurol* 2003, **62**:429.
8. Nath A, Haughey NJ, Jones M, Anderson C, Bell JE, Geiger JD: **Synergistic neurotoxicity by human immunodeficiency virus proteins Tat and gp120: protection by memantine.** *Ann Neurol* 2002, **47**:186–194.
9. Bruce-Keller AJ, Chauhan A, Dimayuga FO, Gee J, Keller JN, Nath A: **Synaptic transport of human immunodeficiency virus-Tat protein causes neurotoxicity and gliosis in rat brain.** *J Neurosci* 2003, **23**:8417–8422.
10. UNAIDS: **Cognitive and Neuropsychological impairment in Early HIV infection: Report of an expert consultation.** Geneva, Switzerland: UNAIDS; 1998.
11. Gatell JM, Clotet B, Pozsamer D, Miró JM, Mallolas J: **Guia practica del SIDA.** Editorial Antares; 2013:425–433. ISBN:9788488825100.
12. Patel SM, Thames AD, Arbid N, Panos SE, Castellon S, Hinkin CH: **The aggregate effects of multiple comorbid risk factors on cognition among HIV-infected individuals.** *J Clinical and Exp Neuropsychology* 2013, **35**:421–434.
13. Eden A, Price RW, Spudich S, Fuchs D, Hagberg L, Gisslen M: **Immune activation of the central nervous system is still present after > 4 years of effective highly active antiretroviral therapy.** *J Infect Dis* 2007, **19**:1778–1783.
14. Heaton RK, Franklin DR, Ellis RJ, McCutchan JA, Letendre SL, Leblanc S, Corkran SH, Duarte NA, Clifford DB, Woods SP, Collier AC, Marra CM, Morgello S, Mindt MR, Taylor MJ, Marcotte TD, Atkinson JH, Wolfson T, Gelman BB, McArthur JC, Simpson DM, Abramson I, Gamst A, Fennema-Notestine C, Jernigan TL, Wong J, Grant I, CHARTER Group, HNRC Group: **HIV-associated neurocognitive disorders before and during the era of combination antiretroviral therapy: differences in rates, nature, and predictors.** *J Neuroviral* 2011, **17**:3.
15. Mateen FJ, Shinohara RT, Carone M, Miller EN, McArthur JC, Jacobson LP, Sacktor N, Multicenter AIDS Cohort Study (MACS) Investigators: **Neurologic disorders incidence in HIV+ vs HIV- men: Multicenter AIDS cohort study, 1996–2011.** *Neurology* 2012, **79**:1873.
16. Crum-Cianflone NF, Moore DJ, Letendre S, Poehlman Roediger M, Eberly L, Weintrob A, Ganesan A, Johnson E, Del Rosario R, Agan BK, Hale BR: **Low prevalence of neurocognitive impairment in early diagnosed and managed HIV-infected persons.** *Neurology* 2013, **80**:371–379.
17. Simioni S, Cavassini M, Annoni J, Rimbault Abraham A, Bourquin I, Schiffer V, Calmy A, Chave JP, Giacobini E, Hirschel B, Du Pasquier RA: **Cognitive dysfunction in HIV patients despite long-standing suppression of viremia.** *AIDS* 2010, **24**:1243–1250.
18. Bonnet F, Amieva H, Marquant F, for the ANRS CO3 Aquitaine Cohort: **Cognitive disorders in HIV-infected patients: are they HIV-related?** *AIDS* 2013, **27**:391–400.
19. McArthur JC, Hoover DR, Bacellar H, Miller EN, Cohen BA, Becker JT, Graham NM, McArthur JH, Selnes OA, Jacobson LP: **Dementia in AIDS patients: incidence and risk factors. Multicenter AIDS cohort Study.** *Neurology* 1993, **43**:2245–2252.
20. Heaton RK, Clifford DB, Franklin DR, Woods SP, Ake C, Vaida F, Ellis RJ, Letendre SL, Marcotte TD, Atkinson JH, Rivera-Mindt M, Vigil OR, Taylor MJ, Collier AC, Marra CM, Gelman BB, McArthur JC, Morgello S, Simpson DM, McCutchan JA, Abramson I, Gamst A, Fennema-Notestine C, Jernigan TL, Wong J, Grant I, CHARTER Group: **HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy: CHARTER Study.** *Neurology* 2010, **75**:2087–2096.
21. Harezlak J, Buchthal S, Taylor M, Schifitto G, Zhong J, Daar E, Alger J, Singer E, Campbell T, Yiannoutsos C, Cohen R, Navia B, HIV Neuroimaging Consortium: **Persistence of HIV-associated cognitive impairment, inflammation, and neuronal injury in era of highly active antiretroviral treatment.** *AIDS* 2011, **25**:625–633.
22. Muñoz-Moreno JA, Fuster MJ, Fumaz CR, Ferrer MJ, Molero F, Jaen A, Clotet B, Dalmau D: **Quejas Cognitivas en Personas con VIH en España: Prevalencia y Variables Relacionadas.** *Med Clin (Barc)* 2013: [Epub ahead of print].
23. Heaton R, Franklin D, Woods S, Marra C, Clifford D, Gelman B, McArthur J, Morgello S, McCutchan A, Grant I, the CHARTER Group: **Asymptomatic mild HIV-associated Neurocognitive Disorder Increases Risk for Future Symptomatic Decline: A CHARTER Longitudinal Study. Nineteenth Conference on Retroviruses and Opportunistic Infections (CROI).** *Seattle* 2012:abstract 77.
24. Robertson KR, Smurzynski M, Parsons TD, Wu K, Bosch RJ, Wu J, McArthur JC, Collier AC, Evans SR, Ellis RJ: **The prevalence and incidence of neurocognitive impairment in the HAART era.** *AIDS* 2007, **21**:1915–1921.
25. Tozzi V, Balestra P, Serraino D, Bellagamba R, Corpolongo A, Piselli P, Lorenzini P, Visco-Comandini U, Vlassi C, Quartuccio ME, Giulianelli M, Noto P, Galgani S, Ippolito G, Antinori A, Narciso P: **Neurocognitive impairment and survival in a cohort of HIV-infected patients treated with HAART.** *AIDS Res Hum Retroviruses* 2005, **21**:706–713.
26. Tozzi V, Balestra P, Bellagamba R, Corpolongo A, Salvatori MF, Visco-Comandini U, Vlassi C, Giulianelli M, Galgani S, Antinori A, Narciso P: **Persistence of neuropsychologic deficits despite long-term highly active antiretroviral therapy in patients with HIV-related neurocognitive impairment: prevalence and risk factors.** *J Acquir Immune Defic Syndr* 2007, **45**:174–182.
27. Valcour V, Shikuma C, Shiramizu B, Watters M, Poff P, Selnes O, Holck P, Grove J, Sacktor N: **Higher frequency of dementia in older HIV-1 individuals: the Hawaii aging with HIV-1 cohort.** *Neurology* 2004, **63**:822–827.
28. Cysique LA, Maruff P, Bain MP, Wright E, Brew BJ: **HIV and age do not substantially interact in HIV-associated neurocognitive impairment.** *J Neuropsychiatry Clin Neurosci* 2011, **1**:83–89.
29. Sun B, Abadjian L, Rempel H, Monto A, Pulliam L: **Differential cognitive impairment in HCV coinfecting men with controlled HIV compared to HCV mono-infection.** *J Acquir Immune Defic Syndr* 2013, **62**:190–196.
30. McCutchan JA, Marquie-Beck JA, Fitzsimons CA, Letendre SL, Ellis RJ, Heaton RK, Wolfson T, Rosario D, Alexander TJ, Marra C, Ances BM, Grant I, CHARTER Group: **Role of obesity, metabolic variables, and diabetes in HIV-associated neurocognitive disorder.** *Neurology* 2012, **78**:485–492.
31. Hilsabeck RC, Castellon SA, Hinkin CH: **Neuropsychological aspects of coinfection with HIV and hepatitis C virus.** *Clin Infect Dis* 2005, **41**(Suppl 1):S38–S44.
32. Fabbiani M, Ciccarelli N, Tana M, Farina S, Baldonero E, Di Cristo V, Colafigli M, Tamburrini E, Cauda R, Silveri MC, Grima P, Di Giambenedetto S: **Cardiovascular risk factors and carotid intima-media thickness are**

- associated with lower cognitive performance in HIV-infected patients. *HIV Med* 2013, **14**:136–144.
33. Valcour V, Shikuma C, Shiramizu B, Watters M, Poff P, Selnes OA, Grove J, Liu Y, Abdul-Majid KB, Gartner S, Sacktor N: **Age, apolipoprotein E4, and the risk of HIV dementia: the Hawaii aging with HIV cohort.** *J Neuroimmunol* 2004, **157**:197–202.
 34. Reiman EM, Webster JA, Myers AJ, Hardy J, Dunckley T, Zismann VL, Joshipura KD, Pearson JV, Hu-Lince D, Huentelman MJ, Craig DW, Coon KD, Liang WS, Herbert RH, Beach T, Rohrer KC, Zhao AS, Leung D, Bryden L, Marlowe L, Kaleem M, Mastroeni D, Grover A, Heward CB, Ravid R, Rogers J, Hutton ML, Melquist S, Petersen RC, Alexander GE, Caselli RJ, Kukull W, Papassotiropoulos A, Stephan DA, Farina S, Baldonero E, Di Cristo V, Colafigli M, Tamburrini E, Cauda R, Silveri MC, Grima P, Di Giambenedetto S: **GAB2 alleles modify Alzheimer's risk in APOE epsilon4 carriers.** *Neuron* 2007, **54**:713–720.
 35. Leoni V, Solomon A, Kivipelto M: **Links between ApoE, brain cholesterol metabolism, tau and amyloid beta-peptide in patients with cognitive impairment.** *Biochem Soc Trans* 2010, **38**:1021–1025.
 36. Morgan EE, Woods SP, Letendre SL, Franklin DR, Bloss C, Goate A, Heaton RK, Collier AC, Marra CM, Gelman BB, McArthur JC, Morgello S, Simpson DM, McCutchan JA, Ellis RJ, Abramson I, Gamst A, Fennema-Notestine C, Smith DM, Grant I, Vaida F, Clifford DB, The CNS HIV Antiretroviral Therapy Effects Research (CHARTER) Group: **Apolipoprotein E4 genotype does not increase risk of HIV-associated neurocognitive disorders.** *J Neurovirol* 2013: [Epub ahead of print].
 37. Letendre S: **Central nervous system complications in HIV disease: HIV-associated neurocognitive disorder.** *Topics in Antiviral Medicine* 2011, **19**:137–142.
 38. Mind exchange working group: **Assessment, diagnosis and treatment of HIV-associated neurocognitive disorder: a consensus report of the mind exchange program.** *Clin Infect Dis* 2013, **56**:1004.
 39. Skinner S, Adewale AJ, DeBlock L: **Neurocognitive screening tools in HIV/AIDS: comparative performance among patients exposed to antiretroviral therapy.** *HIV Med* 2009, **10**:246–252.
 40. Power C, Selnes OA, Grim JA, McArthur JC: **HIV dementia scale: a rapid screening test.** *J Acquir Immune Defic Syndr Hum Retrovirol* 1995, **8**:273–278.
 41. Sacktor NC, Wong M, Nakasujja N, Skolasky RL, Selnes OA, Musisi S, Robertson K, McArthur JC, Ronald A, Katabira E: **The International HIV Dementia Scale: a new rapid retesting test for HIV dementia.** *AIDS* 2005, **19**:1367–1374.
 42. Koski L, Brouillette MJ, Lalonde R, Hello B, Wong E, Tsuchida A, Fellows L: **Computerized testing augments pencil-and-paper tasks in measuring HIV-associated mild cognitive impairment(*).** *HIV Med* 2011, **12**:472–480. Epub 2011 Mar 13.
 43. Zipursky AR, Gogolishvili D, Rueda S, Brunetta J, Carvalho A, McCombe JA, Gill MJ, Rachlis A, Rosenes R, Arbes G, Marcotte T, Rourke SB: **Evaluation of brief screening tools for neurocognitive impairment in HIV/AIDS: a systematic review of the literature.** *AIDS* 2013, **27**:2385–2401.
 44. Carey CL, Woods SP, Rippeth JD, Gonzalez R, Moore DJ, Marcotte TD, Grant I, Heaton RK, HNRC Group: **Initial validation of a screening battery for the detection of HIV-associated cognitive impairment.** *Clin Neuropsychol* 2004, **18**:234–248.
 45. Cysique LA, Maruff P, Darby D, Brew BJ: **The assessment of cognitive function in advanced HIV-1 infection and AIDS dementia complex using a new computerised cognitive test battery.** *Arch Clin Neuropsychol* 2006, **21**:185–194.
 46. Muñoz-Moreno JA, Prats A, Pérez-Álvarez N, Fumaz CR, Garolera M, Doval E, Negro E, Ferrer MJ, Clotet B: **A brief and feasible paper-based method to screen for neurocognitive impairment in HIV-infected patients.** *J Acquir Immune Defic Syndr* 2013, **63**:585–592.
 47. Grupo de expertos de estudio de Sida (GeSIDA) y de la secretaria del plan nacional sobre el SIDA (SPNS): **Documento de consenso sobre el manejo clínico de los trastornos neurocognitivos asociados a la infección por el VIH.** Noviembre 2012.
 48. Brew BJ, Letendre SL: **Biomarkers of HIV related central nervous system disease.** *Int Rev Psychiatry* 2008, **20**:73–88.
 49. Brew BJ: **Markers of AIDS dementia complex: the role of cerebrospinal fluid assays.** *AIDS* 2001, **15**:1883–1884.
 50. Letendre S, Ellis R, Deutsch R, Clifford D, Marra C, McCutchan A, the CHARTER Group: **Correlates of time-to-loss-of-viral-response in CSF and plasma in the CHARTER cohort.** In *17th Conference on Retroviruses & Opportunistic Infections*. 2010. Abstract 430.
 51. Price RW, Epstein LG, Becker JT, Cinque P, Gisslen M, Pulliam L, McArthur JC: **Biomarkers of HIV-1 CNS infection and injury.** *Neurology* 2007, **69**:1781–1788.
 52. Forlenza OV, Diniz BS, Gattaz WF: **Diagnosis and biomarkers of predementia in Alzheimer's disease.** *BMC Med* 2010, **8**:89. Published online 2010 December 22. Review.
 53. Brew BJ, Pemberton L, Blennow K, Wallin A, Hagberg L: **CSF amyloid β 42 and tau levels correlate with AIDS dementia complex.** *Neurology* 2005, **9**:1490–1492.
 54. Ellis RJ, Seubert P, Mottar R, Galasko D, Deusch R, Heaton RK, Heyesd MP, McCutchan JA, Atkinson JH, Grant I, HIV Neurobehavioral Research Center Group: **Cerebrospinal fluid tau protein is not elevated in HIV-associated neurologic disease in humans.** *Neurosci Lett* 1998, **254**:1–4.
 55. Steinbrink F, Evers S, Buerke B, Young P, Arendt G, Koutsilieri E, Reichelt D, Lohmann H, Husstedt IW, German Competence Network HIV/AIDS: **Cognitive impairment in HIV infection is associated with MRI and CSF pattern of neurodegeneration.** *Eur J Neurol* 2013, **20**:420–428.
 56. Letendre SL, McCutchan JA, Childers ME, Woods SP, Lazzaretto D, Heaton RK, Grant I, Ellis RJ, HNRC Group: **Enhancing antiretroviral therapy for human immunodeficiency virus cognitive disorders.** *Ann Neurol* 2004, **56**:416–423.
 57. Robert KH, Franklin DR, Ellis RJ, McCutchan JA, Letendre SL, Leblanc S, Corkran SH, Duarte NA, Clifford DB, Woods SP, Collier AC, Marra CM, Morgello S, Mindt MR, Taylor MJ, Marcotte TD, Atkinson JH, Wolfson T, Gelman BB, McArthur JC, Simpson DM, Abramson I, Gamst A, Fennema-Notestine C, Jernigan TL, Wong J, Grant I, CHARTER Group, HNRC Group: **HIV-associated neurocognitive disorders before and during the era of combination antiretroviral therapy: differences in rates, nature and predictors.** *J Neurovirol* 2010, **10**:3–16.
 58. Mellgren A, Antinori A, Cinque P, Price RW, Eggers C, Hagberg L, Gisslén M: **Cerebrospinal fluid HIV-1 infection usually responds well to antiretroviral treatment.** *Antivir Ther* 2005, **10**:701–707.
 59. Letendre SL, McCutchan JA, Childers ME, HNRC Group, Woods SP, Lazzaretto D, Heaton RK, Grant I, Ellis RJ, HNRC Group: **Enhancing antiretroviral therapy for human immunodeficiency virus cognitive disorders.** *Ann Neurol* 2004, **56**:416–423.
 60. Patel K, Ming X, Williams PL, Robertson KR, Oleske JM, Seage GR 3rd, International Maternal Pediatric Adolescent AIDS Clinical Trials 219/219C Study Team: **Impact of HAART and CNS penetrating antiretroviral regimens on HIV encephalopathy among perinatally infected children and adolescents.** *AIDS* 2009, **23**:1893–1901.
 61. Eden A, Fuchs D, Hagberg L, Nilsson S, Spudich S, Svennerholm B, Price RW, Gisslén M: **HIV-1 viral escape in cerebrospinal fluid of subjects on suppressive antiretroviral treatment.** *J Infect Dis* 2010, **202**:1819–1825.
 62. Letendre S, Marquie-Beck J, Capparelli E, Best B, Clifford D, Collier AC, Gelman BB, McArthur JC, McCutchan JA, Morgello S, Simpson D, Grant I, Ellis RJ, CHARTER Group: **Validation of the CNS penetration-Effectiveness Rank for quantifying antiretroviral penetration into the central nervous system.** *Arch Neurol* 2008, **65**:65–70.
 63. Cusini A, Vernazza PL, Yerly S, Decoster LA, Ledergerber B, Fux CA, Rohrbach J, Widmer N, Hirschel B, Gaudenz R, Cavassini M, Klimkait T, Zenger F, Gutmann C, Opravil M, Günthard HF, Swiss HIV Cohort Study: **Higher CNS penetration-effectiveness of long-term combination antiretroviral therapy is associated with better HIV-1 viral suppression in cerebrospinal fluid.** *JAIDS* 2013, **62**:28–35.
 64. Ene L, Duiculescu D, Ruta SM: **How much do antiretroviral drugs penetrate into the central nervous system?** *J Med Life* 2011, **4**:432–439.
 65. Smurzynski M, Wu K, Letendre S, Robertson K, Bosch RJ, Clifford DB, Evans S, Collier AC, Taylor M, Ellis R: **Effects of central nervous system antiretroviral penetration on cognitive functioning in the ALLRT cohort.** *AIDS* 2011, **25**:357–365.
 66. Sacktor N, Miyahara S, Deng L: **Minocycline treatment for HIV-associated cognitive impairment: results from a randomized trial.** *18th conference on Retroviruses and Opportunistic Infections*; 2011. Abstract 421.
 67. Schifitto G, Navia BA, Yiannoutsos CT, Marra CM, Chang L, Ernst T, Jarvik JG, Miller EN, Singer EJ, Ellis RJ, Kolson DL, Simpson D, Nath A, Berger J, Shriver SL, Millar LL, Colquhoun D, Lenkinski R, Gonzalez RG, Lipton SA, Adult AIDS Clinical Trial Group (ACTG) 301, 700 Teams, HIV MRS Consortium: **Memantine and HIV-associated cognitive impairment: a neuropsychological and proton magnetic resonance spectroscopy study.** *AIDS* 2007, **21**:1877–1886.
 68. Ettenhofer ML, Hinkin CH: **Aging, neurocognition, and medication adherence in HIV infection.** *Am J Geriatr Psychiatry* 2009, **17**:281–290.

69. Hinkin CH, Hardy DJ, Mason KI, Castellon SA, Durvasula RS, Lam MN, Stefaniak M: **Medication adherence in HIV-infected adults: effect of patient age, cognitive status and substance abuse.** *AIDS* 2004, **18**:19–25.
70. Muñoz-moreno JA, Fumaz CR, Ferrer MJ: **Benefits of a cognitive rehabilitation program on neurocognitive impairment in HIV-infected patients. Preliminary finding.** *J Int Neuropsychol Soc* 2007, **13**:S1–S9.
71. Dore GJ, McDonald A, Li Y, National HIV Surveillance Committee: **Marked improvement in survival following AIDS dementia complex in the era of highly active antiretroviral therapy.** *AIDS* 2003, **17**:1539–1545.
72. Sevigny JJ, Albert SM, McDermott MP, Schifitto G, McArthur JC, Sacktor N, Conant K, Selnes OA, Stern Y, McClemon DR, Palumbo D, Kiebertz K, Riggs G, Cohen B, Marder K, Epstein LG: **An evaluation of neurocognitive status and markers of immune activation as predictors of time to death in advanced HIV infection.** *Arch Neurol* 2007, **6**:97–102.
73. Waldrop-valverde D, Jones DL, Gould F, Kumar M, Ownby RL: **Neurocognition, health related reading literacy and numeracy in medication management for HIV infection.** *AIDS Patient Care STDS* 2010, **24**:477–484.
74. Letendre S, Ellis R, Deutsch R, Clifford D, Marra C, McCutchan A: **the CHARTER Group: Correlates of time-to-loss-of-viral-response in CSF and plasma in the CHARTER cohort.** In *17th Conference on Retroviruses & Opportunistic Infections*. 2010. Abstract 430.
75. Scott JC, Woods SP, Vigil O, Heaton RK, Schweinsburg BC, Ellis RJ, Grant I, Marcotte TD, San Diego HIV Neurobehavioral Research Center (HNRC) Group: **A neuropsychological investigation of multitasking in HIV infection: implications for everyday functioning.** *Neuropsychology* 2011, **25**:511–519.
76. Rueda S, Raboud J, Mustard C, Bayoumi A, Lavis JN, Rourke SB: **Employment status is associated with both physical and mental health quality of life in people living with HIV.** *AIDS Care* 2011, **23**:435–443.
77. Tozzi V, Balestra P, Murri R, Galgani S, Bellagamba R, Narciso P, Antinori A, Giulianelli M, Tosi G, Fantoni M, Sampaolesi A, Noto P, Ippolito G, Wu AW: **Neurocognitive impairment influences quality of life in HIV-infected patients receiving HAART.** *Int J STD AIDS* 2004, **15**:254–259.
78. Dore GJ, Correll PK, Li Y, Kaldor JM, Cooper DA, Brew BJ: **Changes to AIDS dementia complex in the era of highly active antiretroviral therapy.** *AIDS* 1999, **13**:1249–1253.
79. Vivithanaporn P, Heo G, Gamble J, Krentz HB, Hoke A, Gill MJ, Power C: **Neurologic disease burden in treated HIV/AIDS predicts survival: a population-based study.** *Neurology* 2010, **75**:1150–1158.
80. Freret T, Gaudreau P, Schumann-Bard P, Billard JM, Popa-Wagner A: **Mechanisms underlying the neuroprotective effect of brain reserve against late life depression.** *J Neural Transmission* 2014: [Epub ahead of print].
81. Shapiro ME, Mahoney JR, Peyser D, Zingman BS, Verghese J: **Cognitive reserve protects against apathy in individuals with human immunodeficiency virus.** *Arch Clin Neuropsychol* 2014, **29**:110–120.

doi:10.1186/2049-9256-2-2

Cite this article as: Sanmarti et al.: HIV-associated neurocognitive disorders. *Journal of Molecular Psychiatry* 2014 **2**:2.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

