LETTER TO THE EDITOR

Quality indicators for primary care and patients' experience: A leap beyond

To the Editor,

Ozaki et al (2019)¹ had evaluated the quality of primary care throughout small facilities using quality indicators (QIs) collected in electronic medical records (EMR) system. They selected 18 QIs based on a comprehensive framework after a modified Delphi method validated by an expert panel members. Besides, the authors also collected data on a patient survey in four dimensions presented in the Appendix: access, communication, coordination, and understanding of patient's background. But, unfortunately, these dimensions are not comparable to other studies in primary health facilities. What do we mean by that? We ask why not use a statistically validated international instrument to assess patients' experience in primary care clinics?

To measure quality of Japan's medical care, one should argue what patients' experience is. They play an important role in the assessment of health services, especially in primary care, where clinics and facilities are usually near each person/family. Starfield and Shi mentioned by the authors on references #1 and #4 had developed and validated a set of questionnaires to evaluate primary healthcare facilities—the Primary Care Assessment Tool (PCAT),² and recently, Aoki et al³ also validated a Japanese adult version, the so-called J-PCAT. It has most of the dimensions considered by the authors, which together allow the measurement of an overall IQ (the overall PCAT score), in addition to allowing the specific calculation for each of the attributes: access—first contact, ongoing care, comprehensiveness (services available), comprehensiveness (services provided), and community orientation.

The use of PCAT is widespread in health services in Brazil, in the states of São Paulo, Rio de Janeiro and especially in Rio Grande do Sul. In agreement with Ozaki et al (2019),¹ when they mention that primary care plays an "increasingly important role in the healthcare system," we propose a leap beyond. Why not outline a national policy for primary care assessment including users' perspective? Thinking about this issue, in early 2019, the Brazilian Ministry of Health sought Brazilian Institute of Geography and Statistics (IBGE)—the Statistics Bureau of Brazil—to include in its main National Health Survey (PNS-2019), the set of questions that forms Brazilian Adult PCAT reduced version. ^{4,5}

Therefore, we encourage the authors in future studies to focus on expanding their research in community clinics in Japan using J-PCAT for the assessment of patients' experience, helping to foster a baseline

to compare nationwide primary care facilities. It could be used not only for health service research, but for a Japanese primary care policy, as Brazilian Federal Government is innovating since 2019, considering it for Family Health Teams (that form the basis of the health system in Brazilian primary care), to evaluate the quality of primary care throughout a part of a pay-for-performance program implemented.

ACKNOWLEDGEMENTS

The authors would like to thank the Brazilian Ministry of Health and Brazilian Institute of Geography and Statistics (IBGE)—the Statistics Federal Bureau of Brazil—to include in the National Health Survey (PNS-2019), PCAT questions.

CONFLICT OF INTEREST

The authors declare no conflict of interests for this article.

Luiz Felipe Pinto PhD^{1,2} Erno Harzheim MD, PhD³
Otávio Pereira D'Avila PhD⁴
Lisiane Hauser PhD⁵

¹Department of Primary Health Care in Medicine, School of Medicine, Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

²Instituto de Higiente e Medicina Tropical (IHMT), Universidade Nova de Lisboa, Nova de Lisboa, Portugal

Email: felipepinto.rio@medicina.ufrj.br

³School of Medicine, Federal University of Rio Grande do Sul (UFRGS), Rio de Janeiro, Brazil

⁴School of Dentist, Federal University of Pelotas, Rio Grande do Sul (UFPel), Rio de Janeiro, Brazil

⁵Brazilian Health Ministry, Rio de Janeiro, Brazil

ORCID

Luiz Felipe Pinto https://orcid.org/0000-0002-9888-606X

REFERENCES

 Ozaki M, Matsumura S, Iwamoto M, Kamitani S, Higashi T, Toyama M et al Quality of primary care provided in community clinics in Japan. J Gen Fam Med. 2019;20:48–54.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. Journal of General and Family Medicine published by John Wiley & Sons Australia, Ltd on behalf of Japan Primary Care Association

204 LETTER TO THE EDITOR

 Johns Hopkins Bloomberg School of Public Health. The Johns Hopkins Primary Care Policy Center. Primary Care Assessment Tools. Baltimore; 2016. [Internet] [cited March 2020]. Available from http://www.jhsph.edu/research/centers-and-institutes/johns-hopkins-primary-care-policy-center/pca_tools.html

- Aoki T, Inouec M, Nakayamad T. Development and validation of the Japanese version of Primary Care Assessment Tool. Fam Pract. 2016;33:112-7.
- 4. Brazil. Brazilian Institute of Geography and Statistics (IBGE). National Health Survey. For the first time, IBGE carries out survey on patient
- check-in in public health system. [Internet] [cited March 2020]. Available from https://agenciadenoticias.ibge.gov.br/en/agencianews/2184-news-agency/news/25542-for-the-first-time-ibge-carri es-out-survey-on-patient-check-in-in-public-health-system
- Brazil. Federal University of Rio Grande do Sul (UFRGS). Launch of Fieldwork of Brazilian National Health Survey (PNS-2019) from Brazilian Institute of Geography and Statistics (IBGE) – Telessaúde, RS - UFRGS. Questionnaire – Primary Health Care – PCAT. [Internet] [cited March 2020]. Available from https://www.ufrgs.br/telessaude rs//documentos/PNS2019_MODULO_H_versao_ingles.pdf.pdf