

Transcriptome analysis and promoter sequence studies on early adipogenesis in 3T3-L1 cells*

Su-jong Kim¹, Ki-Hwan Lee¹, Yong-Sung Lee¹, Eun-Gyeng Mun², Dae-Young Kwon³ and Youn-Soo Cha^{2§}

¹Department of Biochemistry, College of Medicine, Hanyang University, Seoul 133-791, Korea

²Department of Food Science and Human Nutrition, Research Institute of Human Ecology, Chonbuk National University, Jeonju, Jeonbuk 561-756, Korea

³Food Function Research, Division Korea Food Research Institute, San46-1, Baekhyun-dong, Bundang-gu, Sunghnam-si, Gyeonggi-do 463-746, Korea

Received January 22, 2007; Revised March 2, 2007; Accepted March 5, 2007

Abstract

To identify regulatory molecules which play key roles in the development of obesity, we investigated the transcriptional profiles in 3T3-L1 cells at early stage of differentiation and analyzed the promoter sequences of differentially regulated genes. One hundred and sixty-one (161) genes were found to have significant changes in expression at the 2nd day following treatment with differentiation cocktail. Among them, 86 transcripts were up-regulated and 75 transcripts were down-regulated. The 161 transcripts were classified into 10 categories according to their functional roles; cytoskeleton, cell adhesion, immune, defense response, metabolism, protein modification, protein metabolism, regulation of transcription, signal transduction and transporter. To identify transcription factors likely involved in regulating these differentially expressed genes, we analyzed the promoter sequences of up- or - down regulated genes for the presence of transcription factor binding sites (TFBSs). Based on coincidence of regulatory sites, we have identified candidate transcription factors (TFs), which include those previously known to be involved in adipogenesis (CREB, OCT-1 and c-Myc). Among them, c-Myc was also identified by our microarray data. Our approach to take advantage of the resource of the human genome sequences and the results from our microarray experiments should be validated by further studies of promoter occupancy and TF perturbation.

Key Words: Microarray analysis, promoter analysis, adipocyte differentiation, 3T3-L1 cells, transcription factor binding sites

Introduction

Obesity is the most prevalent health problem in developed countries and is a growing problem in developing countries. It serves as a high risk factor for morbidity and mortality from metabolic disorders such as type-2 diabetes, dyslipidemia, hypertension, coronary heart disease, atherosclerosis and numerous types of cancer (Gregoire *et al.*, 1998). Excess fat mass, characterizing obesity, is usually caused by excessive energy storage over a prolonged period of time. At the cellular level, increases in adipose tissue occur as the result of lipid accumulation within pre-existing adipocytes or differentiation of vascular stromal cells within the adipose tissue into adipocytes (Rosen & Spiegelman, 2000).

To identify regulatory molecules which play key roles in the development of obesity, which in turn may serve as better therapeutic targets for obesity and related diseases, numerous investigations have focused on the adipocyte differentiation process. From these studies, adipogenic transcription factors and

adipocyte-specific genes such as members of the CCAAT/enhancer binding protein (C/EBP), peroxisome proliferator-activated receptor (PPAR) families, sterol-regulatory element binding proteins (SREBPs) have been identified. In addition to studies using traditional cloning and gene identification techniques in a "gene by gene" manner, several investigations using microarray were performed on differentiation process of 3T3-L1 preadipocytes into adipocyte (Soukas *et al.*, 2001). As a result of these genome-wide studies, many genes associated with the cell cycle, DNA replication, metabolism, signal transduction, cytoskeleton, protein turnover and splicing factors as well as some genes involved in transcriptional regulation were identified as genes differentially regulated during adipocyte differentiation (Burton *et al.*, 2002; Guo & Liao, 2000; Ron & Habener, 1992; Soukas *et al.*, 2001). Despite of a mountain of gene expression profiles, an understanding of the complete mechanism by which precursor cells become adipocytes is far from complete and the genetic regulatory network underlying adipogenesis has not yet been fully understood. For example, many genes that mark the

* This work (M10510120001-05N1012-00110) was supported by research grants from Korea Science and Engineering Foundation (KOSEF) for Biofoods Research Program, Ministry of Science and Technology.

§ Corresponding Author: Youn-Soo Cha, Tel. 82-63-270-3822, Fax. 82-63-270-3854, Email. cha8@chonbuk.ac.kr

differentiated adipocyte were expressed with different kinetics than SREBPs, C/EBPs, and PPARs. Genes known to have a PPAR γ response element in its promoter, such as insulin-like growth factor-II, β 3-adrenergic receptor and the phosphoenolpyruvate carboxykinase, are expressed only late in differentiation at day 4 and maximally at day 28, which is not consistent with the initial expression pattern of PPAR γ (Soukas *et al.*, 2001). Cytochrome *c* oxidase VIIIH, glucose-6-phosphate isomerase and phosphofructokinase I also demonstrated a similar pattern of expression, however, their pattern differed from the expression profile of the aforementioned transcription factors. These observations lead to the speculation that potentially many other regulatory networks are involved in regulating their expression.

In this study, we investigated the transcriptional profiles in 3T3-L1 cells at the early stage of differentiation and analyzed the promoter sequences of differentially regulated genes for identifying common transcription regulators. This approach might provide useful information to establish genetic regulatory network from gene expression profiles.

Materials and Methods

Materials

All tissue culture materials were purchased from Gibco (Grand Island, NY, U.S.A.). Insulin, dexamethasone and MIX were from Sigma (St. Louis, MO, U.S.A.).

Cell culture

3T3-L1 preadipocytes (A.T.C.C., Manassas, VA, U.S.A.) were cultured in Dulbecco's modified Eagle's medium (DMEM) with 10% (v/v) foetal bovine serum (FBS), 1% (v/v) penicillin-streptomycin (100 units/ml penicillin, 100 μ g/ml streptomycin in 0.85% saline) and 1% (v/v) 100 M pyruvate at 37°C in 95% air, 5% CO₂. To induce differentiation, 2-day postconfluent 3T3-L1 preadipocytes (day 0) were stimulated for 72 h with 0.5 mM MIX, 0.5 M dexamethasone and 10 μ g/ml insulin (MDI) added to DMEM/10% FBS culture medium. On day 3, the MDI medium was replaced with DMEM/10% FBS, which was changed every 2 days thereafter until analysis.

Oil Red O staining

3T3-L1 preadipocyte and adipocyte cells were stained with Oil Red O essentially as described by Ramirez-Zacarias, Castro-Munozledo, and Kuri-Harcuch (Ramirez-Zacarias *et al.*, 1992). Briefly, the 3T3-L1 cells were fixed in 10% formalin for 90 min. After washing thoroughly with distilled water, cells were incubated with a working solution of Oil Red O for 3 h. The staining of lipid droplets in 3T3-L1 preadipocyte and adipocyte cells was

quantified using a phase-contrast microscope and Image-Pro Plus software from Media Cybernetics (Carlsbad, CA).

Quantitative analysis of intracellular triglyceride (TG) accumulation

For analysis of the intracellular TG content, the cells were washed with PBS and harvested using lysis buffer (50 mM Tris, 1 mM EDTA, 1 mM beta-mercaptoethanol, 10 mM MgCl₂, 0.15 mM NaCl, and 1 mM phenylmethylsulfonyl fluoride, pH 7.5). Subsequent to sonicating the samples, spectroscopic quantification of TG was performed using the enzyme coupled glycerol assay kit 337 (Sigma Diagnostics, Deisenhofen, Germany). The protein content was determined by the method of Bradford using bovine serum albumin (BSA) as the standard (Bradford, 1976). TG content was expressed as μ g TG per g protein.

Microarray analysis

Mouse 10K cDNA microarray used in this study consisted of 10,336 spots, as previously described (Ahn *et al.*, 2004). It included 6531 transcripts from the National Institute of Aging (NIA), 1243 transcripts from the Brain Molecular Anatomy Project (BMAP), 2060 transcripts from InCyte Pharmaceuticals (Fremont, CA, USA), and yeast DNA and housekeeping genes as negative control.

Total RNA was prepared from cells using Trizol (Invitrogen, Carlsbad, CA, USA). Fluorescence-labeled cDNA probes were prepared from 20 μ g of total RNA by using an amino-allyl cDNA labeling kit (Ambion, Austin, Texas, USA). At least four replicates were performed, and two of these were repeated with the fluorophores reversed to eliminate false-positive results. The Cy5 and Cy3 probes were mixed, and hybridization was performed at 55°C for 16 h, as previously described (Ahn *et al.*, 2004).

The two fluorescent images (Cy3 and Cy5) were scanned separately by a GMS 418 Array Scanner (Affymetrix, Santa Clara, CA, USA), and the image data were analyzed using ImaGene 4.2 (Biodiscovery, Santa Monica, CA, USA) and MAAS (Gaiagene, Seoul, Korea) software (Kim *et al.*, 2001). For each hybridization, emission signal data were normalized by multiplying the Cy3 signal values by the ratio of the means of the Cy3 and Cy5 signal intensities for all spots on the array. The criteria to eliminate unreliable data were previously described (Kim *et al.*, 2004).

In this analysis, we calculated the median value of gene expression ratio from four independently repeated microarray experiments. We used the modified *t*-tests, SAM method to evaluate statistical significance of changes in gene expression (Tusher *et al.*, 2001). We took the genome-wide significance level at the SAM (δ)=0.66 and adopted a cutoff of 2.0-fold change based on our experiences. Genes showing significant differences in expression levels were classified into different functional categories, based on Gene Ontology (GO) with

modifications.

Real-time quantitative PCR analysis and melting curve analysis

Four micrograms of total RNA was reverse-transcribed in 25 μ l of reaction mixture, containing 2.5 U MuLV reverse transcriptase, 1 U RNase inhibitor, 5 mM MgCl₂, 50 mM KCl, 10 mM Tris-HCl (pH 8.3), 2.5 μ M oligo (dT) primer, and 1 mM dNTPs. The reaction mixture was heated to 42°C for 60 min and then denatured at 85°C for 5 min. cDNA was amplified with ICycler (BioRad, Hercules, CA, USA) in 50 μ l of reaction mixture containing AmpliTaq DNA polymerase (0.04 U/ μ l, Perkin Elmer, Shelton, Connecticut), 50 mM Tris (pH 8.3), 3 mM MgCl₂, 0.25 mM dNTPs, 1/50,000 dilution of SYBR Green I (Molecular Probes, Eugene, OR), and 0.25 μ M appropriate sense and antisense PCR primers. The sequences of the primers were as follows: fatty acid synthase (*Fasn*) forward, 5'-TCC ACC TTT AAG TTG CCC TG-3', reverse, 5'-TCT GCT CTC GTC ATG TCA CC-3' aldolase1 forward, 5'-CTA CAA GGC TCT GAG CGA CC-3', reverse, 5'-ACA GGA AAG TGA CCC CAG TG-3' carnitine palmitoyltransferase 2 (*Cpt2*) forward, 5'-CAC AAC ATC CTG TCC ACC AG-3', reverse, 5'-CAT TGC AGC CTA TCC AGT CA-3' peroxisome proliferator-activated receptor γ (*Pparg*) forward, 5'-CAA GAC TAC CCT TTA AGT GAA-3', reverse, 5'-CTA CTT TGA TCG CAC TTT GGT-3' Iroquois related homeobox 5 (*Drosophila*) (*Irx5*) forward, 5'-CTG TCC CGG CCT CTT TAT TAC A-3', reverse, 5'-ACC GTC TGG TTT AAT CCA TTG A-3' transforming growth factor, beta 3 (*Tgfb3*) forward, 5'-CCT GGC CCT GCT GAA CTT G-3', reverse, 5'-GAC GTG GGT CAT CAC CGA T-3' Transforming growth factor, beta receptor II (*Tgfb2*) 5'-CCG CTG CAT ATC GTC CTG TG-3', reverse, 5'-AGT GGA TGG ATG GTC CTA TTA CA-3' Mitogen-activated protein kinase 6 (*Mapk6*) 5'-CAT GGA CTT AAA ACC ATT GGG CT-3', reverse, 5'-TGA GGA CAA TTT TCT TGA TGG CT-3' Metallothionein 1 (*Mt1*) 5'-AAG AGT GAG TTG GGA CAC CTT-3', reverse, 5'-CGA GAC AAT ACA ATG GCC TCC-3' Metallothionein 2 (*Mt2*) 5'-GCC TGC AAA TGC AAA CAA TGC-3', reverse, 5'-AGC TGC ACT TGT CGG AAG C-3' Caveolin, caveolae protein 1 (*Cav1*) 5'-ATG TCT GGG GGC AAA TAC GTG-3', reverse, 5'-CGC GTC ATA CAC TTG CTT CT-3' Fatty acid binding protein 4, adipocyte (*Fabp4*) 5'-AAG GTG AAG AGC ATC ATA ACC CT-3', reverse, 5'-TCA CGC CTT TCA TAA CAC ATT CC-3' Serine (or cysteine) peptidase inhibitor, clade F, member 1 (*Serpinf1*) 5'-GCC CTG GTG CTA CTC CTC T-3', reverse, 5'-CGG ATC TCA GGC GGT ACA G-3' Serine (or cysteine) peptidase inhibitor, clade E, member 2 (*Serpine2*) 5'-CAC ATG GGA TCG CGT CCA TC-3', reverse, 5'-CAG CAC TTT ACC AAC TCC GTT TA-3' Activating transcription factor 4 (*Atf4*) 5'-ATG GCG CTC TTC ACG AAA TC-3', reverse, 5'-ACT GGT CGA AGG GGT CAT CAA-3' Activating transcription factor 5 (*Atf5*) 5'-CTT CTC TGA TTG GAT GAC TGA GC-3', reverse, 5'-CCT TCT TGA GTA GGG ATG CCA

T-3' Early growth response 1 (*Egr1*) 5'-GTT ATC CCA GCC AAA CGA CTC-3', reverse, 5'-GGT TCA GGC CAC AAA GTG TT-3' β -actin forward, 5'-GGG TCA GAA GGA CTC CTA TG-3' reverse, 5'-GTA ACA ATG CCA TGT TCA AT-3' The following cycling conditions were used: one denaturing cycle at 95°C for 5 min, followed by 30 cycles of 95°C for 30 s, 60°C for 45 s, and 72°C for 1 min. Relative RNA levels were determined by analyzing the changes in SYBR Green I fluorescence during PCR according to the manufacturer's instructions. β -actin was amplified in parallel and the results were used for normalization. The correct size of PCR product was confirmed by electrophoresis on a 2% agarose gel stained with ethidium bromide. Purity of the amplified PCR products was determined by melting point analysis using the ICycler software.

Promoter analysis

Promoter analysis was performed using PAINT v3.0 as previously described (Addya *et al.*, 2004). PAINT v3.0 contains a database of promoters (UpstreamDB) constructed for all the annotated genes (known and putative) in the Ensembl genome database for *Mus musculus*, containing 24,261 genes (Clamp *et al.*, 2003). Promoter analysis for putative TREs, i.e., binding sites for known transcription factors (TFs) was performed by MATCH tool using TRANSFAC database version 7.2. For each gene, 2,000 base pairs (bp) upstream (5' to the transcription start site) were analyzed. For each group of genes analyzed, the *P* values for significance of enrichment of each TRE in that group were calculated using hypergeometric distribution, by comparing the abundance of each TRE to that from a reference set of randomly selected genes.

Results

Histological and biochemical analysis of adipocyte differentiation

The 3T3-L1 preadipocytes differentiated to adipocytes in response to the administration of dexamethasone, indomethacin, 3-isobutyl-1-methyl-xanthine, and insulin as previously described (Pittenger *et al.*, 1999). As shown in Fig. 1(a), a few cells accumulating lipid vesicles were observed at the 2nd day following stimulation, and then the lipid droplet-containing cell population was increased in a time-dependent manner up to day 6. Accumulation of triglycerides in cells was also increased in a time-dependent manner up to day 6 (Fig. 1(b)).

Identification of genes differentially regulated during 3T3-L1 preadipocyte differentiation

To identify genes differentially regulated during 3T3-L1




Fig. 1. Histological and biochemical analysis of adipocyte differentiation Post-confluent 3T3-L1 cells were hormonally treated with differentiation cocktail (1 μ M dexamethasone, 5 μ g/ml insulin, and 0.5 mM IBMX) for 6 days as described in the Methods section. (a) The cells were fixed at the indicated time points and stained with Oil Red O to assess lipid accumulation (100X magnification). (b) (c) Data of triglycerides and protein are expressed as mean \pm S.D (n=3). Asterisks denote significant difference (ANOVA) between the control and differentiation cocktail treatment on days 2, 4 and 6 ($p < 0.05$).

preadipocyte differentiation, about 10,000 gene expression levels in differentiation cocktail-treated 3T3-L1 cells were compared with those of vehicle-treated cells as control. Only the genes, whose mRNA levels were changed 2.0-fold or higher and detected as significant change by SAM method, were designated as differentially expressed genes (Fig. 2). By these criteria, 161 genes were found to have significant changes in expression at the 2nd day following treatment with differentiation cocktail (Table 1, 2). Of these 161 transcripts, 86 transcripts were up-regulated and 75 transcripts were down-regulated. The 161 transcripts were classified into 10 categories according to their functional roles: cytoskeleton, cell adhesion, immunity, defense response, metabolism, protein modification, protein metabolism, regulation of transcription, signal transduction and transporter (Fig. 3).

Verification of the microarray results with Real-time RT-PCR

To validate the differential gene expression revealed by cDNA microarray-based profiling of 3T3-L1 adipogenesis, real-time quantitative RT-PCR was carried out for several selected genes; fatty acid synthase (*Fasn*), aldolase1, carnitine palmitoyltransferase 2 (*Cpt2*), peroxisome proliferator-activated receptor γ (*Pparg*), Iroquois related homeobox 5 (*Drosophila*) (*Irx5*), transforming growth factor, beta 3 (*Tgfb3*), Transforming growth factor, beta receptor II (*Tgfb2*), Mitogen-activated protein kinase 6 (*Mapk6*), Metallothionein 1 (*Mt1*), Metallothionein 2 (*Mt2*), Caveolin, caveolae protein 1 (*Cav1*), Fatty acid binding protein 4, adipocyte (*Fabp4*), Serine (or cysteine) peptidase inhibitor,





Fig. 2. Representative MA plots and SAM plot (a) Representative MA plots comparing the 3T3-L1 preadipocytes vs. the differentiated cells, M represents the log ratio of the two fluorescent dyes used to label probes, and A represents averaged logarithmic intensity. Broken line represents a 2-fold change. (b) SAM scatter plot of observed relative difference versus the expected relative difference. The genes showing significant difference in expression between the 3T3-L1 preadipocytes and the differentiated cells were identified, Broken line represents $\delta = 0.66$.




Fig. 3. Global gene expression profile in functional categories Black bars and white bars represent the percentage of induced and repressed genes, respectively.

Table 1. Genes which were upregulated during adipogenesis

Genebank number	Gene name	Symbol	Fold change				
Cytoskeleton and cell adhesion				BE944401	Forkhead box B2	Foxb2	3.9
Al849746	Talin 1	Tln1	2.2	Al843304	CCAAT/enhancer binding protein (C/EBP), alpha	Cebpa	3.0
BG084556	Myosin Vb	Myo5b	2.3	BG064727	Alpha thalassemia/mental retardation syndrome X-linked homolog (human)	Atrx	2.6
W67022	Keratin complex 1, acidic, gene 10	Krt1-10	2.9	Signal transduction			
BG069796	Kelch-like 2, Mayven (Drosophila)	Klh2	2.9	Al509976	Transforming growth factor, beta receptor III	Tgfb3	2.2
Al844265	Integrin beta 1 binding protein 1	Itgb1bp1	3.0	BG085088	Transforming growth factor, beta receptor II	Tgfb2	2.2
BE948546	Glycoprotein 5 (platelet)	Gp5	2.7	Al849214	Transforming growth factor, beta 3	Tgfb3	3.5
BG086102	Chondroitin sulfate proteoglycan 2	Cspg2	2.3	BG065450	Sorbin and SH3 domain containing 1	Sorbs1	3.2
Immune and defense response				BG080390	Regulator of G-protein signaling 2	Rgs2	6.2
Al852793	Superoxide dismutase 2, mitochondrial	Sod2	2.2	Al850306	Receptor (calcitonin) activity modifying protein 3	Ramp3	2.0
W09930	Lipopolysaccharide binding protein	Lbp	2.3	BG069499	PTK2 protein tyrosine kinase 2	Ptk2	2.1
AW494715	Interferon activated gene 203	Ifi203	2.1	AU021253	Protein phosphatase 1A, magnesium dependent, alpha isoform	Ppm1a	2.2
AA152885	Chemokine (C-X-C motif) ligand 13	Cxcl13	4.6	BG072253	Mitogen-activated protein kinase 6	Mapk6	2.2
Metabolism				BG063925	Metallothionein 2	Mt2	3.0
BG075608	Triosephosphate isomerase 1	Tpi1	2.2	Al427514	Metallothionein 1	Mt1	2.3
BG075934	Transaldolase 1	Taldo1	2.3	BG084827	Insulin-like growth factor binding protein 4	Igfbp4	3.5
BG086835	Sterol carrier protein 2, liver	Scp2	2.2	BG088548	Growth arrest specific 6	Gas6	3.2
Al853169	Stearoyl-Coenzyme A desaturase 1	Scd1	6.1	BE943948	Extra cellular link domain-containing 1	Xlkd1	3.1
BG065475	S-adenosylhomocysteine hydrolase	Ahcy	2.2	AA466094	Cell death-inducing DFFA-like effector c	Cidec	3.2
BG078664	Pyruvate kinase, muscle	Pkm2	2.8	AA138693	Caveolin, caveolae protein 1	Cav1	2.1
AA038254	Pyruvate carboxylase	Pcx	3.2	W13905	Angiotensin-like 4	Angptl4	3.4
BG064823	Phosphoglycerate mutase 1	Pgam1	2.3	AA106031	A kinase (PRKA) anchor protein 4	Akap4	3.6
Al850456	NAD(P) dependent steroid dehydrogenase-like	Nsdhl	2.0	Transporter			
BG075331	Myo-inositol 1-phosphate synthase A1	Isyna1	2.8	AA260521	Uncoupling protein 2 (mitochondrial, proton carrier)	Ucp2	2.4
AW494998	Lipase, hormone sensitive	Lipe	2.1	AA137517	Syntaxin binding protein 2	Stxbp2	2.2
Al835385	Lactate dehydrogenase A	Ldha	3.9	BG070928	Solute carrier family 1 (neutral amino acid transporter), member 5	Slc1a5	2.4
BG076460	Glutamate-cysteine ligase, catalytic subunit	Gclc	3.3	AW121314	NTF2-related export protein 1	Nxt1	2.0
Al835583	Glutamate-ammonia ligase (glutamine synthetase)	Glul	9.2	AA087193	Lipocalin 2	Lcn2	11.4
AA011759	Glutamate-ammonia ligase (glutamine synthetase)	Glul	3.6	AA213017	Flavin containing monooxygenase 3	Fmo3	2.6
BG077710	Galactose-4-epimerase, UDP	Gale	2.1	BE943530	Fatty acid binding protein 4, adipocyte	Fabp4	19.0
AA116513	Fatty acid synthase	Fasn	3.6	BG073601	Diazepam binding inhibitor	Dbi	2.1
W11965	Enolase 3, beta muscle	Eno3	2.2	BG078810	Cytochrome c, somatic	Cycc	2.5
Al847556	Diacylglycerol O-acyltransferase 2	Dgat2	4.8	BG063100	Cytochrome c, somatic	Cycc	2.2
Al605638	Diacylglycerol O-acyltransferase 1	Dgat1	2.1	Unknown			
BG071627	DDHD domain containing 1	Ddhd1	2.1	BG087501	Zinc finger CCHC-type and RNA binding motif 1	Zcrb1	2.2
BG085333	Carnitine palmitoyltransferase 2	Cpt2	2.4	Al850717	Transcribed locus		2.4
AA518639	Aldolase 1, A isoform	Aldoa	2.4	BG071626	Similar to Glyceraldehyde-3-phosphate dehydrogenase (GAPDH)	LOC382450	2.8
BG074754	Acyl-CoA synthetase long-chain family member 1	Acs1	2.1	AA122891	Similar to glyceraldehyde-3-phosphate dehydrogenase	LOC380687	2.5
Protein modification and metabolism				AA137748	RIKEN cDNA 1500003O03 gene	1500003O03 Rik	2.2
AA276457	Renin 1 structural	Ren1	2.1	BG088778	Microsomal glutathione S-transferase 3	Mgst3	2.5
BG062950	Prolyl endopeptidase	Prep	2.7	Al844280	Lymphocyte antigen 6 complex, locus A	Ly6a	4.3
BG087373	FK506 binding protein 5	Fkbp5	3.2	W83922	Leucine-rich repeats and immunoglobulin-like domains 1	Lrig1	2.3
AA259979	Angiotensinogen (serpin peptidase inhibitor, clade A, member 8)	Agt	2.8	BG065088	DEAD (Asp-Glu-Ala-Asp) box polypeptide 21	Ddx21	2.3
Regulation of transcription				BG065476	CDNA sequence AK190093	AK190093	2.3
Al843911	TSC22 domain family 3	Tsc22d3	4.7	BG065432	ATP-binding cassette, sub-family F (GCN20), member 3	Abcf3	2.5
AA511029	Peroxisome proliferator activated receptor gamma	Pparg	3.1				
AA267605	Peroxisome proliferator activated receptor gamma	Pparg	2.7				
BG078091	NMDA receptor-regulated gene 1	Narg1	2.5				
AA009268	Myelocytomatosis oncogene	Myc	2.5				
AW492717	Iroquois related homeobox 5 (Drosophila)	Irx5	4.8				

Table 2. Genes which were downregulated during adipogenesis

Genebank number	Gene name	Symbol	Fold change		
Cytoskeleton and cell adhesion					
AA423149	Cysteine rich protein 61	Cyr61	-2.5		
AA108928	Secreted phosphoprotein 1	Spp1	-3.1		
BE947634	Vascular cell adhesion molecule 1	Vcam1	-3.0		
BG087410	CD9 antigen	Cd9	-2.2		
AI838607	Thrombospondin 1	Thbs1	-3.4		
AA003904	Thrombospondin 2	Thbs2	-2.6		
AI323131	Thrombospondin 3	Thbs3	-2.5		
BG077878	Caldesmon 1	Cald1	-2.4		
W64636	Calponin 1	Cnn1	-2.1		
AA220193	Calponin 2	Cnn2	-3.8		
W09925	Filamin C, gamma (actin binding protein 280)	Finc	-2.0		
AW557788	Filamin, alpha	Fina	-2.2		
W96914	Lysyl oxidase	Lox	-3.8		
BG087143	Microfibrillar-associated protein 2	Mfap2	-2.1		
BG087200	Myosin, light polypeptide 6, alkali, smooth muscle and non-muscle	Myl6	-2.1		
W81878	Periostin, osteoblast specific factor	Postn	-2.0		
BG087142	Procollagen, type I, alpha 2	Col1a2	-4.3		
W89883	Procollagen, type III, alpha 1	Col3a1	-2.6		
AA034564	Procollagen, type V, alpha 2	Col5a2	-2.5		
AI894006	Procollagen, type XI, alpha 1	Col11a1	-2.4		
BG064802	Secreted acidic cysteine rich glycoprotein	Sparc	-2.6		
AI835403	Thymosin, beta 4, X chromosome	Tmsb4x	-2.5		
AA444490	Tissue inhibitor of metalloproteinase 2	Timp2	-2.5		
BG086016	Tropomyosin 1, alpha	Tpm1	-2.7		
Immune and defense response					
AA213167	Heme oxygenase (decycling) 1	Hmox1	-2.5		
AI846176	Histocompatibility 2, D region locus 1	H2-D1	-2.5		
AA033050	Serine (or cysteine) peptidase inhibitor, clade E, member 2	Serpine2	-2.6		
BG087320	Serine (or cysteine) peptidase inhibitor, clade F, member 1	Serpinf1	-5.2		
BG086364	Serine (or cysteine) peptidase inhibitor, clade H, member 1	Serpinh1	-2.1		
AA109951	Beta-2 microglobulin	B2m	-5.4		
BG087171	Complement component 1, r subcomponent	C1r	-2.6		
BG069444	Complement component 1, s subcomponent	C1s	-2.1		
Metabolism					
BG074397	Glutathione S-transferase, mu 1	Gstm1	-2.1		
BG064825	Hexosaminidase A	Hexa	-2.1		
BG069782	Hydroxysteroid (17-beta) dehydrogenase 4	Hsd17b4	-2.5		
BG071742	N-acylsphingosine amidohydrolase 1	Asah1	-3.1		
AA060268	Phospholipase D family, member 3	Pld3	-2.3		
BG076063	Phosphorylase kinase alpha 2	Phka2	-2.5		
Protein modification and metabolism					
W78651	Cystatin C	Cst3	-2.4		
BG074453	Lysosomal membrane glycoprotein 2	Lamp2	-2.0		
AA119072	Tripeptidyl peptidase I	Tpp1	-2.1		
AI836324	Carboxypeptidase E	Cpe	-2.4		
W14837	Legumain	Lgmn	-2.2		
BG087474	Procollagen C-endopeptidase enhancer protein	Pcolce	-2.0		
Regulation of transcription					
AF252627	Activating transcription factor 4	Atf4	-2.1		
AI851042	Activating transcription factor 5	Atf5	-2.2		
AA178779	Interferon regulatory factor 8	Irf8	-3.4		
AA059930	Nuclear factor I/B	Nfib	-2.3		
W14113	Twist gene homolog 1 (Drosophila)	Twist1	-3.4		
BG070825	Early growth response 1	Egr1	-2.3		
AA023645	Four and a half LIM domains 2	Fhl2	-3.7		
Signal transduction					
AA000945	DNA segment, human D4S114	D0H4S114	-4.2		
BG065503	Integrin beta 5	Itgb5	-2.9		
AI836468	Myristoylated alanine rich protein kinase C substrate	Marcks	-3.4		
BG063995	Natriuretic peptide precursor type B	Nppb	-3.2		
W42321	Pentraxin related gene	Ptx3	-2.5		
AA049699	Pleiotrophin	Ptn	-9.3		
AA038395	Ras suppressor protein 1	Rsu1	-2.4		
AI848233	Sprouty homolog 2 (Drosophila)	Spry2	-2.0		
AI840211	A kinase (PRKA) anchor protein 8-like	Akap8l	-2.8		
Transporter					
BG076932	Annexin A1	Anxa1	-2.9		
AA002439	Annexin A5	Anxa5	-3.1		
AA499296	Annexin A6	Anxa6	-2.2		
AI837042	Apolipoprotein D	Apod	-2.5		
AA498131	Aquaporin 8	Aqp8	-2.3		
AU022767	Exportin 4	Xpo4	-2.1		
BG074498	Thiosulfate sulfurtransferase, mitochondrial	Tst	-2.0		
BG085465	Nucleoredoxin	Nxn	-2.0		
Unknown					
BG065334	0 day neonate cerebellum cDNA, RIKEN full-length enriched library, clone: C230070N19 product:myristoylated alanine rich protein kinase C substrate		-3.2		
AA407811	Delta-like 1 homolog (Drosophila)	Dlk1	-3.3		
W18828	Dihydropyrimidinase-like 3	Dpysl3	-2.5		
AA250039	Lectin, galactose binding, soluble 9	Lgals9	-2.1		
W82141	Lysosomal membrane glycoprotein 1	Lamp1	-2.1		
AI836519	Myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, Drosophila); translocated to, 11	Mllt11	-3.6		
AI894053	Reticulocalbin 1	Rcn1	-2.7		

clade F, member 1 (*Serpinf1*), Serine (or cysteine) peptidase inhibitor, clade E, member 2 (*Serpine2*), Activating transcription factor 4 (*Atf4*), Activating transcription factor 5 (*Atf5*) and Early growth response 1 (*Egr1*). These genes were chosen from the metabolism, transcriptional regulation, signal transduction and protein modification categories based on the GO terms because these cellular processes have been reported to be important in adipogenesis (Gregoire *et al.*, 1998; Heath *et al.*, 2000; Ntambi & Kim, 2000; Rosen *et al.*, 2000; Soukas *et al.*, 2001; Yoon *et al.*, 2001). The expressions of *Fasn* and *Cpt2* involved in metabolism were increased by 4.6- and 2.4-fold during adipocyte differentiation, respectively. Transcription factors, *Pparg* and *Irx5* were also increased by 3.5- and 4.2-fold. Several genes that play roles in signal transduction and transport also showed expression patterns similar to those described above *Tgfb3*




Fig. 4. Comparison between cDNA microarray analysis and real time RT-PCR. 3T3-L1 cells were treated with differentiation cocktail for two days and total RNA was isolated. The relative mRNA expression levels of genes were measured by real-time RT-PCR analysis with SYBRO green. Data are expressed as fold changes (means \pm SD), normalized to β -actin mRNA expression, where the values for the 3T3-L1 preadipocytes were set at 1.00. The analyses were performed in triplicate,

(3.5-fold), *Tgfb2* (2.2-fold), *Mapk6* (2.2-fold), *Mt1* (1.7-fold) *Mt2* (3.3-fold), *Cav1* (2.1-fold) and *Fabp4* (27-fold). On the other hand, *Serpine1*, *Serpine2*, *Atf4*, *Atf5* and *Egr1* were decreased after treatment with differentiation cocktail by 28, 19, 28, 40, and 35% (Fig. 4).

In general, when gene expression profiles obtained by both microarray analysis and RT-PCR were compared, their patterns were very similar with regards to the direction (up- or down-regulation) and degree of differences in expression. These observations confirmed that the microarray data were reliable.


Promoter analysis

Promoter analysis was performed for the up- or down-regulated genes using PAINT v3.0 to identify biologically relevant transcription factor binding sites or TFBSs, found in the regulatory regions of these genes. For each gene differentially

Table 3. Statistically over-represented TF binding sites in the genes differentially expressed during adipogenesis

TFs	p-value
TFs overrepresented in the promoters of the upregulated genes	
AP-1/V\$AP1_Q4	0.00206
HEN1/V\$HEN1_01	0.03719
HEN1/V\$HEN1_02	0.01779
NF-kappaB (p65)/V\$NFKAPPAB65_01	0.02393
NF-kappaB/V\$NFKAPPAB_01	0.04667
Oct-1/V\$OCT1_07	0.03527
c-Myc/Max/V\$MYCMAX_02	0.01402
c-Rel/V\$CREL_01	0.0125
TFs overrepresented in the promoters of the downregulated genes	
Tax/CREB/V\$TAXCREB_01	0.04228
v-Myb/V\$VMYB_01	0.03555

(a) Selected TFBSs from upregulated genes



(b) Selected TFBSs from downregulated genes




Fig. 5. Matrix of genes differentially expressed during adipogenesis and TF binding sites statistically over-represented in their promoters.

expressed during adipocyte differentiation, the regions which fell within 2000 base pairs (bp) upstream of the transcription start site (TSS) were analyzed. Table 3 shows TFBSs overrepresented in the promoters of the up- or down-regulated genes and *P*-values. We adopted *P*-value cutoff of 0.05 to select over-represented TFBSs. The binding sites of AP-1, HEN1, NF-kappaB (p65), NF-kappaB, Oct-1, c-Myc-Max and c-Rel were selected as over-represented TFBSs for genes increased during adipocyte differentiation. NGFI-C, Tax/CREB and v-Myb binding sites were selected as over-represented TFBSs for genes decreased during adipogenesis (Table 3, Fig. 5).

Discussion

Differentiation toward adipocytes in differentiation cocktail-treated 3T3-L1 cells was evidenced by lipid accumulation in the cells coupled with up-regulation in mRNAs characteristic of adipocytes [Lipase, hormone sensitive (*Lipe*), Fatty acid synthase (*Fasn*), *Pparg*, and CCAAT/enhancer binding protein (*C/EBP*), alpha (*Cebpa*)] (Gregoire *et al.*, 1998; Heath *et al.*, 2000; Ntambi & Kim, 2000; Rosen *et al.*, 2000; Soukas *et al.*, 2001; Yoon *et al.*, 2001).

At the early differentiation stage, genes involved in cytoskeleton and cell adhesion were affected. Talin 1 (*Tln1*), Myosin Vb (*Myo5b*), Keratin complex 1, acidic, gene 10 (*Krt1-10*), Kelch-like 2, Mayven (Drosophila) (*Klhl2*), Integrin beta 1 binding protein 1 (*Itgb1bp1*), Glycoprotein 5 (platelet) (*Gp5*) and Chondroitin sulfate proteoglycan 2 (*Cspg2*) were up-regulated over 2-fold. Otherwise, Thrombospondin 1, 2, and 3, Caldesmon 1, Calponin 1 and 2, Procollagen I, III, V, and XI, and tissue inhibitor of metalloproteinase 2 (*Timp2*) were down-regulated over 2-fold during early adipogenesis. The differentiation of 3T3-L1 cells involves a dramatic alteration in cell morphology from a flattened fibroblast to a spherical adipocyte. Therefore, dynamic changes in expression of genes involved in cytoskeleton and cell adhesion are not unexpected during the differentiation process.

Genes associated with immune and defense response were also differentially regulated; Superoxide dismutase 2, mitochondrial (*Sod2*), Lipopolysaccharide binding protein (*Lbp*), Interferon activated gene 203 (*Ifi203*) and Chemokine (C-X-C motif) ligand 13 (*Cxcl13*) increased over 2-fold. Heme oxygenase (decycling) 1 (*Hmox1*), Histocompatibility 2, D region locus 1 (*H2-D1*), Serine (or cysteine) peptidase inhibitor, clade E, member 2 (*Serpine2*), Serine (or cysteine) peptidase inhibitor, clade F, member 1 (*Serpinf1*), Serine (or cysteine) peptidase inhibitor, clade H, member 1 (*Serpinh1*), Beta-2 microglobulin (*B2m*), Complement component 1, r subcomponent (*C1r*) and Complement component 1, s subcomponent (*C1s*) decreased.

Genes involved in metabolism, such as Triosephosphate isomerase 1 (*Tpi1*), Transaldolase 1 (*Taldo1*), Sterol carrier protein 2, liver (*Scp2*), Stearoyl-Coenzyme A desaturase 1 (*Scd1*), Pyruvate kinase, muscle (*Pkm2*), Pyruvate carboxylase (*Pcx*), Phosphoglycerate mutase 1 (*Pgam1*), NAD(P) dependent steroid dehydrogenase-like (*Nsdhl*), Myo-inositol 1-phosphate synthase A1 (*Isyna1*), Glutamate-cysteine ligase, catalytic subunit (*Gclc*), Diacylglycerol O- acyltransferase 1 and 2 (*Dgat1*, *Dgat2*), carnitine palmitoyltransferase 2 (*Cpt2*) and acyl-CoA synthetase long-chain family member 1 (*Acs1l*), showed increased expression patterns. Increase in expression levels of *Scd1*, *Pcx*, *Lipe*, *Fasn* and *Pparg* genes is consistent with the previous reports (Heath *et al.*, 2000; Soukas *et al.*, 2001).

In addition to *Pparg* and *Cebpa* mentioned above, 13 genes involved in regulation of transcription were affected during early adipogenesis; TSC22 domain family 3 (*Tsc22d3*), NMDA receptor-regulated gene 1 (*Narg1*), Myelocytomatosis oncogene (*Myc*), Iroquois related homeobox 5 (Drosophila) (*Irx5*), Forkhead box B2 (*Foxb2*), Alpha thalassemia/mental retardation syndrome X-linked homolog (human) (*Atrx*) were up regulated over 2.0-fold. Increases in gene expression and activities of *Pparg*, *Cebpa* and *Myc* have been known to induce many lipogenic genes and their regulators playing pivotal roles in adipogenesis (Heath *et al.*, 2000; Pulverer *et al.*, 2000). Activating transcription factor 4 and 5 (*Atf4*, *Atf5*), Interferon regulatory factor 8 (*Irf8*), Nuclear factor I/B (*Nfib*), Twist gene

homolog 1 (Drosophila) (*Twist1*), Early growth response 1 (*Egr1*) and Four and a half LIM domains 2 (*Fhl2*) were down-regulated over 2-fold. Activating transcription factor 2 was previously shown to be decreased during adipogenesis (Burton *et al.*, 2004). The roles of transcription factors previously unappreciated in the generation of differentiated adipocytes remain to be determined.

Many genes involved in signal transduction exhibit dynamic changes in expression. One of the most interesting results was that genes in the transforming growth factor, beta (*TGF β*) signal pathway were expressed an average 2.5-fold higher in the 3T3-L1 cells at the early differentiation. Transforming growth factor, beta 3 (*Tgfb3*), and transforming growth factor beta receptor II and III (*Tgfb2*, *Tgfb3*) were increased by 2.5-, 1.2 and 1.2-fold during adipocyte differentiation. Regulator of G-protein signaling 2 (*Rgs2*) (up-regulated by 5.2-fold), Metallothionein 1 and 2 (*Mt1* and *Mt2*) (up-regulated by 1.3- and 2.0-fold), Insulin-like growth factor binding protein 4 (*Igfbp4*) (up-regulated by 2.5-fold) showed similar expression patterns. *Rgs2* was previously isolated as a gene which is induced at the earliest stage of adipocyte differentiation (Imagawa *et al.*, 1999; Nishizuka *et al.*, 2001). Induction of *Rgs2* was stage-specific, occurring in the confluent cells but not in the proliferating cells, indicating that the expression of *Rgs2* is closely related to adipocyte differentiation. MTs are also induced in 3T3-L1 mouse fibroblasts during adipocyte differentiation and suggested to play a role in adipocyte differentiation (Schmidt & Beyersmann, 1999; Traythurn *et al.*, 2000a; Traythurn *et al.*, 2000b). The level of MT-2A mRNA in subcutaneous fat tissues was found to be significantly higher in obese subjects (Do *et al.*, 2000). Our results were consistent with these previous reports.

Genes coding for transporter proteins were also affected. Fatty acid binding protein 4 (*Fabp4*), adipocyte increased (18-fold) during adipogenesis, as previously reported (Burton *et al.*, 2004). Annexin A5 was decreased by -2.1-fold, which is consistent with a previous report (Burton *et al.*, 2004).

Regulatory regions in promoters of differentially expressed genes provide insight into potential coordinate regulation of gene expression. PAINT analysis identified binding sites for transcription factors known to serve important roles in cell proliferation, differentiation and developmental processes in other systems; AP-1, HEN1, NF-kappaB family, Oct-1, c-Myc/Max, Tax/CREB, and v-Myb. Rapid and transient induction of AP-1 was shown during the conversion of 3T3-L1 cells to adipocytes (Ishida *et al.*, 1988). Up-regulation of c-Myc during adipose differentiation was observed in our microarray analysis. In addition, increases in gene expression and activities of *Myc* have been shown in previous studies (Heath *et al.*, 2000; Pulverer *et al.*, 2000). Adipogenesis is associated with changes in amount and subunit composition of the NF-kB complexes. NF-kB subunits p65 (*RelA*), p68 (*RelB*), and I κ B were shown to be up-regulated during fat cell differentiation. Correspondingly, basal NF-kB nuclear gel shift and luciferase reporter assays were also shown to be induced in parallel during differentiation (Berg

et al., 2004). Oct-1 was shown to be involved in TNF α -induced decreases in adipocytes differentiation and LPL gene transcription (Morin *et al.*, 1995).

It is noteworthy that TFBS for HEN1 not known to be expressed during adipocyte differentiation were found to be significantly enriched in this promoter analysis (see Table 3).

HEN1 was originally discovered because of its homology within the bHLH motif to the hematopoietic transcription factor and oncogene *SCL* (Begley *et al.*, 1992). HEN1 expression is confined to the nervous system, and has been observed to be induced throughout embryonic development (Begley *et al.*, 1992). The roles of HEN1 in the generation of differentiated adipocytes remain to be determined.

In summary, we investigated the transcriptional profiles in 3T3-L1 cells at early stage of differentiation and analyzed the promoters of differentially regulated genes using bioinformatics. This information will be useful for establishing the regulatory network among genes involved in adipogenesis. Further studies of promoter occupancy and transcription factor perturbation are now required to provide a functional validation of the TFBSs identified in this study, as well as the TFs that bind to them.

Literature cited

- Addya S, Keller MA, Delgrosso K, Ponte CM, Vadigepalli R, Gonye GE & Surrey S (2004). Erythroid-induced commitment of K562 cells results in clusters of differentially expressed genes enriched for specific transcription regulatory elements. *Physiol Genomics* 19:117-130.
- Ahn JI, Lee KH, Shin DM, Shim JW, Lee JS, Chang SY, Lee YS, Brownstein MJ, Lee SH & Lee YS (2004). Comprehensive transcriptome analysis of differentiation of embryonic stem cells into midbrain and hindbrain neurons. *Dev Biol* 265:491-501.
- Berg AH, Lin Y, Lisanti MP & Scherer PE (2004). Adipocyte differentiation induces dynamic changes in NF- κ B expression and activity. *Am J Physiol Endocrinol Metab* 287:E1178-1188.
- Begley CG, Lipkowitz S, Gobel V, Mahon KA, Bertness V, Green AR, Gough NM & Kirsch IR (1992). Molecular characterization of NSCL, a gene encoding a helix-loop-helix protein expressed in the developing nervous system. *Proc Natl Acad Sci U S A* 89:38-42.
- Bradford MM (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248-254.
- Burton GR, Guan Y, Nagarajan R & McGehee RE Jr (2002). Microarray analysis of gene expression during early adipocyte differentiation. *Gene* 293:21-31.
- Burton GR, Nagarajan R, Peterson CA & McGehee RE Jr (2004). Microarray analysis of differentiation-specific gene expression during 3T3-L1 adipogenesis. *Gene* 329:167-185.
- Clamp M, Andrews D, Barker D, Bevan P, Cameron G, Chen Y, Clark L, Cox T, Cuff J, Curwen V, Down T, Durbin R, Eyraas E, Gilbert J, Hammond M, Hubbard T, Kasprzyk A, Keefe D, Lehvaslaiho H, Iyer V, Melsopp C, Mongin E, Pettett R, Potter S, Rust A, Schmidt E, Searle S, Slater G, Smith J, Spooner W, Stabenau A, Stalker J, Stupka E, Ureta-Vidal A, Vastrik I & Birney E (2003). Ensembl 2002: accommodating comparative genomics. *Nucleic Acids Res* 31:38-42.
- Do MS, Nam SY, Hong SE, Kim KW, Duncan JS, Beattie JH & Trayhurn P (2002). Metallothionein gene expression in human adipose tissue from lean and obese subjects. *Horm Metab Res* 34:348-351.
- Gregoire FM, Smas CM & Sul HS (1998). Understanding adipocyte differentiation. *Physiol Rev* 78:783-809.
- Guo X & Liao K (2000). Analysis of gene expression profile during 3T3-L1 preadipocyte differentiation. *Gene* 251:45-53.
- Heath VJ, Gillespie DA & Crouch DH (2000). Inhibition of the terminal stages of adipocyte differentiation by cMyc. *Exp Cell Res* 254:91-98.
- Imagawa M, Tsuchiya T & Nishihara T (1999). Identification of inducible genes at the early stage of adipocyte differentiation of 3T3-L1 cells. *Biochem Biophys Res Commun* 254:299-305.
- Ishida Y, Taniguchi H & Baba S (1988). Rapid and transient induction of c-fos and c-myc during adipose conversion of 3T3-L1 cells: no relationship with 1 α ,25-dihydroxyvitamine D₃-suppressed adipogenesis. *Kobe J Med Sci* 34:263-270.
- Kim JH, Kim HY & Lee YS (2001). A novel method using edge detection for signal extraction from cDNA microarray image analysis. *Exp Mol Med* 33:83-88.
- Kim S, Sohn I, Ahn JI, Lee KH, Lee YS & Lee YS (2004). Hepatic gene expression profiles in a long-term high-fat diet-induced obesity mouse model. *Gene* 340:99-109.
- Morin CL, Schlaepfer IR & Eckel RH (1995). Tumor necrosis factor- α eliminates binding of NF- κ B and an octamer-binding protein to the lipoprotein lipase promoter in 3T3-L1 adipocytes. *J Clin Invest* 95:1684-1689.
- Nishizuka M, Honda K, Tsuchiya T, Nishihara T & Imagawa M (2001). RGS2 promotes adipocyte differentiation in the presence of ligand for peroxisome proliferator-activated receptor gamma. *J Biol Chem* 276:29625-29627.
- Ntambi JM & Young-Cheul K (2000). Adipocyte differentiation and gene expression. *J Nutr* 130:3122S-3126S.
- Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, Moorman MA, Simonetti DW, Craig S & Marshak DR (1999). Multilineage potential of adult human mesenchymal stem cells. *Science* 284:143-147.
- Pulverer B, Sommer A, McArthur GA, Eisenman RN & Luscher B (2000). Analysis of Myc/Max/Mad network members in adipogenesis: inhibition of the proliferative burst and differentiation by ectopically expressed Mad1. *J Cell Physiol* 183:399-410.
- Ramirez-Zacarias JL, Castro-Munozledo F & Kuri-Harcuch W (1992). Quantitation of adipose conversion and triglycerides by staining intracytoplasmic lipids with Oil Red O. *Histochemistry* 97:493-497.
- Ron D & Habener JF (1992). CHOP, a novel developmentally regulated nuclear protein that dimerizes with transcription factors C/EBP and LAP and functions as a dominant-negative inhibitor of gene transcription. *Genes Dev* 6:439-453.
- Rosen ED & Spiegelman BM (2000). Molecular regulation of adipogenesis. *Annu Rev Cell Dev Biol* 16:145-171.
- Rosen ED, Walkey CJ, Puigserver P & Spiegelman BM (2000). Transcriptional regulation of adipogenesis. *Genes Dev* 14:1293-1307.
- Schmidt C & Beyersmann D (1999). Transient peaks in zinc and metallothionein levels during differentiation of 3T3-L1 cells. *Arch Biochem Biophys* 364:91-98.
- Soukas A, Socci ND, Saatkamp BD, Novelli S & Friedman JM

- (2001). Distinct transcriptional profiles of adipogenesis in vivo and in vitro. *J Biol Chem* 276:34167-34174.
- Traythurn P, Duncan JS, Wood AM & Beattie JH (2000a). Regulation of metallothionein gene expression and secretion in rat adipocytes differentiated from preadipocytes in primary culture. *Horm Metab Res* 32:542-547.
- Traythurn P, Duncan JS, Wood AM & Beattie JH (2000b). Metallothionein gene expression and secretion in white adipose tissue. *Am J Physiol Regul Integr Comp Physiol* 279:R2329-2335.
- Tusher VG, Tibshirani R & Chu G (2001). Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci U S A* 98:5116-5121.
- Yoon JC, Puigserver P, Chen G, Donovan J, Wu Z, Rhee J, Adelmant G, Stafford J, Kahn CR, Granner DK, Newgard CB & Spiegelman BM (2001). Control of hepatic gluconeogenesis through the transcriptional coactivator PGC-1. *Nature* 413:131-138.