

ORIGINAL PAPER

Alternative Medicine

Effectiveness of ozone therapy in addition to conventional treatment on mortality in patients with COVID-19

Şahin Çolak¹  | Burcu Genç Yavuz¹  | Mürsel Yavuz² | Burak Özçelik¹ |
Metin Öner¹  | Asu Özgültekin³  | Seniha Şenbayrak⁴ 

¹Haydarpaşa Numune Training and Research Hospital, Department of Emergency Medicine, University of Health Sciences, Istanbul, Turkey

²Kosuyolu Rezonans Clinic, Integrative Medicine, Istanbul, Turkey

³Haydarpaşa Numune Training and Research Hospital, Department of Anesthesiology and Intensive Care, University of Health Sciences, Istanbul, Turkey

⁴Haydarpaşa Numune Training and Research Hospital, Department of Infectious Disease and Clinical Microbiology, University of Health Sciences, Istanbul, Turkey

Correspondence

Şahin Çolak, Haydarpaşa Numune Training and Research Hospital, Department of Emergency Medicine, University of Health Sciences, Istanbul, Turkey.
Email: dr.sahincolak@gmail.com

Abstract

Aim: In this study, we aimed to investigate the effectiveness of ozone therapy, which is one of the integrative medicine applications that has been used safely for many years, on the prevalence of mortality in patients receiving COVID-19 treatment.

Methods: This was a prospective, controlled study conducted on patients with COVID-19 who were hospitalised. In this study, 55 patients were included. The patients were divided into two groups as the ozone and control group. Ozone therapy (major autohemotherapy) was applied to 37 patients who were being treated with the appropriate COVID-19 treatment protocol determined by the infectious diseases committee of our hospital. The ozone treatment protocol consisted of seven sessions (one session per day) of intravenous ozone administration, applied in a volume of 100 mL and a concentration of 30 µg/mL. Only the conventional COVID-19 treatment protocol was applied to 18 patients in the control group. Clinical follow-up was performed until the discharge of the patients from the hospital with successful treatment or until the mortality occurred. Factors affecting mortality were analysed using univariate regression analysis.

Results: Intensive care unit (ICU) hospitalisation was required in 6 of the 37 patients who were treated with ozone (16.2%), while 4 of 18 patients in the control group required ICU treatment (22.2%) ($P = .713$). When the mortality rates between the two groups were compared, mortality was lower in the ozone group ($P = .032$). As a result of univariate logistic regression analysis performed to investigate the factors affecting mortality, treatment with ozone therapy was determined as a risk factor for mortality. Patients receiving ozone therapy appear to have a lower mortality risk (odds ratio [OR]: 0.149, 95% confidence interval [CI] 0.026–0.863, $P = .034$).

Conclusion: In this study, the findings suggested that the administration of ozone therapy along with the conventional medical treatment in patients hospitalised for COVID-19 could reduce mortality.

1 | INTRODUCTION

The COVID-19 outbreak that began at the end of 2019 continues to affect the whole world. Although there was a decrease in the severity of the pandemic in the summer of 2020, the increase in

the number of cases with the arrival of autumn made the whole world uneasy again. SARS-CoV-2 is transmitted from person to person via droplets or direct contact, and the most common symptoms presented during the prodromal phase are fever, dry cough, myalgia and fatigue.^{1,2} Although the cases can be asymptomatic or

have mild symptoms, it has also been reported that approximately 20% of the hospitalised patients who have a more severe clinical presentation further develop acute respiratory distress syndrome (ARDS).^{3,4} The leading causes of death due to the virus are respiratory failure, hyperinflammation, cytokine storm or multiorgan failure.⁵

There is no specific antiviral drug approved by the US Food and Drug Administration (FDA) or European Medicines Agency (EMA) in the treatment of COVID-19 as of 5 February 2021. WHO also shares the opinion that a specific treatment still does not exist.⁶ Because an effective and specific treatment for SARS-CoV-2 has not been developed yet, interest in supportive treatments with proven safety, such as vitamin D supplementation, has increased to prevent mortality and morbidity in the management of the disease.⁴

Ozone therapy has been known for more than 150 years.^{7,8} Its effectiveness, particularly in the treatment of infectious diseases, has been demonstrated in many studies conducted in Cuba, Italy, Germany, Russia and Spain.⁵ The beneficial effects of ozone have been demonstrated in various studies.⁵ These strong, low cost and non-pharmacological effects have also enabled ozone to be widely used for more than 50 pathological diseases, such as degenerative disorders, neurological, orthopaedic and genitourinary disorders.⁸⁻¹⁰ Ozone has been reported to be helpful in the treatment of these pathological disorders by inducing oxidative-antioxidative mechanisms.^{8,11} Besides, ozone provides oxygen substantially to tissues with poor oxygenation.^{6,12}

In this study, we investigated the effectiveness of ozone therapy to reduce mortality rates in patients hospitalised due to COVID 19.

2 | MATERIALS AND METHODS

2.1 | Study population and data collection

This study was approved by the Ethical Committee of the University of Health Sciences Haydarpasa Numune Training and Research Hospital and conducted following the principles of the Declaration of Helsinki. Written informed consent was obtained from all patients or their legal representatives. We performed a prospective quasi-experimental before-and-after pilot study. This study included mild and severe COVID-19 patients hospitalised in the Haydarpasa Numune Training and Research Hospital, with lung involvement and RT-PCR (reverse transcriptase-polymerase chain reaction) positivity for SARS-CoV-2.

The required sample size of this study was calculated using the Gpower 3.1.¹³ Drawing on the literature,⁴ a sample size of 51 patients was required to provide 80% power with 5% alpha and effect size $w = 0.394$. The participants were randomly assigned in a 2:1 allocation to the control ($n:18$) and treatment ($n:37$) groups using a computer-generated randomisation.

Thirty-seven patients who met the following criteria were included in the ozone group of our study. Inclusion criteria were as follows: application to the emergency department with fever and

What's known

- Ozone therapy is one of the most widely used integrative medicine applications.
- Although effective and specific treatment for SARS-CoV-2 has not yet been developed, reliable supportive therapies are drawing attention in the prevention of this disease.
- Ozone therapy was found to be effective to reduce mortality due to Covid-19.

respiratory system complaints, being 18 years or older, lung tomography findings indicating COVID-19 in accordance with the literature,¹⁴⁻¹⁶ positivity for SARS-CoV-2 nucleic acid (RT-PCR) test, acceptance of ozone therapy (by the patient or his/her legal guardian) by written consent. Patients who were breastfeeding, pregnant or patients with a diagnosis of glucose 6-phosphate dehydrogenase (G-6PD) deficiency were excluded from this study.

For the control group, 18 patients were included who met the above-mentioned inclusion criteria but did not consent to the ozone treatment protocol and accepted to participate in this study in the control group by giving written consent. Leucocyte and lymphocyte count, ferritin, D-Dimer, procalcitonin, C-reactive protein and IL-6 measurement tests were performed at the time of admission among patients with findings consistent with COVID-19 in lung tomography, and then patients were hospitalised.

2.2 | Procedures

Patients in the ozone and control groups received the appropriate medical treatment according to the COVID-19 protocol determined by the infectious diseases committee of our hospital and according to their individual clinical status. The main drugs in this treatment protocol consisted of hydroxychloroquine (400 mg every 12 hours on the first day and 200 mg every 12 hours for the next 4 days), enoxaparin, favipiravir and antibiotics if a secondary bacterial infection is considered and antipyretics if required. Other symptomatic treatment measures were also taken according to the patient's clinical picture. Ozone major autohemotherapy (MAH) was applied to the ozone patient group, along with the conventional medical treatment that was deemed appropriate. Ozone was produced by the Turkozone Blue S CE medical device. The ozone bottle and set were disposable, made of medical-grade materials, and fully ozone compatible (Medipac Medical®, Germany and Bexen Medical®, Spain).

MAH was administered to the patients once daily for seven consecutive days. Each time, 100 ml of venous blood was collected and mixed with O₃ gas. In our study in accordance with the literature, the mixture composed of oxygen (95%-100%) and ozone (1%-5%) with a 0.8 lit/min flow rate, and the final pressure of the gas remained at the normal atmospheric pressure.¹⁷ In this study, whole blood samples

were exposed to ozone gas at 30 µg/mL of ozone with five minute effective mixing (that is the best time for the homogeneous balance of ozone gas and blood¹⁷) at a 1:1 ratio of oxygen-ozone to blood volume. The sodium citrate ratio contained in the ozone bottles was 3.13%, in line with the recommendation of the world ozone federation.¹⁸ In our study, to provide standard treatment, 100 mL of blood was taken from each patient in special ozone bottles and ozonized, and reinfusion was performed in 10-15 minutes in accordance with the World Federation of Ozone recommendation.¹⁸

Patients were followed up until they were discharged from the hospital or mortality occurred. In our study, the pre-treatment biochemical test results of the patients were compared, and the mortality rate observed in the groups was calculated. The discharge and mortality rates of the patients in the control group and the patients in the ozone group were compared.

2.3 | Statistical analysis

Statistical analysis was performed using IBM SPSS Statistics for Windows, Version 23.0 (IBM Corp., Armonk, NY). The normality assessment was performed using the Shapiro-Wilk test. Descriptive analyses were presented using mean ± SD (range), median (range) or n (%), where appropriate. Categorical data were analysed using the Pearson chi-square test and Fisher's exact test. Mann-Whitney *U* test and Student's *t* test were utilised for analysis of non-normally and normally distributed numerical data, respectively. Wilcoxon signed ranks test was used to compare the measured parameters of patients before and after the treatment. Univariate and multivariate logistic regression analysis was used to determine independent risk factors associated with mortality. The variables with $P < .1$ in the univariate analyses were further tested in the multivariate models. Odds ratio (OR) with corresponding 95% CIs was reported. A *p* value of less than 0.05 was considered statistically significant.

3 | RESULTS

In this study, 55 patients diagnosed with COVID-19 pneumonia and hospitalised were included. The mean age of the patients was 60.2 ± 14.8 (min: 25, max: 88) and 52.7% ($n = 29$) of the participants were male. The mean age of the patients in the group in which ozone therapy was not applied ($n = 18$) was 64.7 ± 10.4 , while the mean age of the patients in the ozone group ($n = 37$) was 58.03 ± 16.3 . While 44.4% of the patients in the group that did not receive ozone treatment were females, 48.6% of the patients in the ozone group were women. Mean age ($P = .118$) and gender distribution ($P = .769$) of the patients according to the patient groups were similar (Table 1).

When the distribution of patients in both groups for comorbidity was compared, there was no significant difference concerning diabetes mellitus (DM) ($P = .713$), hypertension ($P = .925$), congestive heart failure (CHF) ($P = .999$), coronary artery disease (CAD) ($P = .346$) and neoplasms ($P = .590$). However, chronic renal failure

(CRF) ($P = .043$) was observed with a higher rate in patients receiving ozone therapy (Table 1). Although the rate of chronic obstructive pulmonary disease (COPD) was observed to be higher in patients who received ozone therapy, this difference was not statistically significant ($P = .078$). In this ongoing pandemic, especially patients over the age of 50 and infected with COVID-19 with comorbidities were hospitalised in our hospital in line with the recommendation of the science committee of the Turkish Ministry of Health. For this reason, the demographic characteristics of patients who received and did not receive ozone therapy were similar in our study. Interestingly, patients with CRF were more likely to accept ozone therapy, which might be because they thought that the course of COVID-19 would be worse in CRF disease and relied more on ozone therapy as an alternative therapy. Therefore, randomisation could not provide a homogeneous distribution of all variables examined between the two groups. When the vital signs of the patients during the admission to the hospital were evaluated, no significant difference was found between the groups concerning body temperature ($P = .619$), heart rate ($P = .109$), systolic blood pressure ($P = .663$) and saturation of O₂ ($P = .068$). There was no difference in pre-treatment levels of IL-6 ($P = .993$), D-Dimer ($P = .167$), ferritin ($P = .893$) and procalcitonin ($P = .352$) according to the study groups. The rate of hospitalisation in the intensive care unit (ICU) was similar according to the study groups ($P = .713$) (Table 1).

All hospitalised patients received the best available therapy with the same standard care of the Turkish Ministry of Health protocol. All patients received hydroxychloroquine and enoxaparin treatment in both groups. In addition to hydroxychloroquine treatment, Favipiravir was added to the treatment of 7.3% of the patients, and Ritonavir was added to 3.6% of the patients. No significant difference was determined between patients who received favipiravir and ritonavir treatment regarding ozone and the control group ($P = .590$, $P = .999$, respectively). Intravenous antibiotic treatment was administered to 94.8% of the total patients. Non-invasive treatment was applied to 10.9% of the patients and there was no significant difference between the groups ($P = .999$). Mechanical ventilation was applied to 9.1% of the patients in total. No significant difference was determined between patients who received mechanical ventilation regarding ozone and the control group ($P = .999$).

In our study, the findings showed that the mortality rate in the ozone group ($n = 37$) was significantly lower than the control group ($n = 18$) ($P = .032$). When all participants were evaluated, the mortality rate was 50% ($n = 5$) in patients hospitalised in the ICU and 4.4% ($n = 2$) in patients hospitalised in the regular ward. The mortality rate was higher in patients hospitalised in the ICU ($n = 10$) ($P = .001$). In the group of patients who did not receive ozone treatment ($n = 18$), the mortality rate of the patients hospitalised in the ICU (75%, $n = 3$) was higher than the patients followed up in the regular ward (14.3%, $n = 2$) ($P = .044$). Similarly, in the group that received ozone therapy ($n = 37$), the mortality rate in patients requiring intensive care (33.3%, $n = 2$) was higher than in patients followed up in the ward (0%) ($P = .023$). When the mortality rates of patients who were

TABLE 1 Comparison of the control group and ozone group, regarding age, gender, comorbidities, temperature, pulse, systolic blood pressure, saturation O₂ and laboratory parameters based on the in-hospital mortality

	Control group (n:18)	Ozone group (n:37)	P
Age			
min-max	42.0–83.0	25.0–88.0	.118
mean ± sd	64.7 ± 10.4	58.03 ± 16.3	
Gender			
Male	10 (55.6)	19 (51.4)	.769
Female	8 (44.4)	18 (48.6)	
Co-morbidities			
Diabetes	4 (22.2)	6 (16.2)	.713
Hypertension	9 (50)	19 (51.4)	.925
Coronary artery disease	3 (16.7)	11 (29.7)	.346
COPD	1 (5.6)	11 (29.7)	.078
Congestive heart failure	1 (5.6)	4 (10.8)	.999
Neoplastic disease	2 (11.1)	2 (5.4)	.590
Chronic renal failure	0 (0)	8 (21.6)	.043
Temperature °C (med, min-max)	36.65 (36-38)	36.5 (36-38.3)	.619
Pulse bpm (med, min-max)	90.78 ± 9.97 (74-108)	86.46 ± 8.86 (72-110)	.109
SBP mmHg (med, min-max)	129.5 ± 22.54 (93-170)	132.35 ± 22.65 (90-181)	.663
Saturation O ₂ (med, min-max)	95 (80-99)	93 (82-99)	.068
IL-6 pg/ml (med, min-max)	19.05 (3.5-69)	17.5 (2.84-87.5)	.993
D-Dimer ng/ml (med, min-max)	785 (240-10,190)	1165 (417-7504)	.167
Ferritin ng/ml (med, min-max)	234.5 (95-1165)	334 (6-2907)	.893
Procalcitonin ng/ml (med, min-max)	0.05 (0.05-16)	0.05 (0.05-3.39)	.352
Intensive care unit	4 (22.2)	6 (16.2)	.713
Mortality	5 (27.8)	2 (5.4)	.032

Note: Data are presented as mean ± SD (range), median (range) or n (%). Student's t test, Mann-Whitney U test, Pearson chi-square test and Fisher's exact test.

Abbreviations: COPD, chronic obstructive pulmonary disease; LC, laboratory characteristics; SBP, systolic blood pressure.

hospitalised in ICU (n = 10) were compared concerning the treatment groups, it was observed that the mortality rate of patients who received ozone therapy (33.3%) was lower than patients who did not (75%), while this difference was not statistically significant ($P = .524$). We think that this difference is not statistically significant due to the small number of patients in the study groups. All our patients had pneumonia. Complications we observed in intensive care patients; ARDS (n = 4), multiorgan failure (n = 4), gastrointestinal bleeding (n = 1) and acute cardiac injury (n = 1). Among the patients treated in the ward (n = 45), all the 31 patients who received ozone therapy were discharged after successful treatment, while mortality occurred in 6.1% (n = 2) of 14 patients who did not receive ozone therapy. Although there was no significant difference between the death rates of the patients hospitalised in the regular ward concerning the treatment groups ($P = .092$), there was no death in the ozone group.

Ozone therapy was found effective in the univariate regression analysis performed to determine the factors affecting mortality (OR: 0.149; %95 CI: 0.026-0.863; $P = .034$) (Table 2).

4 | DISCUSSION

In our study, we applied ozone therapy to patients infected with SARS-Cov-2 in addition to conventional treatment and investigated the clinical outcomes of the patients compared to the group that did not receive ozone therapy. When we compared the results in the two groups that were similar regarding age, gender and comorbid diseases, we found that the mortality rate was significantly lower in the ozone treatment group. In the univariate regression analysis of factors affecting mortality, the findings suggest the effectiveness of ozone therapy in COVID-19 treatment.

The specific treatment of COVID-19 has not been developed yet, but the fight against coronavirus with antiviral drugs and symptomatic treatments goes on worldwide. In vaccine studies, no one has yet achieved a definite success so far. The development of alternative treatments to reduce the mortality of COVID-19 continues. In addition, ozone therapy, known for its high oxidant properties, is a method of treatment that has been used safely in many countries in infectious, immunological and

TABLE 2 Univariate regression analysis of factors affecting mortality

Variables	Univariate model	
	OR (95% CI)	P
Age	1.059 (0.990-1.133)	.098
Male	0.635 (0.128-3.146)	.578
Diabetes	4.393 (0.804-23.999)	.088
Hypertension	1.333 (0.269-6.606)	.725
CAD	1.2 (0.205-7.011)	.840
COPD	1.52 (0.256-9.028)	.645
CHF	1.833 (0.175-19.252)	.613
CRF	2.8 (0.440-17.799)	.275
Temperature	0.42 (0.070-2.532)	.344
Pulse	1.046 (0.963-1.136)	.286
SBP	0.985 (0.949-1.023)	.438
Saturation O ₂	1.017 (0.849-1.219)	.851
Ozone therapy	0.149 (0.026-0.863)	.034

Abbreviations: CAD, coronary artery disease; CHF, congestive heart failure; CI, confidence interval; COPD, chronic obstructive pulmonary disease; CRF, chronic renal failure; OR, odds ratio; SBP, systolic blood pressure.

vascular diseases for many years.¹⁰ SARS-CoV-2 is an enveloped virus and the high density of double-bonded molecular bodies in the structure of it, which facilitates such oxidant agents to damage the integrity of the virus.^{6,19} Similar to the Ebola virus, the spike and envelope proteins of SARS-CoV-2 are rich in cysteine and tryptophan amino acids, which make them vulnerable to oxidation.^{20,21} Ozone therapy thus causes oxidation in the cysteine and tryptophan residues of the viral membrane proteins.^{19,22} Apart from the strong oxidant effect, it has also been reported that lymphocytes and monocytes re-infused into the patient during MAH would stimulate the immune system.⁶ Thus, viral replication and the progression of infection can be prevented.^{23,24} The Menendez Cuban group reported in their animal studies that the previously applied ozone therapy to the endotoxic shock model was as effective as dexamethasone treatment in reducing tumor necrosis factor α levels.^{25,26} This information suggests that ozone therapy is quite valuable in preventing a cytokine storm, one of the leading causes of death in patients infected with COVID-19.²⁶⁻²⁹ In our study, patients with RT-PCR positiveness for SARS-CoV-2 were categorised as mild and severe COVID-19 infection according to the literature.^{30,31} Mild type characterised with mild pneumonia cases. Patients with dyspnea, respiratory rate ≥ 30 breaths per minute, blood oxygen saturation $\leq 93\%$ and partial pressure of oxygen (PaO₂)/fraction of inspired oxygen (FiO₂) ratio < 300 were included in the severe type. Among 55 patients infected with COVID-19, 10 required ICU admission with a diagnosis of severe type COVID-19. The proportion of patients hospitalised in ICU between the ozone and control groups was similar (Table 1). We found that the mortality rate in ward and

intensive care patients in the group receiving ozone therapy was significantly lower than the control group who did not receive ozone therapy ($P = .032$).

Many studies have reported that advanced age and comorbid diseases may negatively affect the prognosis of COVID-19 and increase mortality.³² Diabetes, hypertension, cardiovascular disease, chronic respiratory disease, cancer and cerebrovascular disease are mainly known risk factors in this respect.^{32,33} It has also been reported that acute kidney injury (AKI) may develop during the course of patients infected with COVID-19, and this situation may significantly increase the risk of mortality.^{34,35} The development of AKI also may lead to a poor prognosis in patients who are infected with COVID-19 with chronic kidney failure (CRF) or with a history of renal transplant.³⁴ In a study conducted on 101 cases that died due to COVID-19, it was reported that 11% of the patients had CRF and 23% developed AKI.³⁶ In our study, we could not find any significant difference between the groups concerning DM, hypertension, CAD, COPD, CHF or neoplasm rates. At the same time, although 21.6% of patients in the ozone group had a history of CRF ($P = .043$), a lower mortality rate was observed compared to the other group. We think this finding supports the effectiveness of ozone therapy.

Another prognostic factor in patients with COVID-19 is the elevation of D dimer, ferritin and Interleukin-6 levels. These parameters were associated with poor prognosis in many studies.^{37,38} In our study, when the laboratory tests performed before treatment between the ozone group and the control group were compared, IL-6, D-Dimer, Ferritin and procalcitonin levels and vital parameters (fever, pulse, systolic blood pressure [SBP] and saturation) were similar.

In COVID-19, for which a specific treatment is not yet available, clinical management sometimes challenges both physicians and patients. Not every patient gives the same response to every drug. At the same time, the toxic side effects of the drugs used may negatively affect the course of the disease. Ozone therapy, on the other hand, is an inexpensive, reliable and well-known method of treatment for many years. In our study, no side effects that could be associated with ozone treatment were observed in the group which received it. We consider that mortality rates can be further decreased with ozone therapy to be applied in addition to the existing conventional treatment modalities.

The limitations of our study are that this study was conducted in a single centre, and the number of patients was small. Another limitation of our study is that the body mass index (BMI) of the patients could not be calculated. The reason for this is that there are no special scales for every patient in our hospital and the use of common scales is not possible under pandemic conditions. Because each patient does not know his/her own weight and height values, BMI data of the patients could not be included in our study. For this reason, there was no homogeneity between the groups in terms of BMI. Multi-centre studies on a larger patient population, including patients' BMI, will further provide valuable insights into the understanding of the effectiveness and significance of ozone therapy.

5 | CONCLUSION

In conclusion, in this study, we demonstrated that applying ozone therapy to patients hospitalised for COVID-19 could contribute to clinical outcomes. No side effects related to ozone therapy were observed in our study. At the same time, the positive effects of ozone on the control of oxidative stress and immunomodulation have been supported by decreasing mortality rates and univariate regression analysis.

ACKNOWLEDGEMENT

The authors did not apply for a specific grant for this research from any funding agency in the public, commercial or non-profit sectors.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ORCID

Şahin Çolak  <https://orcid.org/0000-0001-8192-9652>

Burcu Genç Yavuz  <https://orcid.org/0000-0001-6693-5288>

Metin Öner  <https://orcid.org/0000-0002-2865-9822>

Asu Özgültekin  <https://orcid.org/0000-0003-4307-9604>

Seniha Şenbayrak  <https://orcid.org/0000-0002-4983-6613>

REFERENCES

- Habibzadeh P, Stoneman EK. The novel coronavirus: a bird's eye view. *Int J Occup Environ Med*. 2020;11:65-71.
- Chia PY, Coleman KK, Tan YK, et al. Detection of air and surface contamination by SARS-CoV-2 in hospital rooms of infected patients. *Nat Commun*. 2020;11:2800.
- Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395:507-513.
- Entrenas Castillo M, Entrenas Costa LM, Vaquero Barrios JM, et al. Effect of calcifediol treatment and best available therapy versus best available therapy on intensive care unit admission and mortality among patients hospitalized for COVID-19: A pilot randomized clinical study. *J Steroid Biochem Mol Biol*. 2020;203:105751.
- Fernández-Cuadros ME, Albaladejo-Florín MJ, Álava-Rabasa S, et al. Effect of rectal ozone (O₃) in severe COVID-19 pneumonia: preliminary results. *SN Compr Clin Med*. 2020;1:9.
- Martínez-Sánchez G, Schwartz A, Donna VD. Potential cytoprotective activity of ozone therapy in SARS-CoV-2/COVID-19. *Antioxidants (Basel)*. 2020;9.
- Elvis AM, Ekta JS. Ozone therapy: a clinical review. *J Nat Sci Biol Med*. 2011;2:66-70.
- Scassellati C, Galoforo AC, Bonvicini C, Esposito C, Ricevuti G. Ozone: a natural bioactive molecule with antioxidant property as potential new strategy in aging and in neurodegenerative disorders. *Ageing Res Rev*. 2020;63:101138.
- Ameli J, Banki A, Khorvash F, Simonetti V, Jafari NJ, Izadi M. Mechanisms of pathophysiology of blood vessels in patients with multiple sclerosis treated with ozone therapy: a systematic review. *Acta Biomed*. 2019;90:213-217.
- Smith NL, Wilson AL, Gandhi J, Vatsia S, Khan SA. Ozone therapy: an overview of pharmacodynamics, current research, and clinical utility. *Med Gas Res*. 2017;7:212-219.
- Fernández-Cuadros ME, Albaladejo-Florín MJ, Peña-Lora D, Álava-Rabasa S, Pérez-Moro OS. Ozone (O₃) and SARS-CoV-2: physiological bases and their therapeutic possibilities according to COVID-19 evolutionary stage. *SN Compr Clin Med*. 2020;1:9.
- Clavo B, Pérez JL, López L, et al. Effect of ozone therapy on muscle oxygenation. *J Altern Complement Med*. 2003;9:251-256.
- Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007;39:175-191.
- Ai T, Yang Z, Hou H, et al. Correlation of chest CT and RT-PCR testing for coronavirus disease 2019 (COVID-19) in China: a report of 1014 cases. *Radiology*. 2020;296:E32-E40.
- Xie X, Zhong Z, Zhao W, Zheng C, Wang F, Liu J. Chest CT for typical coronavirus disease 2019 (COVID-19) pneumonia: relationship to negative RT-PCR testing. *Radiology*. 2020;296:E41-E45.
- Zhai P, Ding Y, Wu X, Long J, Zhong Y, Li Y. The epidemiology, diagnosis and treatment of COVID-19. *Int J Antimicrob Agents*. 2020;55:105955.
- Mehraban F, Seyedarabi A, Ahmadian S, Mirzaaghaei V, Moosavi-Movahedi AA. Personalizing the safe, appropriate and effective concentration(s) of ozone for a non-diabetic individual and four type II diabetic patients in autohemotherapy through blood hemoglobin analysis. *J Transl Med*. 2019;17:227.
- Baeza J, Cabo JR, Gomez M, Menendez S, Re L. WFOTs review on evidence based ozone therapy. In *World Federation of Ozone Therapy*, 2015: 116.
- Tizaoui C. Ozone: a potential oxidant for COVID-19 virus (SARS-CoV-2). *Ozone Sci Eng*. 2020;42:378-385.
- Lopez LA, Riffle AJ, Pike SL, Gardner D, Hogue BG. Importance of conserved cysteine residues in the coronavirus envelope protein. *J Virol*. 2008;82:3000-3010.
- Schoeman D, Fielding BC. Coronavirus envelope protein: current knowledge. *Virus J*. 2019;16:69.
- Broer R, Boson B, Spaan W, Cosset FL, Corver J. Important role for the transmembrane domain of severe acute respiratory syndrome coronavirus spike protein during entry. *J Virol*. 2006;80:1302-1310.
- Re L, Martínez-Sánchez G, Bordicchia M, et al. Is ozone preconditioning effect linked to Nrf2/EpRE activation pathway in vivo? A preliminary result. *Eur J Pharmacol*. 2014;742:158-162.
- Rossmann A, Mandic R, Heinis J, et al. Intraperitoneal oxidative stress in rabbits with papillomavirus-associated head and neck cancer induces tumoricidal immune response that is adoptively transferable. *Clin Cancer Res*. 2014;20:4289-4301.
- Zamora ZB, Borrego A, López OY, et al. Effects of ozone oxidative preconditioning on TNF-alpha release and antioxidant-prooxidant intracellular balance in mice during endotoxic shock. *Mediators Inflamm*. 2005;1:16-22.
- Rowen RJ, Robins H. A plausible "penny" costing effective treatment for corona virus—ozone therapy. *J Infect Dis Epidemiol*. 2020;6:113.
- Liu Q, Zhou YH, Yang ZQ. The cytokine storm of severe influenza and development of immunomodulatory therapy. *Cell Mol Immunol*. 2016;13:3-10.
- Tetro JA. Is COVID-19 receiving ADE from other coronaviruses? *Microbes Infect*. 2020;22:72-73.
- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan. *China Lancet*. 2020;395:497-506.
- Lian J, Jin X, Hao S, et al. Analysis of epidemiological and clinical features in older patients with corona virus disease 2019 (COVID-19) out of Wuhan. *Clin Infect Dis*. 2020;71:740-747.
- Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the chinese center for disease control and prevention. *JAMA*. 2020;323:1239-1242.

32. Liu K, Chen Y, Lin R, Han K. Clinical features of COVID-19 in elderly patients: a comparison with young and middle-aged patients. *J Infect.* 2020;80:e14-e18.
33. Li JY, You Z, Wang Q, et al. The epidemic of 2019-novel-coronavirus (2019-nCoV) pneumonia and insights for emerging infectious diseases in the future. *Microbes Infect.* 2020;22:80-85.
34. Adapa S, Chenna A, Balla M, et al. COVID-19 pandemic causing acute kidney injury and impact on patients with chronic kidney disease and renal transplantation. *J Clin Med Res.* 2020;12:352-361.
35. Cheng Y, Luo R, Wang K, et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int.* 2020;97:829-838.
36. Shi Q, Zhao K, Yu J, et al. Clinical characteristics of 101 COVID-19 nonsurvivors in Wuhan, China: a retrospective study. In. *medRxiv.* 2020. 03.04.20031039.
37. Huang I, Pranata R, Lim MA, Oehadian A, Alisjahbana B. C-reactive protein, procalcitonin, D-dimer, and ferritin in severe coronavirus disease-2019: a meta-analysis. *Ther Adv Respir Dis.* 2020;14:1753466620937175.
38. Guo YR, Cao QD, Hong ZS, et al. The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak—an update on the status. *Mil Med Res.* 2020;7:11.

How to cite this article: Çolak Ş, Genç Yavuz B, Yavuz M, et al. Effectiveness of ozone therapy in addition to conventional treatment on mortality in patients with COVID-19. *Int J Clin Pract.* 2021;75:e14321. <https://doi.org/10.1111/ijcp.14321>