




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Revista de Psiquiatría y Salud Mental

www.elsevier.es/saludmental



SCIENTIFIC LETTER

COVID-19 impact vs. suicide impact in Spain

Comparación del impacto producido por el COVID-19 y por el suicidio en España

Coronavirus disease 2019 (COVID-19) has had a major impact on our lives, both socio-economically and in terms of our physical and mental health. On 30 January 2020, the World Health Organization (WHO) declared COVID-19 a public health emergency.

On 11 March 2020, the pandemic was officially declared. To date, 102,392 people have died worldwide by COVID-19, while many others have been affected directly or indirectly.¹

The pandemic has had a disruptive socio-economic effect: approximately one third of the world's population was locked down and severe restrictions on freedom of movement were imposed, leading to a drastic reduction in economic activity and a parallel rise in unemployment. At the healthcare level, there has been an unprecedented overflow and collapse of hospitals.²

It is hard to imagine a health problem with a greater impact than COVID-19, but the truth is that suicide is one of the most important public health problems and one of the main causes of death in our country.³ The real magnitude of this problem is underestimated and has been increasing in recent years. According to data provided by the National Statistics Institute (INE), 3941 people took their own lives in Spain in 2020, an increase of 7.4% over 2019. The epidemiological and social impact of suicide involves not only mortality due to completed suicide, but also suicidal ideation and suicide attempts, as well as the consequences for the quality of life of the population.⁴

Unlike COVID-19, suicide is a health problem affecting young people. Currently, suicide is the leading cause of unnatural death in our country in people aged 15–29 years.⁵ Of concern, suicide attempts in this age group increased by 250% in 2020 compared to previous years.³

The aim of this study is to quantify the impact of suicide in terms of years of potential life lost (YPLL) and to compare it with the YPLL by COVID-19 in the year 2020 in Spain. YPLL below the age of 75 is an illustrative indicator of the loss suffered by society because of premature deaths.⁶

For this study, we used data on deaths by age group caused by COVID-19 and suicide as provided by the Instituto

de Salud Carlos III [Carlos III Health Institute]⁷ and the Instituto Nacional de Estadística [National Institute of Statistics]³ for ages 0–75 years, from 1 January to 31 December 2020. To calculate the YPLL, we subtracted the average age of each age group to age 75. We multiplied the result by the number of deaths corresponding to each age group and then summed the results to obtain the total YPLL by each cause.

In 2020, YPLLs as a result of suicide were 80,679.5, versus 63,417.43 YPLLs due to COVID-19, a difference of 17,262.1 YPLLs. These results are illustrated in Fig. 1.

We observed that the death rates with the greatest weight in the case of COVID infection are located in the older age groups, the most prominent being between 60 and 69 years of age, age groups in which most deaths are often related to cardiovascular risk factors, pulmonary diseases or complications of other pathologies. However, the greatest number of deaths by suicide occur in the 25–54 age group.⁵ That is why YPLL are higher for suicide than for COVID-19 in our country.

Our results contrast with the study by Porras-Segovia et al. who compared YPLLs from COVID-19 in 2020 with those produced by suicide in 2019 in the United States, finding that COVID-19 had produced more YPLLs than suicide, albeit with a narrow difference.⁸


The scope and implications of our findings encompass multiple aspects. The greatest losses in work productivity are concentrated in young people, as well as a decrease in natural population growth through a reduction of birth rates, which implies a rationalization and restructuring of health spending. We must also remember the suffering that suicidal behavior entails for the person and his or her social environment.⁹

There are still great challenges in suicide prevention. It is necessary to address this problem through effective primary and secondary prevention, diagnosis of the precipitating or underlying pathology, as well as improvements in treatment, early diagnosis and allocation of funds.

Since March 2020, a major effort has been made to tackle the COVID-19 pandemic. In comparison, suicide remains an orphan disease. Evidence shows that, with proper healthcare planning, suicide can be prevented.¹⁰ A comprehensive approach is required for a better understanding of the predisposing and precipitating factors that contribute to suicide behavior, and a coordinated involvement of health professionals and healthcare planning is required to prevent suicide and increase the quality of life of the population.

<https://doi.org/10.1016/j.rpsm.2022.05.006>

1888-9891/© 2022 SEP y SEPB. Published by Elsevier España, S.L.U. All rights reserved.



* Data from the "Instituto Nacional de Estadística" [National Statistics Institute] and the "Instituto de Salud Carlos III" [Health Institute Carlos III] compran YPLL due to COVID-19 and suicide by age in 2020. Area under the curve for suicide: 80,679.5 YPLL. Area under the curve for COVID-19: 63,417.43 YPLL

Figure 1 Years of Potential Life Lost (YPLL) before age 75 due to COVID-19 and suicide*.

Acknowledgements

Our research team wants to thank the support received from "Instituto de Salud Carlos III" (CM19/00026), the "Ministerio de Ciencia, Innovación y Universidades" (RTI2018-099655-B-I00; TEC2017-92552-EXP) and the Alicia Koplowitz Foundation.

References

1. World Health Organization. Novel Coronavirus (2019-nCoV): Situation Report - 11. Data as reported by 31 January 2020.
2. Minué Lorenzo S. La pandemia COVID-19: Lo que hemos aprendido hasta ahora desde España. *Revista da Rede APS* 2020. 2020.
3. INE, 2022. Ine.es. Available from: https://www.ine.es/prensa/edcm_2020.pdf.
4. Sáiz PA, Bobes J. Prevención del suicidio en España: una necesidad clínica no resuelta. *Rev psiquiatr salud ment.* 2014;1-4. Giner Jiménez L. Diferencias en la conducta suicida: estudio comparativo entre los intentos de suicidio y suicidio consumado, 2010.
5. Sanidad.gob.e s. 2022. Mortalidad por suicidio en España, 2020. Available from: https://www.sanidad.gob.es/gl/estadEstudios/estadisticas/estadisticas/estMinisterio/mortalidad/docs/Defunciones_Suicidio2020.pdf.
6. Arriaga EE. Los años de vida perdidos: su utilización para medir los niveles y cambios de la mortalidad. *Notas de población.* 1996.
7. ISCIII, 2022. Isciii.es. Available from: <https://www.isciii.es/QueHacemos/Servicios/VigilanciaSaludPublicaRENAVE/EnfermedadesTransmisibles/Documents/INFORMES/Informes%20COVID-19/INFORMES%20COVID-19%202021/Informe%20n%C2%BA%20108%20Situaci%C3%B3n%20de%20COVID-19%20en%20Espa%C3%B1a%20a%209%20de%20diciembre%20de%202021.pdf>.

8. Porras-Segovia A, Baca-García E, Courtet P, Wainberg M, Oquendo MA. If suicide were CoViD-19: a neglected cause of premature death. *J Clin Psychiatry.* 2021;82:27939.
9. Zalsman G, Hawton K, Wasserman D, van Heeringen K, Arensman E, Sarchiapone M, et al. Suicide prevention strategies revisited: 10-year systematic review. *Lancet Psychiatry.* 2016;3:646-59.
10. Nordentoft M, Erlangsen A. Suicide-turning the tide. *Science.* 2019;365:725, <http://dx.doi.org/10.1126/science.aaz1568>.

Jimena María Merayo-Cano^a,
Alejandro Albán Porras-Segovia^{b,c},
Enrique Baca-García^{a,b,d,e,f,g,h,i,j,*}

^a Department of Psychiatry, University Hospital Fundación Jiménez Díaz, Madrid, Spain

^b Health Research Institute, Fundación Jiménez Díaz, Madrid, Spain

^c Child and Adolescent Mental Health Team, Imperial College London, London, United Kingdom

^d Department of Psychiatry, General Hospital of Villalba, Villalba, Madrid, Spain

^e Department of Psychiatry, Hospital Infanta Elena, Valdemoro, Madrid, Spain

^f Department of Psychiatry, Autonomous University of Madrid, Madrid, Spain

^g Department of Psychiatry, Hospital Rey Juan Carlos, Mostoles, Spain

^h CIBERSAM, Spain

ⁱ Department of Psychology, Universidad Católica del Maule, Talca, Chile

^j Department of Psychiatry, Centre Hospitalier Universitaire de Nîmes, Nîmes, France

* Corresponding author.

E-mail address: ebacgar2@yahoo.es (E. Baca-García).