

## RESEARCH ARTICLE

# One-year descriptive analysis of patients treated at an anti-rabies clinic—A retrospective study from Kashmir

Khalid Bashir<sup>1</sup>, Inaamul Haq<sup>2\*</sup>, S. Muhammad Salim Khan<sup>2</sup>, Mariya Amin Qurieshi<sup>2</sup>

**1** Department of Community Medicine, Government Medical College, Anantnag, Jammu & Kashmir, India,

**2** Department of Community Medicine, Government Medical College, Srinagar, Jammu & Kashmir, India

\* [haqinaam@yahoo.co.in](mailto:haqinaam@yahoo.co.in)



## Abstract

Dog bites in humans are a major public health problem in India in general and Kashmir in particular. Canine rabies is almost non-existent in developed countries and exists mainly in the poorer, low socioeconomic strata of society in the developing world. The objective of this study was to determine the characteristics, pattern, and burden of dog bite injuries in the Kashmir valley. Data from Anti-Rabies Clinic of a tertiary care hospital in Srinagar, the summer capital of the state of Jammu & Kashmir, was collated and analyzed. Analysis of records of all the patients who had reported between April 2016 and March 2017 was done. A total of 6172 patients had reported to the Anti-Rabies Clinic for management of animal bites from 1<sup>st</sup> April 2016 to 31<sup>st</sup> March 2017. Most of the patients were young males. Almost half (47.7%) of the patients were bitten in the afternoon. Lower limbs were the most common site of bite (71.7%). Most of the bites were of Category III (57.6%) followed by Category II (42.3%); only one case of Category I was recorded. Almost all (98.0%) cases reported being bitten by dogs. Conclusions: Category III dog bites on lower limbs were the most common type of animal bites presenting to the Anti-Rabies Clinic of a tertiary care hospital. Children have more chances of a bite on head and neck region. Serious and workable efforts have to be made to reduce the incidence and consequences of animal bites.

## OPEN ACCESS

**Citation:** Bashir K, Haq I, Khan SMS, Qurieshi MA (2020) One-year descriptive analysis of patients treated at an anti-rabies clinic—A retrospective study from Kashmir. *PLoS Negl Trop Dis* 14(8): e0007477. <https://doi.org/10.1371/journal.pntd.0007477>

**Editor:** Sergio Recuenco, Universidad Nacional Mayor de San Marcos, PERU

**Received:** May 16, 2019

**Accepted:** June 20, 2020

**Published:** August 25, 2020

**Copyright:** © 2020 Bashir et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its Supporting Information files.

**Funding:** This study has not been funded by any organization. All the cost of data entry & management and other costs has been borne by the authors.

**Competing interests:** The authors have declared that no competing interests exist.

## Author summary

In Kashmir, dog bite is an important public health problem. Thousands of people become victims of an animal bite, especially dog bite, and some of them develop rabies. Rabies is an invariably fatal viral disease resulting in approximately 59 000 human deaths per year globally, with 95% of cases occurring in Africa and Asia. The only way to prevent a rabies death is vaccination of an animal bite victim. In Kashmir, the burden and characteristics of dog bites are not routinely captured by the health system in place. We, therefore, attempted to find out the burden and characteristics of animal bite victims by analyzing one-year data from an Anti-Rabies Clinic at a tertiary care hospital in Kashmir. We found that 98% of the patients registered at the clinic during the period were victims of a dog bite. The victims were mostly young males but females and children were not shown any

mercy either. Lower limbs were the most frequently reported site of the bite. Our analysis of the data also revealed that children under 15 years were more prone to a bite in the head and neck region. We concluded that the burden of animal bites, especially dog bites, is high in Kashmir and recommended that serious efforts need to be put into practice to decrease the number of animal bite victims and prevent any rabies deaths.

## Introduction

Rabies transmission by dog bites is one of the major public health problems, considering its palpable fear and anxiety, as it is a sinister zoonotic infection transmitted to humans or animals by the bite of a rabid animal. Rabies virus, which belongs to the *Lyssavirus* genus of the *Rhabdoviridae* family, causes fatal encephalitis [1]. An estimated 59,000 people die of rabies annually globally, India accounting for the maximum of these deaths. The economic cost of rabies has been estimated to be 8.6 billion USD annually [2]. Dog bites constitute more than 99% of human rabies cases and the rest is associated with cats, foxes, bats, and other mammals [3]. Among all post-exposure prophylaxis (PEP) recipients, approximately 40% are children <15 years of age and they also have the highest rabies mortality [4]. Dog rabies is almost non-existent in developed countries of Europe, North America, and Australia; but it is prevalent in the developing world. Lack of reliable data and unawareness of the burden and risk factors associated with human rabies together represent a critical challenge for the formulation of policies and strategies to control the disease and has been considered a major cause for underinvestment in rabies control measures in these countries [2]. In developing countries, the real number of sufferers is probably higher than the reported statistics [2, 3]. In addition to the health importance in human-beings, disease outbreak among live-stock causes significant economic losses. Despite the preventability of rabies by effective and safe vaccines, the disease is still a healthcare problem in many countries, especially in Africa and Asia.[5]. Asia constitutes 96.5% of the economic burden of the disease in developing countries, which costs it 6.8 billion USD annually [2, 6].

Rabies is invariably fatal, yet completely preventable if PEP is applied in a timely and correct manner. The World Health Organization (WHO) has prepared standard recommendations for PEP, which include immediate and thorough wound washing with soap and water, followed by administration of the vaccine and additionally, infiltration of Rabies Immunoglobulin in WHO category III bites [3].

In Kashmir, the population of free-roaming dogs is high with a dog-human ratio of 1:14 which is higher in comparison to other parts in India [7]. The present study aims to generate a picture of the alarmingly increasing number of dog bite victims, potential cases of dreaded rabies, whose quality of physical and psychological health is directly or indirectly influenced by the event. Our objective was to determine the epidemiological features, characteristics, and pattern of dog bite victims who received PEP at a tertiary care center in Kashmir for one year. This will help in devising prevention strategies in view of the elimination of dog-mediated human rabies by 2030, as jointly outlined by WHO, FAO, OIE, the Global Alliance for Rabies Control, and the international community [8].

## Methods

This study was conducted at the Anti-Rabies Clinic (ARC) of Shri Maharaja Hari Singh (SMHS) Hospital, Government Medical College, Srinagar, run by the Department of Community Medicine. The ARC receives animal bite cases from the whole of Kashmir valley which

had a population of 6.9 million as per the 2011 census [9]. Animal bite cases are also managed at other hospitals across Kashmir. However, the ARC receives approximately three-fourth of all cases. The ARC maintains records of demographic and clinical details of patients visiting the clinic for treatment. The clinic adheres to WHO-recommended protocol for PEP, which includes prompt wound washing, an anti-rabies vaccine for WHO Category II and III exposures, and use of Immunoglobulin in Category III exposures [3]. Following recommendations of the World Health Organization on PEP following animal bites in 2005, the ARC started the use of the intradermal regimen in 2011. Recently the ARC has started the 2-site intradermal schedule as recommended by the recent WHO document [10].

The ARC follows a protocol for animal bite management which includes tetanus toxoid immunization for category II and III bites and Equine Rabies Immunoglobulin (ERIG) 40 IU/Kg body weight for category III bites. The ARC provides anti-rabies vaccines free of cost to the bite victims. ERIG is not provided free of cost. Patients need to purchase ERIG from the market. The health workers at the ARC are trained in the management of animal bites as well as the administration of the vaccine and ERIG.

We did an analysis of secondary data; records of the patients from 1<sup>st</sup> April 2016 to 31<sup>st</sup> March 2017 were collated and analyzed (S1 Data). All data analyzed were anonymized. Demographic and clinical details of the patients were entered into a Microsoft Excel spreadsheet. From records, we took into consideration different variables for study analysis including age, sex, time, site, severity and WHO category of bite, type of animal involved and the residence of the patient.

In order to estimate the incidence of animal bites in District Srinagar during the study period, we estimated the mid-interval population of the District based on 2011 census data [9] and the growth rates for urban areas of the state of Jammu & Kashmir reported by the Sample Registration System [11].

### Statistical analysis

Data was entered into a Microsoft Excel spreadsheet. Chi-square test was used to test independence between the site of bite versus age and sex. Adjusted standardized residuals (ASR) were used for cell-by-cell comparison of observed and expected frequencies and an ASR > 2 was considered significant [12]. In order to further explore the relationship between site of bite and sex, analysis restricted to cases >15 years of age was done. Chi-squared goodness-of-fit was used to test if the number of cases varied across months. A p-value of less than 0.05 was considered statistically significant. SPSS version 23.0 (IBM Corp. Released 2015. IBM SPSS Statistics for Windows, Version 23.0. Armonk, NY: IBM Corp.) was used for analyzing the data.

### Results

A total of 6172 patients had reported to the ARC from 1<sup>st</sup> April 2016 to 31<sup>st</sup> March 2017. The demographic details of the patients are shown in Table 1. The bite victims were mostly young males. In our study, 3773 out of a total of 6172 cases were from Srinagar. Animal bite cases are being managed at other hospitals in District Srinagar as well. During the study period, 2032 animal bite cases were managed at these 'other' hospitals. This gives us a total of 5805 (3773 + 2032) animal bite cases in District Srinagar during the study period. The estimated mid-interval population of District Srinagar during the study period was 1290644. Thus, the estimated incidence of an animal bite reported to a health facility in District Srinagar during the one-year study period was 450 per 100,000 population.


The number of cases was highest during the months of March to May (Fig 1).

**Table 1. Demographic characteristics of animal bite victims.**

Demographic characteristics	Number of bite victims (n = 6172)	Percentage (%)
<b>Age (years)</b>		
≤5	426	6.9
6–15	986	16.0
16–30	1519	24.6
31–50	2278	36.9
51–65	773	12.5
>65	190	3.1
<b>Sex</b>		
Female	1635	26.5
Male	4537	73.5
<b>Residence</b>		
District Srinagar	3773	61.1
Other districts	2399	38.9

<https://doi.org/10.1371/journal.pntd.0007477.t001>

Table 2 describes the characteristics of animal bites. Almost half of the cases (47.7%) were bitten in the afternoon (1200 hours to 1759 hours). Lower limbs were the most common site of bite (71.7%). Ninety-eight percent of patients were bitten by a dog (6048/6172). Only one patient reported with a WHO category I bite.



**Fig 1. Monthly distribution of animal bite cases.** Dash line indicates the monthly average of animal bite cases. The cases were the highest during the months following the winter months.

<https://doi.org/10.1371/journal.pntd.0007477.g001>

Table 2. Characteristics of animal bites.

Bite characteristics	Number of bite victims (n = 6172)	Percentage (%)
<b>Time of bite*</b>		
Morning (0600 to 0959 hrs)	920	14.9
Forenoon (1000 to 1159 hrs)	1651	26.8
Afternoon (1200 to 1759 hrs)	2946	47.8
Evening (1800 to 2159 hrs)	524	8.5
Night (2200 to 0559 hrs)	121	2.0
<b>Site of bite injury†</b>		
Head and Neck	237	3.8
Trunk	156	2.5
Upper Limb	1339	21.7
Lower Limb	4424	71.7
Others	23	0.4
<b>Biting animal</b>		
Dog	6048	98.0
Other	124	2.0
<b>Category of bite</b>		
I	1	0.02
II	2613	42.3
III	3558	57.6

\*Time of bite was not available for 10 cases

†Percentages do not add up to 100 because of multiple bites in some cases

<https://doi.org/10.1371/journal.pntd.0007477.t002>

Table 3 analyzes the relationship between the site of bite versus age and sex. The site of the bite was significantly associated with age ( $p < 0.0001$ ). Lower limbs were the most common site of bite across all age and sex groups. However, under-five children were more likely to be bitten on the head, neck and trunk—adjusted standardized residual (ASR) 8.3 and 8.2, respectively. Young adults were more likely to be bitten on lower limbs than expected (ASR 3.7). Males were more likely to be bitten on lower limbs (ASR 2.4) and females were more likely to be bitten on upper limbs (ASR 2.7) than expected.

Analysis restricted to patients >15 years of age (Table 4) further revealed that as compared to females, males were more likely than expected to be bitten on the head and neck region (ASR 2.9). Among children up to 15 years of age, there was no significant relationship between sex and the site of animal bite.

## Discussion

This study shows that dog bite-related injuries are very common in Kashmir. At the ARC, category III dog bites on the lower limbs were the most common.

We analyzed data from the only ARC in Kashmir which mostly provides services to people from District Srinagar. The estimated incidence of animal bites in Srinagar (450 per 100,000 population) is much higher than the incidence reported from Kenya [13] (284 per 100,000), Tanzania [14] (74 per 100,000) and Ghana [15] (54.1 per 100,000). This may, however, be an underestimation of the actual animal bite incidence since many animal bite victims might not have reported to the health care services, as has been reported in an earlier multicentric study from India [16].

Table 3. Relationship of site of bite versus age and sex.

		Site of bite				Total	p-value
		Head and neck	Trunk	Upper limbs	Lower limbs		
Age (years)	< = 5	48 (11.3%) [8.3]	36 (8.5%) [8.2]	96 (22.6%) [0.5]	245 (57.6%) [-6.8]	425	<0.0001
	6–15	43 (4.4%) [1.0]	43 (4.4%) [4.1]	185 (18.9%) [-2.3]	708 (72.3%) [0.3]	979	
	16–30	40 (2.6%) [-2.8]	22 (1.5%) [-3.0]	306 (20.2%) [-1.6]	1147 (75.7%) [3.7]	1515	
	31–50	73 (3.2%) [-1.9]	34 (1.5%) [-3.9]	524 (23.1%) [2.1]	1633 (72.1%) [0.2]	2264	
	51–65	25 (3.3%) [-0.9]	15 (2.0%) [-1.1]	182 (23.7%) [1.4]	547 (71.1%) [-0.5]	769	
	>65	7 (3.7%) [-0.1]	4 (2.1%) [-0.4]	39 (20.5%) [-0.4]	140 (73.7%) [0.5]	190	
Sex	Male	180 (4.0%) [1.0]	108 (2.4%) [-1.0]	942 (20.8%) [-2.7]	3288 (72.8%) [2.4]	4518	0.0306
	Female	56 (3.4%) [-1.0]	46 (2.8%) [1.0]	390 (24.0%) [2.7]	1132 (69.7%) [-2.4]	1624	
<b>Total</b>		236	154	1332	4420	6142*	

\*Those with bites at multiple sites and 'other' sites were excluded from this analysis

Figures in parentheses are row percentages

Figures in square brackets are adjusted standardized residuals

<https://doi.org/10.1371/journal.pntd.0007477.t003>

The cases were mostly young men (Table 1), but children and elderly were not spared. The youngest victim was an infant and the oldest victim was 97 years old. About 23% were children <15 years of age. Males outnumbered females. In Kashmir, males usually go out more frequently for work or for social visits as compared to females. Most of the available literature on dog bites has reported a male predominance [13, 14, 17–22]. The percentage of child victims varies across studies from 25% to 72% [13–20, 23, 24].

Table 4. Sex versus site of bite in animal bite victims >15 years of age.

Sex	Site of bite				Total	p-value
	Head and neck	Trunk	Upper limbs	Lower limbs		
Male	123 (3.5%) [2.9]	51 (1.4%) [-1.3]	741 (21.0%) [-3.2]	2611 (74.0%) [2.3]	3527	0.0002
Female	22 (1.8%) [-2.9]	24 (2.0%) [1.3]	309 (25.5%) [3.2]	856 (70.7%) [-2.3]	1211	
<b>Total</b>	145	75	1051	3467	4738*	

\*Those with bites at multiple sites and 'other' sites were excluded from this analysis

Figures in parentheses are row percentages

Figures in square brackets are adjusted standardized residuals

<https://doi.org/10.1371/journal.pntd.0007477.t004>

The ARC is situated in District Srinagar and that might be the reason why most of the cases in our study were from Srinagar. District hospitals and some of the sub-district level hospital in other districts of the region provide bite-management services to people in those areas.

Our analysis shows a definite rise in the number of cases during the months of March to May (Fig 1). Kashmir is a valley with the winter season starting in December and ending in February. People usually prefer to stay indoors during the period. Outdoor movement of people increases as the winter ends. This might have led to increased interaction with dogs and hence an increase in dog bite cases during the months following winter. A somewhat similar trend has been reported from Iran and Korea [17, 25]; Park et al [25], however, relate it to the length of the day. However, in a five-year study from Srilanka, Kularatne et al reported a more or less even pattern across seasons [20].

The afternoon was the most common time of animal bites (47.7%) followed by forenoon (26.7%) (Table 2). Every seventh case was bitten when he/she ventured out early morning, which people usually do for the morning prayers or for a visit to the bakery or some other work.

The most common site of the bite was lower limbs followed by upper limbs (Table 2). Seven patients had bites at more than one region. Consumption of milk from a rabid cow was the most common among the “others” category. Our results are similar to results from Kenya [13] and Nigeria [19]. Interestingly, an earlier study from Jos Plateau State of Nigeria [18] reported that 85% of victims were bitten on arms.

We found a significant relationship between a victim’s age and the site of an animal bite ( $p < 0.0001$ ) (Table 3). Lower limbs were the most common site of bite across all age and sex groups. A breakdown of table 3 using adjusted standardized residuals revealed that under-five and young children were more likely to be victims of an animal bite on the head, neck, and trunk.

Our results revealed a significant relationship between sex and the site of an animal bite ( $p = 0.0308$ , Table 3). In order to further delve into the relationship, we restricted our analysis to patients >15 years of age (Table 4) hypothesizing that any sex differentials in this context will come into play after puberty. The analysis revealed an interesting relationship. As compared to females, males were more likely to report with a bite in the head and neck region. Furthermore, upper limbs were a more likely site for an animal bite among females.

A dog was the biting animal in 98% of cases. Records lacked information about dog ownership. However, our experience with dog bites at the ARC has been that most of the cases are due to street dogs as people in the region do not usually keep a dog as a pet. Other animals included leopard, cat, and cow. Sudarshan MK et al [16] have reported dog bites to be the most frequent cause of rabies cases in India.

More than half of the cases (57.6%) were classified as category III bites and only one category I bite was recorded. The under-representation of category I bites might be because of several reasons—lack of knowledge among the population about the need for PEP in category I bites, the capacity of the health workers to recognize, treat, and report such bites, and the referral nature of the ARC.

Animal bite victims suffer huge losses in the form of lost wages, travel costs, and direct treatment costs. Moreover, psychological and emotional denting is something which is difficult to translate [2]. Scarring is a common consequence related to dog bites, and the resulting emotional distress due to cosmetic reasons should not be undermined, particularly for wounds on the face.

Measures which can reduce the burden of human rabies include provision of PEP to animal bite victims, pre-exposure prophylaxis of people at risk, increasing awareness about rabies and the need for PEP, and vaccinating dogs against rabies.



The biggest challenges in Africa and Asia, including Kashmir in particular, are free-roaming dog populations, limited veterinary and human health infrastructure, and absence of efficient communication between the veterinary and the human health sectors [26, 27]. The absence of effective control over rabies in the dog population is proving costly in terms of DALYs, premature death and the cost of PEP to public and private health sectors [27, 28]. There are conflicting reports about the number of stray dogs in Srinagar with numbers varying from as low as 22,000 to as high as 150,000 [29, 30]. Whatever the actual number of stray dogs, there is a dire need to devise a strategy to control the alarmingly growing dog population. Unfortunately, there is no provision in our region for isolating suspected dogs or for diagnosing animal rabies. Measures like mass dog vaccination campaigns to improve herd immunity, as recommended by veterinarians, can reduce the burden of human rabies [31]. Animal rabies control requires more attention and must be done through local municipalities. Rabies elimination programs focused mainly on mass vaccination of dogs are largely justified by the future savings of human rabies prevention programs.

### Limitations

This study was based on an analysis of records from an anti-rabies clinic. We, therefore, could not obtain information about the characteristics of the biting animal such as ownership or the circumstances of bite such as provocation. Information was not available about the receipt of tetanus vaccination and ERIG. Because of the secondary nature of the data, we could not validate the categorization of bite. However, during the study period bite categorization was done by trained health workers. The percentage of patients who completed PEP was also not available. We also could not evaluate the impact of animal bites on the physical and psychological health of the victims. An economic burden evaluation was also not possible. We could not draw the real picture of dog bite victims in Kashmir, as data analysis was limited to our hospital and many patients visit hospitals in the peripheries as well while some others miss PEP due to lack of knowledge.

### Conclusions

This study highlights the magnitude of animal bite injuries in Kashmir. Ninety-eight percent of patients had suffered injury from dogs. Category III bites on the lower limbs are the most common type of bite reported at the Anti-Rabies Clinic. Dog bites don't spare children and geriatric population. Children are more prone to be bitten on the head, neck, and trunk. It is time to take proactive measures to stop the menace of the growing animal bite cases and potential rabies deaths.

### Supporting information

**S1 STROBE Checklist. STROBE checklist.**

(DOC)

**S1 Data. Anonymized data sheet.**

(XLSX)

### Author Contributions

**Conceptualization:** Khalid Bashir, Inaamul Haq, S. Muhammad Salim Khan, Mariya Amin Qurieshi.

**Data curation:** Inaamul Haq.



**Formal analysis:** Khalid Bashir, Inaamul Haq.

**Investigation:** Inaamul Haq, Mariya Amin Qurieshi.

**Methodology:** Khalid Bashir, Inaamul Haq, S. Muhammad Salim Khan, Mariya Amin Qurieshi.

**Project administration:** Khalid Bashir, Inaamul Haq, S. Muhammad Salim Khan, Mariya Amin Qurieshi.

**Resources:** Inaamul Haq.

**Software:** Khalid Bashir, Inaamul Haq.

**Supervision:** Khalid Bashir, Inaamul Haq, S. Muhammad Salim Khan, Mariya Amin Qurieshi.

**Validation:** Khalid Bashir, Inaamul Haq, S. Muhammad Salim Khan, Mariya Amin Qurieshi.

**Visualization:** Khalid Bashir, Inaamul Haq.

**Writing – original draft:** Khalid Bashir, Inaamul Haq.

**Writing – review & editing:** Khalid Bashir, Inaamul Haq, S. Muhammad Salim Khan, Mariya Amin Qurieshi.

## References

1. Hemachudha T, Ugolini G, Wacharapluesadee S, Sungkarat W, Shuangshoti S, Laothamatas J. Human rabies: neuropathogenesis, diagnosis, and management. *Lancet Neurol*. 2013 May; 12(5):498–513. [https://doi.org/10.1016/S1474-4422\(13\)70038-3](https://doi.org/10.1016/S1474-4422(13)70038-3) PMID: 23602163
2. Hampson K, Coudeville L, Lembo T, Sambo M, Kieffer A, Attlan M, et al. Estimating the Global Burden of Endemic Canine Rabies. Carvalho MS, editor. *PLoS Negl Trop Dis*. 2015 Apr 16; 9(4):e0003709. <https://doi.org/10.1371/journal.pntd.0003709> PMID: 25881058
3. Expert Consultation on Rabies, World Health Organization, editors. WHO Expert Consultation on Rabies: second report; [Geneva from 18 to 20 September 2012]. Geneva: WHO; 2013. 139 p. (WHO technical report series).
4. WHO | Rabies vaccines: WHO position paper [Internet]. WHO. [cited 2019 Feb 13]. Available from: [http://www.who.int/rabies/resources/who\\_wer8532/en/](http://www.who.int/rabies/resources/who_wer8532/en/).
5. Bahonar A., Rashidi H, Simani S, Fayaz A, Haghdoost A., Rezaei-nassab M, et al. Relative frequency of animal rabies and factors affecting it in Kerman province, 1993–2003. *J Sch Public Health Inst Public Health Res* [Internet]. 2007; 5(1). Available from: <http://sjsph.tums.ac.ir/article-1-186-en.html>.
6. Expert Consultation on Rabies, World Health Organization, editors. WHO Expert Consultation on Rabies: first report; [Geneva from 5 to 8 October 2004]. Geneva: WHO; 2005. 88 p. (WHO technical report series).
7. Khan SMS. Socioeconomic burden due to ever-increasing stray dog population in Jammu and Kashmir, India. Poster session presented at: Reducing the Economic and Social Burden of Rabies in Asia—Strategic Approach. 3rd Rabies in Asia Conference: RIACON; 2011 Nov 28–30; Colombo, Sri Lanka.
8. WHO | Global elimination of dog-mediated human rabies [Internet]. WHO. [cited 2019 Feb 13]. Available from: [http://www.who.int/rabies/resources/who\\_htm\\_ntd\\_nzd\\_2016.02/en/](http://www.who.int/rabies/resources/who_htm_ntd_nzd_2016.02/en/).
9. Census of India Website: Office of the Registrar General & Census Commissioner, India [Internet]. [cited 2019 Feb 13]. Available from: [http://www.censusindia.gov.in/2011census/population\\_enumeration.html](http://www.censusindia.gov.in/2011census/population_enumeration.html).
10. Expert Consultation on Rabies, World Health Organization, editors. WHO Expert Consultation on Rabies: third report; [Bangkok from 26 to 28 April 2017]. Geneva: WHO; 2018. 183 p. (WHO technical report series).
11. Census of India Website: Office of the Registrar General & Census Commissioner, India [Internet]. [cited 2019 Feb 13]. Available from: [http://censusindia.gov.in/vital\\_statistics/SRS\\_Bulletins/Bulletins.html](http://censusindia.gov.in/vital_statistics/SRS_Bulletins/Bulletins.html).
12. Agresti A. An introduction to categorical data analysis. 3rd ed. Hoboken, NJ: John Wiley & Sons; 2019. 375 p.

13. Ngugi JN, Maza AK, Omolo OJ, Obonyo M. Epidemiology and surveillance of human animal-bite injuries and rabies post-exposure prophylaxis, in selected counties in Kenya, 2011–2016. *BMC Public Health* [Internet]. 2018 Dec [cited 2019 Feb 13];18(1). Available from: <https://bmcpublihealth.biomedcentral.com/articles/10.1186/s12889-018-5888-5>.
14. De Nardo P, Gentilotti E, Vairo F, Nguhuni B, Chaula Z, Nicastrì E, et al. A retrospective evaluation of bites at risk of rabies transmission across 7 years: The need to improve surveillance and reporting systems for rabies elimination. Markotter W, editor. *PLoS ONE*. 2018 Jul 2; 13(7):e0197996. <https://doi.org/10.1371/journal.pone.0197996> PMID: 29965994
15. Punguyire DT, Osei-Tutu A, Aleser EV, Letsa T. Level and pattern of human rabies and dog bites in Techiman Municipality in the Middle Belt of Ghana: a six year retrospective records review. *Pan Afr Med J* [Internet]. 2017 [cited 2019 Feb 13];28. Available from: <http://www.panafrican-med-journal.com/content/article/28/281/full/>.
16. Sudarshan MK, Madhusudana SN, Mahendra BJ, Rao NSN, Narayana DHA, Rahman SA, et al. Assessing the burden of human rabies in India: results of a national multi-center epidemiological survey. *Int J Infect Dis* 2007; 11:29–35. <https://doi.org/10.1016/j.ijid.2005.10.007> PMID: 16678463
17. Khazaei S, Karami M, Veisani Y, Solgi M, Goodarzi S. Epidemiology of Animal Bites and Associated Factors with Delay in Post-Exposure Prophylaxis; A Cross-Sectional Study. *Bull Emerg Trauma*. 2018 Jun 1; 6(3):239–44. <https://doi.org/10.29252/beat-060309> PMID: 30090820
18. Alabi O, Nguku P, Chukwukere S, Gaddo A, Nsubuga P, Umoh J. Profile of dog bite victims in Jos Plateau State, Nigeria: a review of dog bite records (2006–2008). *Pan Afr Med J*. 2014; 18(Suppl 1):12.
19. Ogundare EO, Olatunya OS, Oluwayemi IO, Inubile AJ, Taiwo AB, Agaja OT, et al. Pattern and outcome of dog bite injuries among children in Ado-Ekiti, Southwest Nigeria. *Pan Afr Med J* [Internet]. 2017 [cited 2019 Feb 13];27. Available from: <http://www.panafrican-med-journal.com/content/article/27/81/full/>.
20. Kularatne SAM, Ralapanawa DMP, Weerakoon K, Bokalamulla UK, Abagaspiya N. Pattern of animal bites and post exposure prophylaxis in rabies: A five year study in a tertiary care unit in Sri Lanka. *BMC Infect Dis* [Internet]. 2015 Dec [cited 2019 Feb 13];16(1). Available from: <http://www.biomedcentral.com/1471-2334/16/62>.
21. Sudarshan MK, Mahendra BJ, Madhusudana SN, Narayana DHA, Rahman A, Rao NSN, et al. An epidemiological study of animal bites in India: results of a WHO sponsored national multi-centric rabies survey. *J Commun Dis* 2006; 38(1):32–9. PMID: 17370688
22. Ichhpujani RL, Mala C, Veena M, Singh J, Bardwaj M, Bhattacharya D, et al. Epidemiology of animal bites and rabies cases in India: a multicentric study. *J Commun Dis* 2008; 40(1):27–36. PMID: 19127666
23. Gautret P, Le Roux S, Faucher B, Gaudart J, Brouqui P, Parola P. Epidemiology of urban dog-related injuries requiring rabies post-exposure prophylaxis in Marseille, France. *Int J Infect Dis*. 2013 Mar; 17(3):e164–7. <https://doi.org/10.1016/j.ijid.2012.09.011> PMID: 23116608
24. Abubakar S, Bakari A. Incidence of dog bite injuries and clinical rabies in a tertiary health care institution: A 10-year retrospective study. *Ann Afr Med*. 2012; 11(2):108. <https://doi.org/10.4103/1596-3519.93534> PMID: 22406671
25. Park JW, Kim DK, Jung JY, Lee SU, Chang I, Kwak YH, et al. Dog-bite injuries in Korea and risk factors for significant dog-bite injuries: A 6-year cross-sectional study. *PLoS ONE* 2019; 14(2): e0210541. <https://doi.org/10.1371/journal.pone.0210541> PMID: 30789915
26. Knobel DL, Cleaveland S, Coleman PG, Fèvre EM, Meltzer MI, Miranda MEG, et al. Re-evaluating the burden of rabies in Africa and Asia. *Bull World Health Organ*. 2005;11.
27. Meltzer MI, Rupprecht CE. A Review of the Economics of the Prevention and Control of Rabies: Part 2: Rabies in Dogs, Livestock and Wildlife. *Pharmacoeconomics*. 1998; 14(5):481–98. <https://doi.org/10.2165/00019053-199814050-00003> PMID: 10344914
28. Anderson A, Shwiff SA. The Cost of Canine Rabies on Four Continents. *Transbound Emerg Dis*. 2015 Aug; 62(4):446–52. <https://doi.org/10.1111/tbed.12168> PMID: 24112194
29. In Srinagar dog population goes down as dog bite cases increases [Internet]. [cited 2019 Feb 13]. Available from: <https://www.outlookindia.com/website/story/in-srinagar-dog-population-goes-down-says-jk-govt-as-dog-bite-cases-increases-fr/311678>.
30. Kashmir: When the dogs came marching in—The Express Tribune Blog [Internet]. [cited 2019 Feb 13]. Available from: <https://blogs.tribune.com.pk/story/11594/kashmir-when-the-dogs-came-marching-in/>.
31. Gongal G, Wright AE. Human Rabies in the WHO Southeast Asia Region: Forward Steps for Elimination. *Adv Prev Med*. 2011; 2011:1–5.