

Readers are invited to submit letters for publication in this department. Submit letters online at <http://joem.edmgr.com>. Choose "Submit New Manuscript." A signed copyright assignment and financial disclosure form must be submitted with the letter. Form available at www.joem.org under Author and Reviewer information.

Predictors of COVID-19 Vaccine Uptake in Health Care Workers: Correspondence

To the Editor:

We read the publication on "Predictors of COVID-19 Vaccine Uptake in Healthcare Workers: A Cross-sectional Study in Greece."¹ Galanis et al¹ discovered that increasing self-perceived knowledge about the COVID-19 pandemic and increased faith in COVID-19 vaccines were linked to COVID-19 vaccine uptake and recommended that policymakers and scientists devise creative measures to promote COVID-19 vaccine uptake among

healthcare workers. Vaccine apprehension is still a major issue around the world. COVID-19 vaccination apprehension could be caused by a variety of circumstances. Healthcare workers' understanding may be limited at times, which may contribute to vaccination rejection.² Furthermore, the background status of the COVID-19 outbreak may have an impact on acceptance. The temporal likelihood of acceptance was noted in a recent work.³ As a result, a continuous monitoring of the problem and an in-depth investigation of the problem are required. As a result, it is required to conduct repeated continuous monitoring of the problem and an in-depth investigation of healthcare workers' knowledge and attitude.

Rujittika Mungmunpantipantip, PhD
Private Academic Consultant
Bangkok, Thailand

Viroj Wiwanitkit, MD
Dr DY Patil University
Pune, India

REFERENCES

1. Galanis P, Moisoglou I, Vraka I, Siskou O, Konstantakopoulou O, Katsiroumpa A, Kaitelidou D. Predictors of COVID-19 vaccine uptake in healthcare workers: a cross-sectional study in Greece. *J Occup Environ Med.* 2022;64(4):e191–e196.
2. Apajitt P, Wiwanitkit V. Recognition on possibility of no-fever and no-history of travel to risk country among nurse in rural area of a country that COVID-19 already existed: observation. *Int J Prev Med.* 2020;11:47.
3. Xiao J, Cheung JK, Wu P, Ni MY, Cowling BJ, Liao Q. Temporal changes in factors associated with COVID-19 vaccine hesitancy and uptake among adults in Hong Kong: serial cross-sectional surveys. *Lancet Reg Health West Pac.* 2022;23:100441.

The authors report no financial support.

The authors report no conflicts of interest.

Address correspondence to: Rujittika Mungmunpantipantip, PhD, Private Academic Consultant, 111 Bangkok 122, Bangkok 103300 Thailand (rujittika@gmail.com).

ORCID 0000-0003-0078-7897 (R.M.)

ORCID 0000-0003-1039-3728 (V.W.)

Copyright © 2022 American College of Occupational and Environmental Medicine

DOI: 10.1097/JOM.0000000000002582