



BRIEF COMMUNICATION

Occupational inequalities in mortality in Korea: an analysis using nationally representative mortality follow-up data from the late 2000s and after

Eunjeong Noh¹, Young-Ho Khang^{1,2}¹Institute of Health Policy and Management, Seoul National University Medical Research Center, Seoul, Korea; ²Department of Health Policy and Management, Seoul National University College of Medicine, Seoul, Korea

Many Korean and international studies have found higher mortality rates and poorer health conditions among manual workers than among non-manual workers. However, a recent study using unlinked data argued that since the economic crisis in Korea in the late 2000s, the mortality estimates of male Korean non-manual workers have been higher than those of manual workers. Our work using individually linked data from the late 2000s and after aimed to examine mortality inequality by occupational class. We analyzed Korea National Health and Nutrition Examination Survey data that were individually linked to cause-of-death data. Cox regression analysis was used to identify the hazard ratios for mortality by occupational class. Of 11,766 males aged between 35 and 64, 397 died between 2007 and 2018: 142 died from cancer, 68 from cardiovascular disease, 88 from external causes, and 99 from other causes. After controlling for age, the mortality estimates for manual workers were 1.85 times higher than those for upper non-manual workers ($p < 0.05$). We observed no evidence of reversed mortality inequality among occupational classes in Korea since the 2000s; this previously reported finding might have been due to numerator-denominator bias arising from the use of unlinked data.

KEY WORDS: Health inequalities, Mortality, Occupations, Republic of Korea, Socioeconomic factors

INTRODUCTION

Non-manual workers are known to have better health and lower mortality rates than manual workers [1-7]. However, a recent study using unlinked data reported that this general pattern appeared to be different in Korea and Japan [8]. That study claimed that non-manual workers' mortality estimates among Korean males aged 35-64 have risen since the 2000s economic crisis, be-

coming higher than those of manual workers. This purported reversed occupational inequality in mortality could be attributed to a sharp increase in cancer-related and suicide-related deaths among non-manual workers since the economic crisis [8]. The authors of the study suggested that the national economic crisis in the late 2000s might have negatively influenced the health risk factors affecting non-manual workers' mortality. However, another recent study using data on males aged 35-64—the same sex and age group used in the prior study [8]—in Korea from 2007 to 2009 found that socioeconomic indicators, such as education, income, parental education, and economic activity, as well as work environment indicators, were more favorable among non-manual workers than among manual workers, and the prevalence of unfavorable health-related behavioral indicators, such as smoking, high-risk alcohol consumption, depression, and suicidal ideation, was still higher among manual workers than among their counterparts [9]. The previously reported finding of a reversed pattern of mortality inequality among occupational classes in Korea may have been caused by numerator-denominator bias, since the

Correspondence: Young-Ho KhangDepartment of Health Policy and Management, Seoul National University College of Medicine, 103 Daehak-ro, Jongno-gu, Seoul 03080, Korea
E-mail: yhkhang@snu.ac.kr

Received: Jan 4, 2022 / Accepted: Apr 6, 2022 / Published: Apr 6, 2022

This article is available from: <https://e-epih.org/>

© This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

© 2022, Korean Society of Epidemiology

study used population and death counts from different unlinked data sources [9-12]. Although individually linked data from the late 1990s and early 2000s in Korea have provided evidence for occupational mortality inequalities favoring non-manual workers [5,13,14], no research has been done using individually linked cohort data from the late 2000s. An analysis of linked cohort data from the late 2000s to the late 2010s could help determine whether there is, in fact, a reversed pattern of occupational inequality in mortality in Korea. This study aimed to determine mortality estimates by occupational class in Korea since the 2000s.

MATERIALS AND METHODS

Population and sample

This study linked data from the 2007-2015 Korea National Health and Nutrition Examination Surveys (KNHANES) to the 2007-2018 cause-of-death data provided by the Korea Disease Control and Prevention Agency (KDCA). The KNHANES is an annual nationwide survey conducted to understand the health and nutritional status of Korean citizens based on Section 16 of the National Health Promotion Act [15]. Of the 53,101 KNHANES respondents from 2007 to 2015, 51,603 (22,083 males, 29,520 females) consented to linkage with cause-of-death data and had valid unique personal identification numbers. Among them, 12,505 males aged 35-64 were eligible for the study purpose, as a prior study examined occupational mortality inequality among males aged 35-64 [8]. With the exclusion of 739 individuals due to missing information on main indicators, the final sample included 11,766 individuals [15]. Individual data linkage was performed internally at the KDCA, and data without any personal information were provided for our analysis.

Variables

The occupational information provided by the KNHANES from 2007 to 2015 was the independent variable. Four occupational groups were identified based on the same definitions used by Tanaka et al. [8]: the upper non-manual group included managers and professionals; the lower non-manual group included clerks and service and sales workers; the manual group included craft workers, plant and machine operators and assemblers, and elementary

occupations; and the “others” group included workers in other jobs (e.g., agricultural, forestry and fishery workers and unemployed individuals). Data on all-cause and cause-specific deaths by occupational class between 2007 and 2018 were identified. Four major causes of death were considered, including cancer (International Classification of Diseases, 10th revision [ICD-10] codes: C00-D48), cardiovascular disease (ICD-10 codes: I00-199), external causes (ICD-10 codes: V01-Y89), and other causes.

Statistical analysis

Using information on survival time and death status, Cox proportional hazards regression models were employed to analyze the hazard ratios (HRs) of all-cause and cause-specific mortality among occupational classes, after adjusting for age.

Ethics statement

The study protocol was approved by the Institutional Review Board (IRB) of Seoul National University Hospital (IRB No. E-2006-058-1131). Informed consent was waived by the IRB.

RESULTS

From 2007 to 2018, 397 of the 11,766 individuals died (3.4%): 142 (35.8%) died from cancer (ICD-10 codes: C00-D48), 68 (17.1%) from cardiovascular diseases (I00-199), 88 (22.2%) from external causes (V01-Y89), and 99 (24.9%) from other causes. The mean \pm standard deviation (SD) age of all subjects was 47.84 ± 10.48 years, and the mean follow-up period for all subjects was 7.40 ± 4.89 years. Table 1 presents the mean age and mean follow-up period by occupational class. The mean age was similar among upper non-manual and lower non-manual workers, but relatively older among manual workers and the “others” group. The mean follow-up period by occupational class showed a graded pattern reflecting mortality risk differentials among occupational classes. The all-cause mortality estimate of the manual workers after age adjustment was 1.85 times higher than those of upper non-manual workers ($p < 0.05$). The HR of lower non-manual workers tended to be higher than that of upper non-manual workers (Table 1).

Age stratified analyses (Supplementary Material 1) showed that the mortality risk of manual workers was generally greater than

Table 1. Numbers of subjects and deaths, mean age, mean follow-up duration, and age-adjusted HRs and their 95% CIs for all-cause mortality¹ by occupational class in Korean males aged 35 to 64 (n=11,766)

Variables	No. of subjects	No. of deaths	Age, mean \pm SD (yr)	Follow-up duration, mean \pm SD (yr)	HR (95% CI)	p-value
Non-manual						
Upper	2,276	35	45.68 \pm 8.49	7.50 \pm 4.03	1.00 (reference)	
Lower	3,069	52	45.34 \pm 8.68	7.40 \pm 3.87	1.31 (0.81, 2.13)	0.28
Manual	3,931	121	48.28 \pm 9.34	7.37 \pm 4.23	1.85 (1.20, 2.85)	0.01
Others	2,490	189	52.97 \pm 9.97	7.35 \pm 4.36	3.38 (2.14, 5.33)	<0.001

HR, hazard ratio; CI, confidence interval; SD, standard deviation.

¹Mortality follow-up data from the 2007 and 2015 Korea National Health and Nutrition Examination Surveys.

Table 2. Age-adjusted HRs and their 95% CIs for mortality¹ from 4 broad causes by occupational class in Korean males aged 35 to 64 (n=11,766)

Variables	Non-manual		Manual	Others
	Upper	Lower		
No. of subjects	2,276	3,069	3,931	2,490
No. of cancer deaths	14	17	34	77
HR (95% CI)	1.00 (reference)	1.09 (0.47, 2.50)	1.34 (0.62, 2.90)	2.39 (1.19, 4.79)
p-value		0.85	0.46	0.01
No. of cardiovascular deaths	5	10	21	32
HR (95% CI)	1.00 (reference)	1.46 (0.47, 4.53)	2.06 (0.74, 5.74)	3.25 (1.10, 9.57)
p-value		0.52	0.17	0.03
No. of external causes	10	18	34	26
HR (95% CI)	1.00 (reference)	1.56 (0.67, 3.60)	2.07 (0.95, 4.51)	3.17 (1.25, 8.06)
p-value		0.30	0.07	0.02
No. of other deaths	6	7	32	54
HR (95% CI)	1.00 (reference)	1.11 (0.30, 4.16)	2.53 (0.80, 7.98)	5.79 (1.80, 18.64)
p-value		0.87	0.11	<0.001

HR, hazard ratio; CI, confidence interval.

¹Mortality follow-up data from the 2007 and 2015 Korea National Health and Nutrition Examination Surveys.

that of non-manual workers among both males aged 35-49 and males aged 50-64. In Korean females aged 35-64, the mortality differentials between manual and non-manual workers were not significant, partly because of the small numbers of deaths, but the mortality risk of manual workers tended to be greater than that of non-manual workers (Supplementary Material 2).

The cause-specific analysis results showed that, largely because of the small numbers of cause-specific deaths, manual workers did not have statistically significantly higher cause-specific mortality estimates than upper non-manual workers. However, those in the “others” group, who were not classified as manual workers but were either agricultural, forestry, or fishery workers or were unemployed, had much higher mortality estimates (all statistically significant) than upper non-manual workers (Table 2).

DISCUSSION

This study utilized cohort data that linked the 2007-2015 KNHANES data with 2007-2018 cause-of-death data. By tracking deaths and the causes of death in Korean males aged 35-64, all-cause mortality estimates according to occupational class and mortality estimates based on 4 causes of death were derived. Even after the economic crisis in the late 2000s, the mortality of male Korean manual workers aged 35-64 remained higher than that of non-manual workers. Therefore, there was no evidence to support the finding of reversed mortality inequality by occupational class reported by the previous study [8], which used unlinked census and death registry data. Our study is consistent with the universal conclusions of previous domestic and international studies using mortality follow-up data, which have shown that the mortality estimates of manual workers are higher than those of non-manual workers [1-7].

Additionally, the prior study proposed that the Korean economic crisis in the 2000s could have negatively affected non-manual workers' psychosocial health factors, such as stress and depression, which might have contributed to increased mortality due to suicide and cancer [8]. However, although cause-specific mortality estimates were not statistically significantly higher in manual workers than in non-manual workers due to the small numbers of cause-specific deaths in the present study, the HRs for cancer and external causes tended to be higher in manual workers than in non-manual workers (Table 2). The mortality estimates of the “others” group were exceptionally higher than those of non-manual workers, and this discrepancy was statistically significant. Moreover, our recent study using data from the late 2000s showed that psychosocial factors (e.g., depression, stress, suicidal ideation), work environment, and health behaviors related to cancer (e.g., smoking, high-risk alcohol consumption) were more favorable in non-manual workers than in manual workers [9].

In summary, like numerous previous studies on mortality and the prevalence of health risk factors by occupational class, this study found that in Korea, manual workers continued to have higher mortality rates than non-manual workers in the late 2000s and after. This result contradicts the prior claim of reversed mortality inequality by occupational class in Korea, which might be attributed to the denominator-numerator bias that has been commonly raised as a concern in health inequality research using unlinked data. If the census and death data by occupational class are not individually linked, as was the case in the study that proposed the inequality reversal, it is highly likely that the mortality estimates of non-manual workers will be overestimated and those of manual workers underestimated [9-12]. Analyzing the linked data provided via the nationally representative KNHANES (2007-2015) and individual mortality data (2007-2018), this study confirmed

that even after the economic crisis of the 2000s, patterns of mortality inequality in Korea remain disadvantageous for those in lower occupational classes.

SUPPLEMENTARY MATERIALS

Supplementary materials are available at <http://www.e-epih.org/>.

CONFLICT OF INTEREST

The authors have no conflicts of interest to declare for this study.

FUNDING

This research was supported by a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea (grant No. HI18C0446).

ACKNOWLEDGEMENTS

We thank the Korea Disease Control and Prevention Agency for providing the linked data.

AUTHOR CONTRIBUTIONS

Both authors contributed equally to conceiving the study, analyzing the data, and writing this paper.

ORCID

Eunjeong Noh: <https://orcid.org/0000-0003-3669-4676>; Young-Ho Khang: <https://orcid.org/0000-0002-9585-8266>

REFERENCES

- Mackenbach JP, Stirbu I, Roskam AJ, Schaap MM, Menvielle G, Leinsalu M, et al. Socioeconomic inequalities in health in 22 European countries. *N Engl J Med* 2008;358:2468-2481.
- Yeoman K, Sussell A, Retzer K, Poplin G. Health risk factors among miners, oil and gas extraction workers, other manual labor workers, and nonmanual labor workers, BRFSS 2013-2017, 32 states. *Workplace Health Saf* 2020;68:391-401.
- Toch-Marquardt M, Menvielle G, Eikemo TA, Kulhánová I, Kulik MC, Bopp M, et al. Occupational class inequalities in all-cause and cause-specific mortality among middle-aged men in 14 European populations during the early 2000s. *PLoS One* 2014;9:e108072.
- Stringhini S, Sabia S, Shipley M, Brunner E, Nabi H, Kivimaki M, et al. Association of socioeconomic position with health behaviors and mortality. *JAMA* 2010;303:1159-1166.
- Khang YH, Kim HR. Socioeconomic inequality in mortality using 12-year follow-up data from nationally representative surveys in South Korea. *Int J Equity Health* 2016;15:51.
- Khang YH, Kim HR. Relationship of education, occupation, and income with mortality in a representative longitudinal study of South Korea. *Eur J Epidemiol* 2005;20:217-220.
- Son M, Armstrong B, Choi JM, Yoon TY. Relation of occupational class and education with mortality in Korea. *J Epidemiol Community Health* 2002;56:798-799.
- Tanaka H, Nusselder WJ, Bopp M, Brønnum-Hansen H, Kalediene R, Lee JS, et al. Mortality inequalities by occupational class among men in Japan, South Korea and eight European countries: a national register-based study, 1990-2015. *J Epidemiol Community Health* 2019;73:750-758.
- Noh E, Khang YH. Analysis of factors contributing to occupational health inequality in Korea: a cross-sectional study using nationally representative survey data. *Arch Public Health* 2021;79:113.
- Khang YH. The surprising result of manual workers in Korea enjoying lower mortality than non-manual workers is likely due to numerator-denominator bias. Comment on: "Mortality inequalities by occupational class among men in Japan, South Korea and eight European countries: a national register-based study, 1990-2015" [cited 2021 Dec 15]. Available from: <https://jech.bmj.com/content/early/2019/06/07/jech-2018-211715.responses?versioned=true>.
- Kim HR, Khang YH. Reliability of education and occupational class: a comparison of health survey and death certificate data. *J Prev Med Public Health* 2005;38:443-448 (Korean).
- Williams GM, Najman JM, Clavarino A. Correcting for numerator/denominator bias when assessing changing inequalities in occupational class mortality, Australia 1981-2002. *Bull World Health Organ* 2006;84:198-203.
- Khang YH, Lee SI, Lee MS, Jo MW. Socioeconomic mortality inequalities in Korea labor & income panel study. *Health Policy Manag* 2004;14:1-20 (Korean).
- Khang YH, Kim HR. Explaining socioeconomic inequality in mortality among South Koreans: an examination of multiple pathways in a nationally representative longitudinal study. *Int J Epidemiol* 2005;34:630-637.
- Yun S, Oh K. Introduction to the Korea National Health and Nutrition Examination Survey (KNHANS) linked Cause of Death Data. *Public Health Wkly Rep* 2020;13:2071-2080 (Korean).