




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



The effects of a novel adjuvant complex/*Eimeria* profilin vaccine on the intestinal host immune response against live *E. acervulina* challenge infection[☆]

Sung-Hyen Lee^{a,1}, Hyun S. Lillehoj^{a,*}, Seung I. Jang^a, Kyung-Woo Lee^a, Robert J. Yancey^b, Paul Dominowski^b

^a Animal Parasitic Diseases Laboratory, Animal and Natural Resources Institute, Agricultural Research Service-U.S. Department of Agriculture, Beltsville, MD 20705, USA

^b Pfizer Inc., 333 Portage Street, Kalamazoo, MI 49007, USA

ARTICLE INFO

Article history:

Received 1 April 2010

Received in revised form 12 June 2010

Accepted 30 June 2010

Available online 15 July 2010

Keywords:

Adjuvant
Eimeria profilin
Coccidiosis
Immunity
Cytokine
Chicken

ABSTRACT

The effects of a novel adjuvant composed of Quil A, cholesterol, dimethyl dioctadecyl ammonium bromide, and Carbopol (QCDC) on protective immunity against avian coccidiosis following immunization with an *Eimeria* recombinant protein were determined. Broiler chickens were subcutaneously immunized with isotonic saline (control), *Eimeria* recombinant profilin alone, or profilin emulsified with QCDC at 1 and 7 days post-hatch, and orally challenged with live *Eimeria acervulina* at 7 days following the last immunization. Body weight gains, gut lesion scores, fecal oocyst outputs, profilin serum antibody titers, lymphocyte proliferation, and intestinal cytokine transcript levels were assessed as measures of protective immunity. Chickens immunized with profilin plus QCDC showed increased body weight gains and decreased intestinal lesion scores compared with the profilin only or control groups. However, no differences were found in fecal oocyst shedding among the three groups. Profilin serum antibody titers and antigen-induced peripheral blood lymphocyte proliferation in the profilin/QCDC group were higher compared with the profilin only and control groups. Finally, while immunization with profilin alone or profilin plus QCDC uniformly increased the levels of intestinal transcripts encoding all cytokines examined (IL-1 β , IL-10, IL-12, IL-15, IL-17F, and IFN- γ) compared with the control group, transcripts for IL-10 and IL-17F were further increased in the profilin/QCDC group compared with the profilin only group. In summary, this study provides the first evidence of the immunoenhancing activities of QCDC adjuvant in poultry.

Published by Elsevier Ltd.

1. Introduction

Avian coccidiosis, an intestinal disease caused by multiple species of *Eimeria*, has had significant economic effects on the poultry industry [1]. Prophylactic medication has been successfully used to control avian coccidiosis, but alternative strategies are sought due to the increasing emergence of drug-resistant coccidia in commercial production settings [2,3]. Although there is a growing reliance on the use of *Eimeria* parasite vaccines for coccidiosis control [4,5], live vaccines pose the risk of unintended infection under the immunosuppressive conditions associated with high-density commercial rearing conditions. On the other hand, while

subunit protein and DNA vaccines do not present the possibility of field infections, both are of limited immunogenicity and there is a lack of suitable immunoenhancing adjuvants for the poultry industry [6–9].

Profilin is a major immunogenic protein of *Eimeria* and vaccination of chickens with this recombinant polypeptide induces partial protection against avian coccidiosis [10,11]. Profilins are essential components of all Apicomplexan parasites, and are involved in actin-dependent gliding motility, parasite migration across biological barriers, and host cell invasion [10]. The profilin gene encodes a ~20 kDa protein that is highly conserved among the various genera of Apicomplexans and expressed during most stages of the parasite's life cycle [6,11,12]. Toxofilin, the profilin of *Toxoplasma gondii*, binds to TLR11, inducing a potent IL-12 response in murine dendritic cells, and plays a critical role in CD4⁺ T cell-driven immunity to the pathogen [10,13]. In our previous study, profilin was identified from the merozoites of *Eimeria acervulina* as an immunogenic protein that induced high levels of IFN- γ production by splenic T cells [14]. Subcutaneous immunization of young broiler chickens with recombinant profilin protein in the absence of adjuvant, or *in ovo* vaccination of 18-day-old embryos with a DNA plasmid encod-

[☆] Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

* Corresponding author. Tel.: +1 301 504 8771; fax: +1 301 504 5103.

E-mail address: Hyun.Lillehoj@ars.usda.gov (H.S. Lillehoj).

¹ This work was carried out during sabbatical leave from National Academy of Agricultural Science, Rural Development Administration, Suwon, Gyeonggi 441-853, Korea.

ing the profilin gene, provided limited, but statistically significant protection against challenge infection with live *E. acervulina* [6,8].

Recently, substantial progress has been made in the formulation of novel adjuvants that enhance the immunogenicity of peptides and proteins without unwanted side effects [15–17]. For example, saponins such as Quil A are derived from the South American tree *Quillaja saponaria molina* and are currently used in several veterinary applications [18]. Quil A is combined with cholesterol and phospholipids to form immunostimulatory complexes (ISCOMs) that avoid the hemolytic and viricidal properties of Quil A alone [19]. However, ISCOMs may stimulate an undesirable autoimmune reaction, and some hydrophilic proteins cannot be efficiently incorporated into their structure. Alternatively, dimethyl dioctadecyl ammonium bromide (DDA) and avidine are lipophilic quaternary amine adjuvants that stimulate robust cell-mediated and humoral immune responses [20]. The efficacy of DDA as an adjuvant for proteins, haptens, viruses, protozoa, and bacteria has been reported [21–23]. By itself, DDA is sparingly soluble in water and is more effective when used with polymers such as dextran, polyethylene glycol, or polyacrylic acids like Carbopol. QCDC is a new adjuvant complex composed of Quil A, cholesterol, DDA, and Carbopol that enhances immune responses against multiple pathogen vaccines in a variety of veterinary settings [24]. However, the adjuvanticity of QCDC in poultry has not been reported. Therefore, the current study was undertaken to evaluate the adjuvant effect of QCDC during vaccination against avian coccidiosis.

2. Materials and methods

2.1. Experimental animals

Experiments were approved by the Agricultural Research Service Institutional Animal Care and Use Committee (ARS IACUC). One-day-old broiler chickens (Ross/Ross, Longenecker Hatchery, Elizabethtown, PA) were housed in Petersime starter brooder units and randomly assigned to 4 groups (20 birds/group). Chickens were kept in brooder pens in an *Eimeria*-free facility for 2 weeks and then transferred into large hanging cages (2 birds/cage) at a separate location where they were infected and kept until the end of the experimental period.

2.2. Expression and purification of recombinant profilin

The profilin gene was originally cloned by immunoscreening an *E. acervulina* cDNA library using a rabbit antiserum against *E. acervulina* merozoites [6]. The 1086-base pair profilin cDNA was subcloned into the pMAL plasmid (New England Biolabs, Ipswich, MA) with an NH₂-terminal maltose-binding protein epitope tag and a Factor Xa protease cleavage site between maltose-binding protein and profilin. Transformed *Escherichia coli* DH5a bacteria were grown to mid-log phase, induced with 1.0 mM of isopropyl- β -D-thiogalactopyranoside for 3 h at 37 °C, collected by centrifugation, and disrupted by sonication on ice (Misonix, Farmingdale, NY). The recombinant profilin protein was isolated on an amylose affinity column (New England Biolabs) according to the manufacturer's instructions, digested with Factor Xa to release profilin from the solid phase, and repassed through the amylose column to remove any contaminating maltose-binding protein. Final purity was confirmed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and Western blotting with profilin-specific rabbit antibody.

2.3. Profilin immunization

One-day-old chickens were immunized subcutaneously with sterile isotonic saline or with 100 μ g of profilin alone or profilin

plus QCDC in a 50 μ l emulsion. The formulation of QCDC consisted of 12.0 μ g/ml of Quil A (E.M. Sergeant Pulp & Chemical Co., Clifton, NJ), 12.0 μ g/ml of cholesterol (FabriChem, Trumbull, CT), 0.6 μ g/ml of DDA (Fluka, Buchs, Switzerland), and 0.75 mg/ml of Carbopol[®] 974P (Lubrizol, Wickliffe, OH). Booster immunization was given at the same injection site as the primary immunizations at 7 days following primary immunization.

2.4. *Eimeria* infection and measurements of body weight gains and fecal oocyst shedding

At 7 days following secondary immunization, chickens (20/group) were orally infected with 1.0×10^4 sporulated oocysts of *E. acervulina* as described [25,26]. Body weights were measured between 0 and 9 days post-infection. For the determination of fecal oocyst shedding, birds (12/group) were placed in oocyst collection cages and fecal samples were collected between 6 and 9 days post-infection. Oocyst numbers were determined using a McMaster chamber according to the following formula: total oocysts/bird = [oocyst count \times dilution factor \times (fecal sample volume/counting chamber volume)]/2 [27,28].

2.5. Measurement of profilin antibody responses

Blood samples (4 birds/group) were collected by cardiac puncture immediately following euthanasia at 7 days post-secondary immunization. Sera was obtained by low speed centrifugation, and used in an ELISA to measure the profilin-specific antibody response as described [27,29]. In brief, 96-well microtiter plates were coated overnight with 1.0 μ g/well of purified recombinant profilin. As a negative control, wells were coated with a recombinant microneme protein from *E. tenella*, EtMIC2 [8]. EtMIC2 was cloned from *E. tenella* and was shown during host cell invasion to be localized at the point of parasite entry and secreted from the host–parasite interface. The plates were washed with phosphate-buffered saline containing 0.05% Tween (PBS-T) and blocked with PBS containing 1% BSA. Diluted sera (1:100) were added (100 μ l/well), incubated with agitation for 2 h at room temperature, and washed with PBS-T. Bound antibodies were detected with peroxidase-conjugated rabbit anti-chicken IgG and 3,3',5,5'-tetramethylbenzidine substrate (Sigma, St. Louis, MO). Optical density at 450 nm (OD₄₅₀) was measured with an automated microplate reader (Bio-Rad, Richmond, CA). All samples were analyzed in quadruplicate to ensure accuracy.

2.6. Determination of intestinal lesion scores

At 9 days post-infection, lesion scores (5 birds/group) were determined using a numerical scale from 0 (normal) to 4 (severe) [28,30] and evaluated by 3 independent observers.

2.7. Lymphocyte proliferation

Peripheral blood lymphocytes (PBL) were collected from uninfected chickens (4 birds/group) by cardiac puncture immediately after euthanasia at 7 days following secondary immunization as described [31]. The cells were adjusted to 5.0×10^6 cells/ml in RPMI medium containing 10% fetal bovine serum, 100 U/ml penicillin, and 100 μ g/ml streptomycin and incubated in a humidified incubator at 41 °C and 5% CO₂ for 48 h with 10 μ g/ml of profilin in 96-well plates. Cell proliferation was measured using WST-8 (Cell-Counting Kit-8[®], Dojindo Molecular Technologies, Gaithersburg, MD) as described [32] and was expressed as OD₄₅₀ (profilin) – OD₄₅₀ (medium alone). Each sample was analyzed in quadruplicate.

Table 1
Oligonucleotide primers used in quantitative RT-PCR.

Type	mRNA target	Primer sequences	PCR product size (bp)	GenBank accession no.
Reference	GAPDH	F: 5'-GGTGGTCTAAGCGTGTAT-3' R: 5'-ACCTCTGTCTCTCCACA-3'	264	K01458
Pro-inflammatory	IL-1 β	F: 5'-TGGGCATCAAGGGCTACA-3' R: 5'-TCGGGTGGTTGGTGATG-3'	244	Y15006
	IL-17F	F: 5'-CTCCGATCCCTTATTCTCCTC-3' R: 5'-AAGCGGTGTGGTCTCAT-3'	292	AJ493595
Th1	IFN- γ	F: 5'-AGCTGACGGTGGACCTATTATT-3' R: 5'-GGCTTTGCGCTGGATTG-3'	259	Y07922
	IL-12	F: 5'-AGACTCCAATGGGCAAATGA-3' R: 5'-CTCTTCGGCAAATGGACAGT-3'	274	NM.213571
	IL-15	F: 5'-TCTGTCTCTCTGTTCTGAGTGATG-3' R: 5'-AGTGATTGCTTCTGCTTTGGTA-3'	243	AF139097
Th2	IL-10	F: 5'-CGGAGCTGAGGGTGAA-3' R: 5'-GTGAAGAAGCGGTGACAGC-3'	272	AJ621614

2.8. Quantification of cytokine mRNA levels

Intestinal duodenum tissues were obtained from uninfected chickens (4 birds/group) at 7 days post-secondary immunization. The tissues were cut longitudinally and washed 3 times with ice-cold Hanks' balanced salt solution containing 100 U/ml of penicillin and 100 μ g/ml of streptomycin. The mucosal layer was carefully scraped away using a surgical scalpel and total RNA was extracted using TRIzol (Invitrogen, Carlsbad, CA). Five micrograms of total RNA were treated with 1.0 U of DNase I and 1.0 μ l of 10 \times reaction buffer (Sigma) and incubated for 15 min at room temperature. Then, 1.0 μ l of stop solution was added to inactivate DNase I, and the resulting mixture was heated to 70 $^{\circ}$ C for 10 min. RNA was reverse-transcribed using the StrataScript first-strand synthesis system (Stratagene, La Jolla, CA) according to the manufacturer's recommendations. Quantitative RT-PCR oligonucleotide primers for chicken cytokines (IL-1 β , IL-17 F, IFN- γ , IL-12, IL-15 and IL-10) and the GAPDH internal control are listed in Table 1. Amplification and detection were carried out using equivalent amounts of total RNA using the Mx3000P system and Brilliant SYBR Green qPCR master mix (Stratagene). Standard curves were generated using log₁₀ diluted standard RNA and the levels of individual transcripts were normalized to those of GAPDH by the Q-gene program [33]. Each sample was analyzed in triplicate. To normalize individual replicates, the logarithmic-scaled threshold cycle (C_t) values were transformed to linear units of normalized expression prior to calculating means and SEM for the references and individual targets, followed by the determination of mean normalized expression (MNE) using the Q-gene program [32,34].

2.9. Statistical analyses

All data was expressed as mean \pm SEM values of 4–20 chickens/group with 3–4 replicates/sample. Comparisons of the mean values were performed by one-way analysis of variance followed by the Duncan's multiple range test using SPSS software (SPSS 15.0 for Windows, Chicago, IL). Differences between groups were considered statistically significant at $P < 0.05$.

3. Results

3.1. The effects of vaccination with profilin plus QCDC on body weight gain and fecal oocyst shedding

In the absence of profilin immunization, *E. acervulina*-infected chickens showed significantly reduced body weight gains (596.4 ± 11.3 g) compared with the uninfected controls (691.0 ± 15.0 g) (Fig. 1A). Chickens immunized with profilin alone and infected with parasites exhibited weight gains (665.3 ± 8.4 g)

equal to uninfected controls and chickens immunized with profilin plus QCDC exhibited additional weight gains (719.1 ± 17.0 g) compared with the profilin alone group. In contrast, no differences were found in fecal oocyst shedding among the 3 immunized and infected groups (Fig. 1B).

3.2. The effect of vaccination with profilin plus QCDC on profilin serum antibody responses

Serum antibody levels in the profilin plus QCDC group increased compared with those of the control or profilin only groups (Fig. 2A).




Fig. 1. The effects of vaccination with profilin plus QCDC on body weight gain and fecal oocyst shedding. Chickens were immunized 1 and 7 days post-hatch with PBS (control), profilin alone, or profilin plus QCDC and orally infected with 1.0×10^4 sporulated oocysts of *E. acervulina* at 7 days post-secondary immunization. (A) Body weight gains were calculated between 0 and 9 days post-infection. Each bar represents the mean \pm SEM values ($N=20$). (B) Fecal oocyst numbers were measured between 6 and 9 days post-infection. Each bar represents the mean \pm SEM values ($N=12$). Bars not sharing the same letters are significantly different according to the Duncan's multiple range test ($P < 0.05$). NS indicates that the bar is not significant among the experimental groups.




Fig. 2. The effect of vaccination with profilin plus QCDC on profilin serum antibody responses. Chickens were immunized and infected as in Fig. 1 and serum antibodies against profilin (A) and EtMIC2 (B) recombinant protein were measured at 7 days post-secondary immunization. Each bar represents the mean \pm SEM values from quadruplicate samples/bird ($N=4$ birds/group). Bars not sharing the same letters are significantly different according to the Duncan's multiple range test ($P<0.05$). NS indicates that the bar is not significant among the experimental groups.

As a negative control, the levels of antibodies reactive with an unrelated *Eimeria* protein antigen, EtMIC2, were equal in the 3 immunization groups (Fig. 2B).

3.3. The effect of vaccination with profilin plus QCDC on intestinal lesion scores

Intestinal lesion scores in the profilin plus QCDC group (1.62 ± 0.2) decreased compared with those of the control group (2.5 ± 0.2) (Fig. 3). Comparison of scores between the profilin only (2.0 ± 0.2) and profilin plus QCDC groups, or between the control and profilin only groups, revealed no significant differences.

3.4. The effect of vaccination with profilin plus QCDC on lymphocyte proliferation

Profilin-stimulated PBL proliferative responses in the profilin plus QCDC group increased compared with those of the control or profilin only groups (Fig. 4).

3.5. The effect of vaccination with profilin plus QCDC on intestinal cytokine transcript levels

The levels of transcripts encoding proinflammatory (IL-1 β , IL-17F), Th1 (IFN- γ , IL-12, IL-15), or Th2 (IL-10) cytokines in the duodenum increased in the profilin only and the profilin/QCDC groups compared with the control group (Fig. 5). The levels of transcripts encoding IL-17F and IL-10 further increased in the profilin/QCDC group compared with the profilin only group.

4. Discussion

This study documents the immunopotentiating effects of a new adjuvant formulation, QCDC, and its ability to augment the immunogenicity of an *Eimeria* recombinant protein vaccine in




Fig. 3. The effect of vaccination with profilin plus QCDC on intestinal lesion scores. Chickens were immunized and infected as in Fig. 1, and lesions were scored from 0 (normal) to 4 (severe) at 9 days post-infection. Each bar represents the mean \pm SEM values from 3 independent observers ($N=5$ birds/group). Bars not sharing the same letters are significantly different according to the Duncan's multiple range test ($P<0.05$).

chickens. The major findings are that immunization of chickens with profilin plus QCDC reduced body weight loss caused by *Eimeria* infection, induced higher serum antibody titers to profilin, decreased gut pathology, increased antigen-driven PBL proliferation, and increased IL-10 and IL-17F transcript levels in the intestine compared with chickens vaccinated with profilin alone.

Novel vaccination strategies using DNA or proteins derived from mucosal pathogens have successfully induced protective immunity against a variety of infectious diseases including avian coccidiosis [6,35]. For example, DNA immunization with plasmids encoding either the profilin or EtMIC2 *Eimeria* protein has been shown to induce both antigen-specific antibody and T cell responses, and to confer partial protection against live parasite challenge infection [6,8]. However, the levels of protective immunity against avian coccidiosis that are induced by subunit protein vaccines have been limited, due in large part to the complex parasite life cycle that includes intracellular and extracellular as well as sexual and asexual stages [1,14,23]. Each of these stages is associated with a unique pattern of gene expression by the parasite and not all *Eimeria* proteins are expressed in all stages. In addition, individual chickens can be simultaneously or sequentially colonized by multiple *Eimeria* species, and cross-protection of host immunity among the different parasites is severely limited. As a result of these practical limitations, recent interest has developed in using vaccine adjuvants as immunostimulators to enhance the immunogenicity of coccidia subunit vaccines.




Fig. 4. The effect of vaccination with profilin plus QCDC on lymphocyte proliferation. Chickens were immunized as in Fig. 1, PBL were isolated at 7 days post-secondary immunization, and cell proliferation in response to stimulation with 10 μ g/ml of profilin was measured and expressed as ΔOD (OD_{450} profilin – OD_{450} medium alone). Each bar represents the mean \pm SEM values from quadruplicate samples/bird ($N=4$ birds/group). Bars not sharing the same letters are significantly different according to the Duncan's multiple range test ($P<0.05$).




Fig. 5. The effect of vaccination with profilin plus QCDC on intestinal cytokine transcript levels. Chickens were immunized as in Fig. 1, intestinal duodenum tissues were isolated at 7 days post-secondary immunization, and transcripts encoding IL-1 β (A), IL-17A (B), IFN- γ (C), IL-10 (D), IL-12 (E), and IL-15 (F) were quantified by real time RT-PCR, and normalized to GAPDH transcript levels. Each bar represents the mean \pm SEM values from triplicate samples/bird ($N=4$ birds/group). Bars not sharing the same letters are significantly different according to the Duncan's multiple range test ($P<0.05$).

QCDC is an adjuvant complex composed of Quil A, cholesterol, DDA, and Carbopol, and its adjuvant effects have been demonstrated against *E. coli*, the feline leukemia virus, bovine viral diarrhea virus, mycoplasma hyopneumoniae, avian and canine influenza viruses, canine coronavirus, and bovine rotavirus [24]. Quil A by itself has been used as an adjuvant in many veterinary animal species, and defined saponins such as QS-21 are currently used in human clinical trials [36–39]. DDA provides a significant adjuvant effect when co-administered systemically or locally with antigen and stimulates predominantly Th1 cells with no reported toxic effects in humans [40–46]. In avians, DDA enhanced humoral and cell-mediated immune responses to a Newcastle disease virus vaccine and induced greater lymphocyte proliferation compared with complete Freund's adjuvant [22]. Chickens immunized with *Eimeria* merozoite antigens in combination with DDA displayed longer-lasting coccidial immunity compared with the *Corynebacterium parvum* adjuvant [23]. Carbopol is a mucoadhesive polymer that has been extensively investigated due to its high viscosity at low concentrations and low toxicity [47]. Carbopol enhanced the efficacy of recombinant canarypox viral vectors for equine herpes virus vaccination and significantly improved virus neutralizing antibody responses to this virus [48]. No prior studies have examined the effects of Carbopol on poultry vaccines.

The beneficial effects of profilin plus QCDC immunization on body weight gain and intestinal pathology indicate that the mode of action of this vaccine complex may involve effector mecha-

nisms influencing these two clinical parameters of coccidiosis. In this regard, the importance of cell-mediated immunity in conferring protection against avian coccidiosis has been well documented [26,34,49–51], and humoral immunity also appears to be of some assistance [25,28]. Serum antibody levels showed higher titers against profilin in the group immunized with profilin plus QCDC, whereas no significant differences in anti-EtMIC2 antibody levels were seen among these groups (Fig. 2B). The reason why profilin plus QCDC had no effect on parasite fecundity/shedding compared with profilin alone is unclear, but this observation is in agreement with previously published reports indicating the absence of a direct correlation between *Eimeria*-induced body weight loss and fecal oocyst shedding [29,49].

In concordance with the evidence that supports the role of cell-mediated immunity in protection against avian coccidiosis, other studies have verified the significance of the gut cytokine response, including the role of proinflammatory, Th1, and Th2 cytokines following *Eimeria* infection [34,51–55]. In fact, increasing levels of gene transcripts encoding IL-10 and IL-17F were associated with enhanced protection against coccidiosis [55], although their role in local immune response needs to be investigated. IL-1 β , a proinflammatory cytokine produced by macrophages, monocytes and dendritic cells, exerted an adjuvant effect when given with a profilin DNA vaccine and reduced fecal oocyst shedding following oral infection with viable parasites [7]. IL-17 is a cytokine that is secreted exclusively by activated memory T cells which induce fibro-

lasts to secrete other proinflammatory or hematopoietic cytokines [56–61]. IL-17 was highly up-regulated following primary infection with *E. acervulina* or *E. maxima*, but not *E. tenella* [52,60]. Chicken IL-17 induced IL-6 production and both cytokines together may play an immunoregulatory role in the initiation, maintenance, and control of intestinal inflammatory responses [61]. IFN- γ has also been associated with protective immune responses to coccidiosis [26,62,63]. Administration of recombinant chicken IFN- γ significantly increased host protection against coccidiosis and reduced the intracellular development of *Eimeria* in chickens [64]. In contrast, IL-10 acts as a differentiation factor for a novel subset of T cells that function as suppressors [65]. IL-10 inhibits the synthesis of proinflammatory cytokines (including IL-1 β , TNF- α , and IL-6), thus down-regulating inflammatory T cell responses [66,67]. Following *E. acervulina* infection, IL-10 expression was moderately increased [52]. Finally, IL-15, a cytokine that promotes the survival of T and NK cells [53,68], enhanced protective immunity to coccidiosis challenge when co-administered with the profilin vaccine [7].

In summary, the enhanced protection against experimental avian coccidiosis conferred by QCDC is a result of augmented cellular and humoral immune responses. Future studies to elucidate the molecular and cellular immune mechanisms mediated by QCDC in various avian clinical conditions will be beneficial to the commercial industry.

Acknowledgments

This project was partially supported by the Agricultural Research Service Current Research Information System (ARS CRIS) project 1265-32000-086 and a formal agreement established between ARS and Pfizer Animal Health. QCDC was supplied by the Pfizer Inc. (Kalamazoo, MI). The authors thank Margie Nichols and Stacy Torreyson for their significant contributions to this research.

References

- [1] Lillehoj HS, Lillehoj EP. Avian coccidiosis. A review of acquired intestinal immunity and vaccination strategies. *Avian Dis* 2000;44:408–25.
- [2] Williams RB. Anticoccidial vaccines for broiler chickens: pathways to success. *Avian Pathol* 2002;31:317–53.
- [3] Pogonka T, Klotz C, Kovacs F, Lucius R. A single dose of recombinant *Salmonella typhimurium* induces specific humoral immune responses against heterologous *Eimeria tenella* antigens in chicken. *Int J Parasitol* 2003;33:81–8.
- [4] Williams RB, Carlyle WW, Bond DR, Brown IA. The efficacy and economic benefits of Paracox, a live attenuated anticoccidial vaccine, in commercial trials with standard broiler chickens in the United Kingdom. *Int J Parasitol* 1999;29:341–55.
- [5] Williams RB, Catchpole J. A new protocol for a challenge test to assess the efficacy of live anticoccidial vaccines for chickens. *Vaccine* 2000;18:1178–85.
- [6] Song KD, Lillehoj HS, Choi KD, Yun CH, Parcells MS, Huynh JT, et al. A DNA vaccine encoding a conserved *Eimeria* protein induces protective immunity against live *Eimeria acervulina* challenge. *Vaccine* 2000;19:243–52.
- [7] Min W, Lillehoj HS, Burnside J, Weining KC, Staeheli P, Zhu JJ. Adjuvant effects of IL-1 β , IL-2, IL-8, IL-15, IFN- α , IFN- γ , TGF- β 4 and lymphotactin on DNA vaccination against *Eimeria acervulina*. *Vaccine* 2001;20:267–74.
- [8] Ding X, Lillehoj HS, Dalloul RA, Min W, Sato T, Yasuda A, et al. *In ovo* vaccination with the *Eimeria tenella* EtMIC2 gene induces protective immunity against coccidiosis. *Vaccine* 2005;23:3733–40.
- [9] Du A, Wang S. Efficacy of a DNA vaccine delivered in attenuated *Salmonella typhimurium* against *Eimeria tenella* infection in chickens. *Int J Parasitol* 2005;35:777–85.
- [10] Yarovsky F, Zhang D, Andersen JF, Bannenberg GL, Serhan CN, Hayden MS, et al. TLR11 activation of dendritic cells by a protozoan profilin-like protein. *Science* 2005;308:1626–9.
- [11] Plattner F, Yarovsky F, Romero S, Didry D, Carlier MF, Sher A, et al. *Toxoplasma* profilin is essential for host cell invasion and TLR11-dependent induction of an interleukin-12 response. *Cell Host Microbe* 2008;3:61–3.
- [12] Danforth HD, Augustine PA. *Eimeria tenella*: use of monoclonal antibody in determining the intracellular fate of the refractile body organelles and the effect on *in vitro* development. *Exp Parasitol* 1989;68:1–7.
- [13] Yarovsky F, Kanzler H, Hienny S, Coffman RL, Sher A. Toll-like receptor recognition regulates immunodominance in an antimicrobial CD4⁺ T cell response. *Immunity* 2006;25:525–7.
- [14] Lillehoj HS, Choi KD, Jenkins MC, Vakharia VN, Song KD, Han JY, et al. A recombinant *Eimeria* protein inducing interferon-gamma production: comparison of different gene expression systems and immunization strategies for vaccination against coccidiosis. *Avian Dis* 2000;44:379–89.
- [15] Baldrige JR, Ward JR. Effective adjuvants for the induction of antigen-specific delayed-type hypersensitivity. *Vaccine* 1997;15:395–401.
- [16] Gupta RK, Siber GR. Adjuvants for human vaccines-current status, problems and future prospects. *Vaccine* 1995;13:1263–76.
- [17] Richards RL, Alving CR, Wassef NM. Liposomal subunit vaccines: effects of lipid A and aluminum hydroxide on immunogenicity. *J Pharm Sci* 1996;85:1286–9.
- [18] Laccaille-Dubois M, Wagner H. A review of the biological and pharmacological activities of saponins. *Phytomedicine* 1996;2:363–86.
- [19] Ozel M, Hoglund S, Gelderblom HR, Morein B. Quaternary structure of the immunostimulating complex (ISCOM). *J Ultrastruct Mol Struct Res* 1989;102:240–8.
- [20] Gall D. The adjuvant activity of aliphatic nitrogenous bases. *Immunology* 1966;11:369–86.
- [21] Korsholm KS, Agger EM, Foged C, Christensen D, Dietrich J, Andersen CS, et al. The adjuvant mechanism of cationic dimethyldioctadecylammonium liposomes. *Immunology* 2007;121:216–26.
- [22] Katz D, Inbar I, Samina I, Peleg BA, Heller DE. Comparison of dimethyl dioctadecyl ammonium bromide, Freund's complete adjuvant and mineral oil for induction of humoral antibodies, cellular immunity and resistance to Newcastle disease virus in chickens. *FEMS Immunol Med Microbiol* 1993;7:303–13.
- [23] Lillehoj HS, Lindblad EB, Nichols M. Adjuvant activity of dimethyl dioctadecyl ammonium bromide, complete Freund's adjuvant and *Corynebacterium parvum* with respect to host immune response to coccidial antigens. *Avian Dis* 1993;37:731–40.
- [24] Dominowski PJ, Mannan RM, Krebs RL, Thompson JR, Childers TA, Olsen MK, et al. Novel adjuvant compositions. U.S. Patent application number 20,090,324,641 (<http://www.freepatentsonline.com/y2009/0324641.html>).
- [25] Lee SH, Lillehoj HS, Park DW, Jang SI, Morales A, Garcia D, et al. Induction of passive immunity in broiler chickens against *Eimeria acervulina* by hyperimmune egg yolk immunoglobulin Y. *Poult Sci* 2009;88:562–6.
- [26] Lee SH, Lillehoj HS, Cho SM, Park DW, Hong YH, Chun HK, et al. Immunomodulatory properties of dietary plum on coccidiosis. *Comp Immunol Microb* 2008;31:389–402.
- [27] Lee SH, Lillehoj HS, Park DW, Hong YH, Lin JJ. Effects of *Pediococcus*- and *Saccharomyces*-based probiotic (MitoMax[®]) on coccidiosis in broiler chickens. *Comp Immunol Microb* 2007;30:261–8.
- [28] Lee SH, Lillehoj HS, Park DW, Jang SI, Morales A, Garcia D, et al. Protective effect of hyperimmune egg yolk IgY antibodies against *Eimeria tenella* and *Eimeria maxima* infections. *Vet Parasitol* 2009;163:123–6.
- [29] Lee SH, Lillehoj HS, Dalloul RA, Park DW, Hong YH, Lin JJ. Influence of *Pediococcus*-based probiotic on coccidiosis in broiler chickens. *Poult Sci* 2007;86:63–6.
- [30] Johnson J, Reid WM. Anticoccidial drugs: lesion scoring techniques in battery and floor-pen experiments with chickens. *Exp Parasitol* 1970;28:30–6.
- [31] Lee SH, Lillehoj H, Chun HK, Tuo W, Park H, Cho SM, et al. *In vitro* treatment of chicken peripheral blood lymphocytes, macrophages, and tumor cells with extracts of Korean medicinal plants. *Nutr Res* 2007;27:362–6.
- [32] Lee SH, Lillehoj HS, Cho SM, Chun HK, Park HJ, Lim CI, et al. Immunostimulatory effects of oriental plum (*Prunus salicina* Lindl.). *Comp Immunol Microb* 2009;32:407–17.
- [33] Muller PY, Janovjak H, Miserez AR, Dobbie Z. Processing of gene expression data generated by quantitative real-time RT-PCR. *Biotechniques* 2002;32:1372–9.
- [34] Lee SH, Lillehoj HS, Jang SI, Kim DK, Ionescu C, Bravo D. Effect of dietary curcuma, capsicum, and lentinus on enhancing local immunity against *Eimeria acervulina* infection. *J Poult Sci* 2010;47:89–95.
- [35] Watts AM, Kennedy RC. DNA vaccination strategies against infectious diseases. *Int J Parasitol* 1999;29:1149–63.
- [36] Livingston P. Ganglioside vaccines with emphasis on CM2. *Semin Oncol* 1998;25:638–45.
- [37] Musselli C, Livingston PO, Ragupathi G. Keyhole limpet hemocyanin conjugate vaccines against cancer: the Memorial Sloan Kettering experience. *J Cancer Res Clin* 2001;127:R20–6.
- [38] Waite DC, Jacobson EW, Ennis FA, Edelman R, White B, Kammer R, et al. Three double-blind randomized trials evaluating the safety and tolerance of different formulations of the saponin adjuvant QS-21. *Vaccine* 2001;19:3957–67.
- [39] Gilewski T, Adluri S, Ragupathi G, Zhang S, Yao TJ, Panageas K, et al. Vaccination of high-risk breast cancer patients with mucin-1 (MUC1) keyhole limpet hemocyanin conjugate plus QS-21. *Clin Cancer Res* 2001;6:1693–701.
- [40] Klinguer C, Beck A, De-Lys P, Bussat MC, Blaecke A, Derouet F, et al. Lipophilic quaternary ammonium salt acts as a mucosal adjuvant when co-administered by the nasal route with vaccine antigens. *Vaccine* 2001;19:4236–44.
- [41] Katz D, Kraaijeveld CA, Snippe H. Synthetic lipid compounds as antigen-specific immunostimulators for improving the efficacy of killed-virus vaccines. In: Stewart-Tull DES, editor. The theory and practical application of adjuvants. Chichester: Wiley; 1995. p. 38–50.
- [42] Hilgers LA, Snippe H. DDA as an immunological adjuvant. *Res Immunol* 1992;143:494–503.
- [43] Lindblad EB, Elhay MJ, Silva R, Appelberg R, Andersen P. Adjuvant modulation of immune responses to tuberculosis subunit vaccines. *Infect Immun* 1997;65:623–9.
- [44] Veronesi R, Correa A, Alterio D. Single dose immunisation against tetanus promising results in human trials. *Rev Inst Med Trop Sao Paulo* 1970;12:46–54.

- [45] Stanfield JP, Gall D, Bracken PM. Single-dose antenatal tetanus immunisation. *Lancet* 1973;1:215–9.
- [46] Chambers JD, Thomas CR, Hobbs JR. Induction of specific transplantation tolerance in man by autoblaster immunisation. *Blut* 1980;41:229–36.
- [47] Minke JM, Audonnet JC, Jessett DM, Fischer L, Guigal PM, Coupier H, et al. Canarypox as a vector for influenza and EHV-1 genes: challenges and rewards. In: Proceedings of the second international veterinary vaccines and diagnostic conference, Oxford. 2000. p. 36.
- [48] Minke JM, Fischer L, Baudu P, Guigal PM, Sindle T, Mumford JA, et al. Use of DNA and recombinant canarypox viral (ALVAC) vectors for equine herpes virus vaccination. *Vet Immunol Immunopathol* 2006;111:47–57.
- [49] Lillehoj HS, Okamura M. Host immunity and vaccine development to coccidia and *Salmonella* infections in chickens. *J Poult Sci* 2003;40:151–93.
- [50] Hong YH, Lillehoj HS, Lillehoj EP, Lee SH. Changes in immune-related gene expression and intestinal lymphocyte subpopulations following *Eimeria maxima* infection of chickens. *Vet Immunol Immunopathol* 2006;114:259–72.
- [51] Lee SH, Lillehoj HS, Cho SM, Park DW, Hong YH, Lillehoj EP, et al. Protective effects of dietary safflower (*Carthamus tinctorius*) on experimental coccidiosis. *J Poult Sci* 2008;46:155–62.
- [52] Hong YH, Lillehoj HS, Lee SH, Dalloul RA, Lillehoj EP. Analysis of chicken cytokine and chemokine gene expression following *Eimeria acervulina* and *Eimeria tenella* infections. *Vet Immunol Immunopathol* 2006;114:209–23.
- [53] Lillehoj HS, Min W, Choi KD, Babu US, Burnside J, Miyamoto T, et al. Molecular, cellular, and functional characterization of chicken cytokines homologous to mammalian IL-15 and IL-2. *Vet Immunol Immunopathol* 2001;82:229–44.
- [54] Lillehoj HS, Kim CH, Keeler CL, Zhang S. Immunogenomic approaches to study host immunity to enteric pathogens. *Poult Sci* 2007;86:1491–500.
- [55] Jang SI, Lillehoj HS, Lee SH, Lee KW, Park MS, Bauchan GR, et al. Immunoenhancing effects of Montanide™ ISA oil-based adjuvants on recombinant coccidia antigen vaccination against *Eimeria acervulina* infection. *Vet Parasitol* in press.
- [56] Yao Z, Painter SL, Fanslow WC, Ulrich D, Macduff BM, Spriggs MK, et al. Human IL-17: a novel cytokine derived from T cells. *J Immunol* 1995;155:5483–6.
- [57] Yao Z, Fanslow WC, Seldin MF, Rousseau AM, Painter SL, Comeau MR, et al. Herpesvirus Saimiri encodes a new cytokine, IL-17, which binds to a novel cytokine receptor. *Immunity* 1995;3:811–21.
- [58] Broxmeyer HE. Is interleukin 17, an inducible cytokine that stimulates production of other cytokines, merely a redundant player in a sea of other biomolecules? *J Exp Med* 1996;183:2411–5.
- [59] Fossiez F, Djossou O, Chomarat P, Flores-Romo L, Ait-Yahia S, Maat C, et al. T cell interleukin-17 induces stromal cells to produce proinflammatory and hematopoietic cytokines. *J Exp Med* 1996;183:2593–603.
- [60] Hong YH, Lillehoj HS, Park DW, Lee SH, Han JY, Shin JH, et al. Cloning and functional characterization of chicken interleukin-17D. *Vet Immunol Immunopathol* 2008;126:1–8.
- [61] Min W, Lillehoj HS. Isolation and characterization of chicken interleukin-17 cDNA. *J Interferon Cytokine Res* 2002;22:1123–8.
- [62] Lillehoj HS, Ding X, Quiroz MA, Bevenssee E, Lillehoj EP. Resistance to intestinal coccidiosis following DNA immunization with the cloned 3-1 *Eimeria* gene plus IL-2, IL-15, and IFN- γ . *Avian Dis* 2005;49:112–7.
- [63] Lillehoj HS, Min WG, Dalloul RA. Recent progress on the cytokine regulation of intestinal immune response to *Eimeria*. *Poult Sci* 2004;83:611–23.
- [64] Lillehoj HS, Choi KD. Recombinant chicken interferon-gamma-mediated inhibition of *Eimeria tenella* development *in vitro* and reduction of oocyst production and body weight loss following *Eimeria acervulina* challenge infection. *Avian Dis* 1998;42:307–14.
- [65] Rothwell L, Young JR, Zoorob R, Whittaker CA, Hesketh P, Archer A, et al. Cloning and characterization of chicken IL-10 and its role in the immune response to *Eimeria maxima*. *J Immunol* 2004;173:2675–82.
- [66] Malefyt W, Haanen J, Spits H, Roncarolo MG, Velde A, Figdor C, et al. Interleukin 10 (IL-10) and viral IL-10 strongly reduce antigen-specific human T cell proliferation by diminishing the antigen-presenting capacity of monocytes via downregulation of class II major histocompatibility complex expression. *J Exp Med* 1991;174:915–24.
- [67] Groux H, Powrie F. Regulatory T cells and inflammatory bowel disease. *Immunol Today* 1999;20:442–5.
- [68] Choi KD, Lillehoj HS, Song KD, Han JY. Molecular and functional characterization of chicken IL-15. *Dev Comp Immunol* 1999;23:165–77.