

RESEARCH ARTICLE

Open Access

Genes and pathways for CO₂ fixation in the obligate, chemolithoautotrophic acidophile, *Acidithiobacillus ferrooxidans*, Carbon fixation in *A. ferrooxidans*

Mario Esparza^{1,4}, Juan Pablo Cárdenas¹, Botho Bowien³, Eugenia Jedlicki², David S Holmes^{1*}

Abstract

Background: *Acidithiobacillus ferrooxidans* is chemolithoautotrophic γ -proteobacterium that thrives at extremely low pH (pH 1-2). Although a substantial amount of information is available regarding CO₂ uptake and fixation in a variety of facultative autotrophs, less is known about the processes in obligate autotrophs, especially those living in extremely acidic conditions, prompting the present study.

Results: Four gene clusters (termed *cbb1-4*) in the *A. ferrooxidans* genome are predicted to encode enzymes and structural proteins involved in carbon assimilation via the Calvin-Benson-Bassham (CBB) cycle including form I of ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO, EC 4.1.1.39) and the CO₂-concentrating carboxysomes. RT-PCR experiments demonstrated that each gene cluster is a single transcriptional unit and thus is an operon. Operon *cbb1* is divergently transcribed from a gene, *cbbR*, encoding the LysR-type transcriptional regulator CbbR that has been shown in many organisms to regulate the expression of RubisCO genes. Sigma⁷⁰-like -10 and -35 promoter boxes and potential CbbR-binding sites (T-N₁₁-A/TNA-N₇TNA) were predicted in the upstream regions of the four operons. Electrophoretic mobility shift assays (EMSAs) confirmed that purified CbbR is able to bind to the upstream regions of the *cbb1*, *cbb2* and *cbb3* operons, demonstrating that the predicted CbbR-binding sites are functional *in vitro*. However, CbbR failed to bind the upstream region of the *cbb4* operon that contains *cbbP*, encoding phosphoribulokinase (EC 2.7.1.19). Thus, other factors not present in the assay may be required for binding or the region lacks a functional CbbR-binding site. The *cbb3* operon contains genes predicted to encode anthranilate synthase components I and II, catalyzing the formation of anthranilate and pyruvate from chorismate. This suggests a novel regulatory connection between CO₂ fixation and tryptophan biosynthesis. The presence of a form II RubisCO could promote the ability of *A. ferrooxidans* to fix CO₂ at different concentrations of CO₂.

Conclusions: *A. ferrooxidans* has features of *cbb* gene organization for CO₂-assimilating functions that are characteristic of obligate chemolithoautotrophs and distinguish this group from facultative autotrophs. The most conspicuous difference is a separate operon for the *cbbP* gene. It is hypothesized that this organization may provide greater flexibility in the regulation of expression of genes involved in inorganic carbon assimilation.

Background

Acidithiobacillus ferrooxidans is a mesophilic, obligately chemolithoautotrophic, γ -proteobacterium that gains energy and reducing power from the oxidation of

ferrous iron and reduced inorganic sulfur compounds (RISCs) [1]. It grows optimally at pH 2, although growth as low as pH 1 has been reported [2]. The microorganism is a key player in the solubilization of copper in industrial bioleaching operations and makes an important contribution to the biogeochemical cycling of nutrients and metals in pristine and manmade acidic environments. In such environments, CO₂ would be expected to exist preferentially as a dissolved gas in

* Correspondence: dsholmes2000@yahoo.com

¹Center for Bioinformatics and Genome Biology, MIFAB, Fundación Ciencia para la Vida and Depto. de Ciencias Biológicas, Facultad de Ciencias Biológicas, Universidad Andres Bello, Santiago, Chile
Full list of author information is available at the end of the article

equilibrium with the atmosphere and not in the bicarbonate form typically found at circum-neutral pHs [3].

A. ferrooxidans has previously been shown [4,5] to have candidate genes (*cbbL* and *cbbS*) for the large and small subunits of ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO, EC 4.1.1.39) that catalyses CO₂ fixation by the Calvin-Benson-Bassham (CBB) cycle in many organisms [6]. *cbbL* and *cbbS* are linked to genes predicted to encode carboxysome shell proteins [7] and a divergently transcribed gene encoding the LysR-type transcription regulator CbbR [4]. The intergenic region between *cbbR* and *cbbL* is predicted to harbor binding sites for CbbR [4]. In addition, microarray transcript profiling experiments have detected differential expression of several genes in *A. ferrooxidans* potentially involved in the CBB cycle depending on the growth substrate used [8].

These observations taken together, suggest that, in *A. ferrooxidans*, CbbR can regulate the expression of RubisCO and the carboxysome genes and therefore is likely to be involved in the regulation of carbon fixation as has been observed in other autotrophic bacteria including: *Xanthobacter flavus* [9], *Ralstonia eutropha* H16 [10], *Chromatium vinosum* [11], *Nitrobacter vulgaris* [12], *Halothiobacillus neapolitanus* [13], *Thiobacillus denitrificans* [14], *Rhodobacter sphaeroides* [15], *Rhodobacter capsulatus* [16], *Rhodospirillum rubrum* [17], *Hydrogenovibrio marinus* [18], *Nitrosomonas europaea* [19] and *Thiomicrospira crunogena* XCL-2 [20]. However, no coherent model has been developed for *A. ferrooxidans* to explain all the data and little experimental evidence has been provided to support several of the aforementioned observations, prompting the current investigation.

Methods

Bacterial strains and culture conditions

Information regarding bacterial strains and plasmids used in this study is provided in Table 1. *A. ferrooxidans* was cultured in 9 K medium (adjusted to pH 3.5 with

H₂SO₄) containing 5 g/l elemental sulfur at 30°C under aerobic conditions on a rotary shaker at 150 rpm as described previously [21]. *Escherichia coli* harboring plasmids was grown at 37°C in LB broth with ampicillin (Amp: 100 µg/ml).

General DNA techniques and sequencing of DNA

A. ferrooxidans cultures were centrifuged at 800 × g to remove solid sulfur precipitates prior to cell harvest. Unattached cells were pelleted at 8000 × g for 10 min. The cell pellet was resuspended in 9 K salt solution for washing and washed cells were collected by centrifugation at 8000 × g for 10 min as described previously [21]. Standard procedures [22] were employed to isolate genomic and plasmid DNA from bacteria, to transform plasmid DNA into *E. coli*, and for general DNA handling. Restriction endonucleases and DNA-modifying enzymes were used as recommended by the manufacturers. Plasmid DNA was prepared by means of the QIAprep Spin Mini Kit (Qiagen). Polymerase chain reaction (PCR) products were amplified with Taq DNA polymerase (Fermentas) and purified from agarose gels using the QiaEx DNA Purification Kit (Qiagen). Each PCR reaction contained in a volume of 25 µl 1 ng of template DNA, 0.5 µM of required primers and 0.2 mM of each deoxyribonucleotide in 1× PCR buffer containing 1.5 mM MgCl₂ (Fermentas). PCR conditions were as follows: initial denaturing step at 95°C for 5 min followed by 30 amplification cycles (denaturation at 95°C for 30 s, annealing at the appropriate temperature depending on the specific primer pairs for 20 s, elongation at 72°C) and a final elongation step at 72°C for 10 min. DNA sequencing of pBAD-*cbbR* was carried out by the Göttingen Genomics Laboratory (Göttingen, Germany).

Isolation of RNA and RT-PCR

Total RNA was isolated from cells of *A. ferrooxidans* grown to mid-log phase in 9 K medium supplemented with sulfur, as described previously [23]. The RNA

Table 1 List of bacterial strains and plasmids used in this study

| Strain or plasmid | Relevant characteristic | Source or reference |
|---------------------------------------|--|---------------------|
| Bacterial strains | | |
| <i>Acidithiobacillus ferrooxidans</i> | Type strain | ATCC 23270 |
| <i>E. coli</i> TOP10 | F ⁻ <i>mcrA</i> Δ(<i>mrr-hsdRMS-mcrBC</i>) φ80 <i>lacZ</i> Δ <i>M15</i> Δ <i>lacX74</i> <i>recA1</i> <i>araD139</i> Δ(<i>ara-leu</i>) 7697 <i>galU</i> <i>galk</i> <i>rpsL</i> (Str ^R) <i>endA1</i> <i>nupG</i> | Invitrogen |
| Plasmids | | |
| pBAD-TOPO [®] | Amp ^R promoter <i>araBAD</i> (P _{BAD}) C-terminal: V5 epitope tag-polyhistidine (6 × His) | Invitrogen |
| pBAD- <i>cbbR</i> | pBAD-TOPO::927-bp fragment containing <i>cbbR</i> from <i>A. ferrooxidans</i> ATCC 23270 expressed from P _{BAD} promoter | This study |

Abbreviations used: ATCC, American Type Culture Collection. Amp^R, ampicillin resistance; Str^R, streptomycin resistance.

preparation was treated with DNase I (Fermentas) before proceeding with the cDNA synthesis step. One microgram of total cellular RNA was used for each reaction. Reverse transcription-PCR (RT-PCR) was performed on purified RNA using the One-Step RT-PCR kit (Qiagen). The sequences of the RT and PCR primers used are provided in Table 2. As controls, reactions were carried out that included RNA but lacked reverse transcriptase to assess genomic DNA contamination and that lacked RNA but contained 1 ng of genomic DNA.

Cloning and expression of *cbbR*

A DNA fragment corresponding to the coding region of *cbbR* of *A. ferrooxidans* was amplified by PCR using primers (Integrated DNA Technologies) *cbbRfw* and *cbbRrev* (Table 2). The amplified product was cloned into the expression vector pBAD-TOPO (Invitrogen) according to the manufacturer's instruction. The resulting plasmid pBAD-*cbbR* was introduced by electroporation into *E. coli* TOP10 (Invitrogen) competent cells [22]. *E. coli* was grown at 37°C in 10 ml LB containing 100 µg/ml ampicillin to an OD₆₀₀ of 0.8. Overproduction of the recombinant His₆-tagged CbbR protein was initiated by adding arabinose to a final concentration of 0.1% (w/v) with continued shaking at 200 rpm for 12 h.

Purification of CbbR

Cells from 1.5 l of induced culture were harvested by centrifugation (8,000 × g for 10 min at 4°C) and at -20°C. After thawing the cell pellet was resuspended in 40 ml denaturing buffer containing 6 M guanidine-HCl, 100 mM NaH₂PO₄ and 10 mM Tris-HCl, pH 8.0, and incubated at room temperature with continuous stirring for about 30 min until inclusion body proteins were solubilized. Any remaining insoluble material was removed by centrifugation at 18,000 × g and 7°C for 20 min. The resulting supernatant was filtered through a 0.45-µm membrane and the recombinant protein subsequently purified by affinity chromatography on a 2.5-ml Ni-nitrilotriacetic acid column under amalgam conditions (denaturing conditions-native conditions). Initially the protein was adsorbed to the matrix under denaturing conditions at room temperature after equilibration with binding denaturing buffer (BDB) containing 8 M urea, 100 mM NaH₂PO₄ and 10 mM Tris-HCl, pH 8.0. The column was first washed with BDB and then with washing denaturing buffer (WDB) containing 8 M urea, 100 mM NaH₂PO₄, 10 mM Tris-HCl, pH 6.0. The elution of CbbR was carried out under native conditions at 4°C after equilibrating the column with native buffer (20 mM imidazole, 300 mM NaCl, 50 mM NaH₂PO₄, pH 8.0). His₆-CbbR was eluted at a flow rate of 1 ml/min with eluting native buffer (250 mM imidazole, 300 mM NaCl, 50 mM NaH₂PO₄, pH 8.0). The eluted

fractions were monitored at 280 nm. Fractions with the highest protein content were pooled, dialysed twice against 50 mM HEPES-NaOH, pH 7.8 containing 200 mM KCl, 10 mM MgCl₂, 1 mM dithiothreitol, 0.05 mM phenylmethylsulfonyl fluoride and 50% (w/v) glycerol. The final protein concentration was 4 mg/ml. Protein preparations were analyzed by SDS-polyacrylamide gel electrophoresis in 12% (w/v) polyacrylamide slab gels under reducing conditions in the presence of 100 mM β-mercaptoethanol. Gels were stained with Coomassie Brilliant Blue R-250. Protein contents were determined using the method of Bradford [24], with bovine serum albumin as a standard. CbbR was stored at -20°C.

Production of antisera to CbbR

Multiple intradermal injections were applied to immunize a female Californian giant rabbit (3.0 kg) as described by [25]. A fresh CbbR preparation (0.5 ml; 1 mg/ml) was emulsified in one volume of complete Freund adjuvant (Commonwealth Serum Laboratories, Melbourne, Australia). The emulsion was prepared under aseptic conditions and 1.0 ml was initially injected into four sites on the back of the animal. Booster injections were given in the same way 75 days after the primary immunization, except that incomplete Freund adjuvant was used. The immune response was monitored by Western Blotting assays with serum separated from test blood samples (1.0 to 2.0 ml) that were obtained from an ear vein every 15 to 20 days after each immunization.

Electrophoretic mobility shift assays (EMSA)

DNA fragments containing the four potential *cbb* operon promoter regions were amplified by PCR and simultaneously biotinylated using the biotin 5'-labelled primers (Table 2). DNA-binding assays were performed at 30°C in a final volume of 17 µl containing 12 mM HEPES-NaOH, pH 7.9, 4 mM Tris-HCl, pH 7.9, 1 mM EDTA, 60 mM KCl, 1 mM dithiothreitol, 10% (w/v) glycerol, 5 µg/µl of bovine serum albumin and 2 µg/µl of poly(dI-dC). The indicated amount of CbbR protein (~290 µM) was incubated with the biotin end-labeled target DNA (20 pmol) for 15 min. A 50-fold excess of unlabeled DNA probe was used to challenge the labeled probe. In supershift experiments, a 1:500 dilution of CbbR-specific antiserum was added to the reaction after DNA binding of CbbR and incubated for an additional 15 min. After the binding reactions, samples were loaded onto a low-ionic strength nondenaturing polyacrylamide gel (4.8% [w/v], which had been prerun at a constant current of 200 mA for more than 90 min, and electrophoresed at 150 mA for about 60 min in 0.5× TBE buffer (89 mM Tris base, 89 mM boric acid and 2 mM EDTA). The separated CbbR-DNA complexes were

Table 2 Sequences of primers used in co-transcription RT-PCR assays, cloning experiments and EMSA assays

| Primers used in RT-PCR assays | | | | | |
|---|----------------------------|--------------------------|-------------------------|--------------------------|--------------------------|
| <i>cbb1</i> operon | | | | | |
| Number ^a | Gene | Forward primer (5'-3') | Number ^a | Gene | Reverse primer (5'-3') |
| 1 | <i>cbbR</i> | CAACGCCGTGTGCTCGAA | 2 | <i>cbbL1</i> | CTAGACTTTTTCACGGCCATGCTT |
| 3 | <i>cbbL1</i> | CTGCCAATCGTGTGCGCGC | 4 | <i>csoS2</i> | CGCACGGGAAAGCGACTT |
| 5 | <i>csoS2</i> | CCTATGGTGCCGTGCCAAC | 6 | <i>csoS3</i> | GTGCATGACGCACGCC |
| 7 | <i>csoS3</i> | GTCAGCGGGTCAAAGCCG | 8 | <i>csoS1A</i> | GCCGCTTGGTCATCG |
| 9 | <i>csoS1B</i> | GGAGCAGATGCGTGTGAGCG | 11 | <i>parA</i> | AGTAGAACCCCGCCGAGCCAA |
| 10 | <i>bfrA</i> | CGCGCAGAAGAGTTACAAGCCTTG | 12 | <i>parA</i> | CTGATCGAACCTGAGGATCGG |
| 13 | <i>parA</i> | GTGCTGCGGTTGAAGGGGT | 14 | <i>hyp2</i> | GTGGAGTTCGATAATGGCGGAG |
| 15 | <i>hyp2</i> | CGAGAAGCCTCCGCCATTATC | 16 | <i>cbbQ1</i> | GCCTGTGGGTCTTTTCAGCAT |
| 18 | <i>cbbO1</i> | TGACGCCAGGAAAGCGGTG | 17 | <i>cbbO1</i> | CAGGGATTTTCAGGCTGGTCCG |
| 19 | <i>cbbO1</i> | GCAGAGGCTGCCAGAAAAGCT | 20 | <i>cbbA</i> | AAGCACCTACCGGTATCCGT |
| | | 21 | <i>bioD^a</i> | CAGTGCCACCGCCACCC | |
| <i>cbb2</i> operon | | | | | |
| Forward primer (5'-3') | | | Reverse primer (5'-3') | | |
| 1 | <i>tatC^a</i> | ACGACGGCGTCTAGAACCGCC | 2 | <i>cbbL2</i> | CCGTAATCCTCTAGACCCGCGTT |
| 3 | <i>cbbL2</i> | CATCGAGAAGGAAGGCAAGGC | 4 | <i>cbbS2</i> | CGCAACCTGTTGACGGATCTG |
| 5 | <i>cbbS2</i> | ACCGGAAAACGCCTTCGGC | 6 | <i>cbbQ2</i> | GGTCAATGGGCCATCTCGCC |
| 7 | <i>cbbQ2</i> | AGGGTGTGAGGCGAAGGCC | 8 | <i>cbbO2</i> | GTACGATGGGCGTGTGCGC |
| 9 | <i>cbbO2</i> | GCCTACAGCGAGGAGCCATG | 10 | <i>yfeA^a</i> | GCGGAGCCTTGTCCCTCGG |
| <i>cbb3</i> operon | | | | | |
| Forward primer (5'-3') | | | Reverse primer (5'-3') | | |
| 1 | <i>hyp4^a</i> | TACGAAGGCGGCTCCCCG | 2 | <i>hyp3</i> | CGACGGCAATCGGAGTCTTT |
| 3 | <i>hyp3</i> | CGGGTGATCGCGCTGGAT | 4 | <i>cbbT</i> | CAGAATGCCGTCGTGACCA |
| 5 | <i>cbbT</i> | ATCGGCATCGACCACCTT | 6 | <i>cbbK</i> | TCCATCATAACGAGGACA |
| 7 | <i>cbbK</i> | CCTACATCAGTACGGGTG | 9 | <i>cbbA</i> | CACCTGCTCCAGGTTGTT |
| 8 | <i>pykA</i> | TTGATCCTCATCACCATCGG | 11 | <i>cbbE</i> | GATATGGATATAGTCGGCACCC |
| 10 | <i>cbbA</i> | GCAGGCCAGCAAGATCAA | 14 | <i>trpE</i> | GCCGACAAGGGAGTATCGA |
| 12 | <i>cbbE</i> | CTATCGAACTGGAAGTGGATGG | 16 | <i>trpG</i> | CGATAGCCGCCACGTCG |
| 13 | <i>cbbZ</i> | TCGGCGATTACGTAACG | 17 | <i>trpC^a</i> | AGGGCCACTGCCGCTGC |
| 15 | <i>trpE</i> | GAAACCATGAACAAACGCCG | | | |
| <i>cbb4</i> operon | | | | | |
| Forward primer (5'-3') | | | Reverse primer (5'-3') | | |
| 1 | <i>ompA^a</i> | GGTATTTCCATTTTTGGGGTGGC | 3 | <i>sahA</i> | CGGCAATGCGGACTTTCCTTAC |
| 2 | <i>metK</i> | TTGGGAGCGGACCGACAAG | 5 | <i>metF</i> | AAGCATACTCGGGACCCAAAG |
| 4 | <i>sahA</i> | CTTCGCGGGGTGCTGA | 7 | <i>cbbP</i> | GACGGGATGTTTTTGGACATGG |
| 6 | <i>metF</i> | CACCGAGCCTGCATTTTACACC | 9 | <i>ynbD</i> | GGCTACAGCCACCACGGGAT |
| 8 | <i>cbbP</i> | ATGTTGCCGGGCAGTTTATGTC | 10 | <i>fbnA^a</i> | GCGAGGTGGACTGGACGGA |
| Primers used in EMSA assays and cloning experiments | | | | | |
| Letter designation | Forward primer (5'-3') | | Reverse primer (5'-3') | | |
| (a) Pcbb1 | CGGCAGCGAAGATCTTGAGTTGGTGC | | (b) Pcbb1 | CTCCGGCCTCATACTTTTTTACGG | |
| (c) <i>cbbRfw</i> | TCTATCCGTCATGCAACCTTG | | (d) <i>cbbRrev</i> | GCGCCATTCCTTTTACCACTG | |
| (e) Pcbb2 | ACGACGGCGGCAAGCACC GCC | | (f) Pcbb2 | CCGTAATCCTTTCACCCCGGTT | |
| (g) Pcbb3 | CATTGAACAGGGTCAGCTCCTGG | | (h) Pcbb3 | ATCGGAGTCTTTGATCATGCGCC | |
| (i) Pcbb4 | TTTGGGGTGGCAGCAAGAAGT | | (j) Pcbb4 | GAAACGGATTTCAGAGGTGAAAAG | |

^aGenes lie outside the operons and are not shown in Figure 2.

electrophoretically transferred from the gel onto a nylon membrane in semi-dry blotting apparatus (Biometra, Göttingen). CbbR-DNA binding was detected using a streptavidin-horseradish peroxidase conjugate and a chemiluminescent substrate (Pierce) followed by autoradiography.

Bioinformatic analyses

Metabolic pathways involved in CO₂ assimilation were retrieved from KEGG <http://www.genome.ad.jp/kegg/>. Protein sequences derived from known genes involved in CO₂ assimilation were used as query sequences to search the genome sequence of *A. ferrooxidans* ATCC 23270, using TblastN and BlastP, respectively, with default parameters. When a prospective candidate gene was identified, its predicted protein sequence was then used to formulate a BlastP <http://www.ncbi.nlm.nih.gov> search of the nonredundant database at NCBI. Only bidirectional best hits were accepted as evidence for putative orthologs. Candidate genes and their translated proteins were further characterized employing the following bioinformatic tools: ClustalW [26] for primary structure similarity relations, PSI-PRED [27] for secondary structure predictions, Prosite [28] for motif predictions, ProDom [29] and Pfam [30] for domain predictions. Information regarding the organization of genes in *A. ferrooxidans* was obtained from [2]. Logos were generated using the web-based application available at <http://weblogo.berkeley.edu/logo.cgi>. The height of each letter in bits corresponds to its relative abundance at each position. Promoters of the σ^{70} -type and rho-independent transcriptional stops were predicted for operons *cbb1-4* using the programs BPROM <http://www.softberry.com> and Transterm [31], respectively.

The organization of gene clusters in facultative and obligate autotrophs involved in the CBB cycle was derived from information available in IMG-JGI <http://www.jgi.doe.gov/> and MicrobesOnline <http://www.microbesonline.org/>, with additional information added for *H. marinus* [18] and *A. ferrooxidans*, *Acidithiobacillus caldus* and *Acidithiobacillus thiooxidans* (this study). The phylogenetic cladogram of these bacteria was constructed from 16 S rRNA sequences obtained from KEGG Orthology K01977 <http://www.genome.jp/kegg/ko.html> and from GenBank <http://www.ncbi.nlm.nih.gov/> for *A. caldus* (GI454888), *A. thiooxidans* (GI454888) and *H. marinus* (GI3882094). 16 S rRNA alignments were carried out using ClustalW and the cladogram was constructed by the NJ method using the program MEGA 4.0 [32]. The robustness of the tree was evaluated by bootstrapping using 1000 replicas. The tree was rooted using the 16 S rRNA of the ϵ -proteobacterium *Helicobacter pylori*.

Results, Discussion and Conclusions

The genome of *A. ferrooxidans* ATCC 23270 encodes CbbR, a LysR-type transcription factor

A gene *cbbR* was predicted in the genome of *A. ferrooxidans* ATCC 23270 (type strain) that potentially encodes a protein with significant amino acid sequence similarity and domain structure to other well-documented CbbRs of the LysR family of transcription factors (Additional file 1). *cbbR* is divergently transcribed from *cbbL1*, a gene predicted to encode the large subunit of form I RubisCO. The genetic linkage between *cbbR* and *cbbL1* is known to be conserved in a number of autotrophic bacteria that fix CO₂ via the CBB cycle such as *Acidithiobacillus ferrooxidans* Fe1 [4], *Hydrogenophilus thermoluteolus* [33], *Nitrosomonas europaea* [19], *Rhodobacter sphaeroides* [34], *Rhodobacter capsulatus* [35], *R. eutropha* H16 [36], *Rhodospirillum rubrum* [17], *Thiobacillus denitrificans* [14] and *Xanthobacter flavus* [9]. We here extend this list to include: *Alkalilimnicola ehrlichii*, *Halorhodospira halophila*, *Methylobium petroleiphilum*, *Nitrobacter winogradskyi*, *Nitrosococcus oceani*, *Nitrospira multiformis*, *Thiomicrospira crunogena* and *Xanthobacter autotrophicus* (Additional file 2).

The *cbbR-cbbL1* intergenic region of *A. ferrooxidans* strain Fe1 has been shown to contain divergent σ^{70} -type promoters and to exhibit two CbbR binding sites that partially overlap these promoters ([4], Figure 1A). The binding sites conform to the pseudo-palindromic motif TNA-N₇-TNA [13] that is a subset of the consensus LysR-type transcription factor binding site T-N₁₁-A [37]. Logos were derived from a multigenome comparison of the *cbbR-cbbL1* intergenic region of a number of bacteria (Additional file 3) and were aligned with the CbbR sites of *A. ferrooxidans* strain Fe1, allowing the prediction of the CbbR binding sites of *A. ferrooxidans* ATCC 27230 (Figure 1B and 1C).

Organization and expression of gene clusters predicted to be involved in CO₂ fixation and associated pathways of central carbon metabolism

A cluster of 16 genes, termed *cbb1*, was predicted to be involved CO₂ fixation. RT-PCR experiments showed that *cbb1* is transcribed as a single unit and thus can be considered to be an operon (Figure 2A). Operon *cbb1* consists of *cbbL1* and *cbbS1*, potentially encoding the large and small subunits of form IAc RubisCO, seven *cso* genes predicted to be involved in α -carboxysome formation, two genes (*cbbQ1* and *cbbO1*) presumed to be involved in RubisCO activation and *cbbA*, potentially encoding a fructose-1,6-bisphosphate aldolase. Gene descriptions are provided in Table 3.

Three additional gene clusters termed *cbb2* (four genes), *cbb3* (twelve genes) and *cbb4* (five genes) were




Figure 1 The *cbbR-cbbL1* intergenic regions of *A. ferrooxidans* strains Fe1 and ATCC 23270. (A) DNA sequence of *cbbR-cbbL1* intergenic region of *A. ferrooxidans* Fe1 showing two TNA-N₇-TNA CbbR-binding regions (boxed sequences) and experimentally verified nucleotides protected by CbbR binding (*) and σ^{70} promoter regions (-10 and -35 sites) (Modified from [5], with permission of the publisher). (B) Logos derived from multiple sequence alignments of the *cbbR-cbbL1* intergenic region of eight bacteria showing conservation of the CbbR-binding sites (more information in additional file 3). (C) Prediction of CbbR-binding sites and σ^{70} promoter regions in the *cbbR-cbbL1* intergenic region of *A. ferrooxidans* ATCC 23270 by comparison with experimentally verified regions of *A. ferrooxidans* Fe1 and using the information derived from Logos.

identified that are predicted to encode functions related to CO₂ fixation and central carbon metabolism (Table 3). RT-PCR experiments revealed that gene clusters *cbb2*, *cbb3* and *cbb4* are transcribed as single units, respectively, and thus constitute operons (Figure 2B-D). *cbb2* contains genes (*cbbL2* and *cbbS2*) encoding additional copies of the large and small subunit of form IAq RubisCO and associated RubisCO activation genes (*cbbQ2* and *cbbO2*) (Figure 2B). The deduced amino acid sequences of these genes are similar but not identical to the corresponding proteins encoded in the *cbb1* operon; CbbL1 and CbbL2 exhibit 84% amino acid sequence identity, whereas CbbS1 and CbbS2 share 56% identity and CbbQ1 and CbbO1 have 84% and 59% identity with CbbQ2 and CbbO2, respectively.

Genes predicted to be encoded by operons *cbb3* and *cbb4* are listed in Table 3 and their organization within these operons is shown in Figure 2.

The two enzymes that are unique to the CBB cycle are RubisCO (encoded by operons *cbb1* and *cbb2*) and phosphoribulokinase (encoded by operon *cbb4*). RuBisCO catalyzes the first step of the cycle, the carboxylation of ribulose-1,5-bisphosphate (RuBP) with CO₂. Phosphoribulokinase catalyzes the last step of the cycle which is the regeneration of the CO₂ acceptor molecule, RuBP, by phosphorylation of ribulose 5-phosphate with ATP. Other steps of the cycle, encoded in operon *cbb3*, are catalyzed by enzymes common to glycolytic and gluconeogenic pathways in central carbon metabolism [8,36].

Promoters of the σ^{70} -type and rho-independent transcriptional stops were predicted for operons *cbb1-4*


(Figure 2). In addition, potential CbbR-binding sites were identified in the four operons based on the detection of conserved TNA-N₇-TNA and T-N₁₁-A motifs described above for operon *cbb1* (Figure 2).

CbbR binds *in vitro* to the predicted σ^{70} -like promoter regions of operons *cbb1-4*

Binding of CbbR to DNA fragments containing the predicted promoters of the four operons *cbb1-4* was evaluated *in vitro* by electrophoretic mobility shift assays (EMSAs). For this purpose the *cbbR* gene was cloned and expressed in *E. coli*. Purified CbbR was used to prepare antisera (anti-CbbR antibodies) whose activity was checked by Western blotting against purified CbbR (data not shown). Biotin-labeled promoter DNA for the EMSA assays was prepared by PCR using primers specified in Table 2 and whose locations within the four operons are shown in Figure 2.

Results show that CbbR was able to retard the promoter regions of the *cbb1*, *cbb2* and *cbb3* operons but not the *cbb4* operon (Figure 3). When a 50-fold molar excess of unlabelled fragment was included in the binding assay retardation of the labelled fragments was abolished. Furthermore, the addition of anti-CbbR antibodies to the reaction produced a supershift in migration, indicating that the shift was caused specifically by the binding of CbbR.

Binding of CbbR to the predicted promoter regions of operons *cbb1-3* suggests that it is involved in their regulation. The reason for the failure of CbbR to retard the DNA fragment containing the predicted promoter of the *cbb4* operon is not known. Perhaps this fragment



requires the presence of additional factors for CbbR binding that are not present in the *in vitro* cocktail used for the EMSA analysis. Alternatively, the predicted CbbR binding site is not functional.

Gene organization of the *cbb* operons

The *cbb3* operon includes not only genes involved in carbon assimilation but also harbors genes with similarity to *trpE* and *trpG* that are predicted to encode the components I and II of anthranilate synthase, the first enzyme of the tryptophan biosynthesis pathway. Anthranilate synthase catalyzes the conversion of chorismate to

anthranilate with the concomitant release of pyruvate [38,39]. In some cases, this conversion can be accomplished by TrpE alone [40].

In order to determine if the association between *trpEG* and the *cbb* genes is restricted to *A. ferrooxidans*, an examination of gene organization was carried out in all sequenced genomes of facultative and obligate autotrophic proteobacteria. Twenty-six proteobacterial organisms (11 α -, 7 β - and 8 γ -) were analyzed, including 10 obligate autotrophs. Linkage between *trpE/G* and *cbbE* and/or *cbbZ* was found in all sequenced obligate autotrophs, all of which belong to the β - or

Table 3 Predicted genes of *cbb* operons

| *Accession | ^a Gene name | ^b Predicted function | ^c Best BlastP hit | ^d % Similarity | ^e Score | ^f E-value | ^g Domains and motifs |
|---------------------------|------------------------|---|---------------------------------------|---------------------------|--------------------|----------------------|--|
| Operon <i>cbb1</i> | | | | | | | |
| ACK78724.1 | <i>cbbR</i> | LysR family transcriptional regulatory protein CbbR | <i>Nitrococcus mobilis</i> | 76 | 363 | 7e-99 | PD462572, PD756396, Pfam03466, Pfam00126, COG0583 |
| ACK79627.1 | <i>cbbL1</i> | Ribulose biphosphate carboxylase large subunit 1 [4.1.1.39] | <i>Halothiobacillus neapolitanus</i> | 94 | 882 | 0 | PD417314, PD000044, Pfam00016, Pfam02788, COG1850 |
| ACK77836.1 | <i>cbbS1</i> | Ribulose biphosphate carboxylase small subunit 1 [4.1.1.39] | <i>Methylococcus capsulatus</i> | 80 | 161 | 8e-39 | PD000290, Pfam00101, COG4451 |
| ACK78689.1 | <i>csoS2</i> | Carboxysome structural peptide | <i>Thiobacillus denitrificans</i> | 59 | 325 | 9e-87 | PD579361, tat signal peptide |
| ACK80925.1 | <i>csoS3</i> | Carboxysome structural peptide | <i>Thiobacillus denitrificans</i> | 65 | 537 | 5e-151 | PD191834, Pfam08936 |
| ACK80352.1 | <i>csoS4A</i> | Carboxysome peptide A | <i>Thiobacillus denitrificans</i> | 93 | 139 | 6e-32 | PD012510, Pfam03319, COG4576, tat signal peptide |
| ACK79436.1 | <i>csoS4B</i> | Carboxysome peptide B | <i>Thiobacillus denitrificans</i> | 82 | 119 | 7e-26 | PD012510, Pfam03319, COG4576 |
| ACK78722.1 | <i>csoS1C</i> | Microcompartments protein | <i>Nitrosomonas eutropha</i> | 97 | 142 | 6e-33 | PD003442, Pfam00936, COG4577 |
| ACK79154.1 | <i>csoS1A</i> | Microcompartments protein | <i>Nitrosomonas eutropha</i> | 97 | 144 | 1e-33 | PD003442, Pfam00936, COG4577 |
| ACK79584.1 | <i>csoS1B</i> | Microcompartments protein | <i>Nitrosomonas eutropha</i> | 95 | 146 | 3e-34 | PD003442, Pfam00936, COG4577 |
| ACK79096.1 | <i>bfrA</i> | Bacterioferritin | <i>Thiobacillus denitrificans</i> | 70 | 135 | 6e-31 | PDA00179, Pfam00210, COG1633 |
| ACK77923.1 | <i>hyp1</i> | Hypothetical protein | <i>Thiobacillus denitrificans</i> | 81 | 68 | 2e-10 | PDA1E0I5 |
| ACK80576.1 | <i>parA</i> | Partition protein A | <i>Thiobacillus denitrificans</i> | 72 | 196 | 6e-49 | PD194671, Pfam01656, COG1192 |
| ACK78664.1 | <i>hyp2</i> | Hypothetical protein | <i>Acidithiobacillus ferrooxidans</i> | 100 | 156 | 1e-09 | |
| ACK80060.1 | <i>cbbQ1</i> | Rubisco activation protein | <i>Nitrosomonas europaea</i> | 92 | 489 | 5e-137 | PD490543, Pfam08406, Pfam07728, COG0714, COG5271 |
| ACK80817.1 | <i>cbbO1</i> | Rubisco activation protein | <i>Thiobacillus denitrificans</i> | 74 | 940 | 0 | PD140693, PD679436, Pfam00092, COG4867, COG4548 |
| ACK80290.1 | <i>cbbA</i> | Fructose-biphosphate aldolase [4.1.2.13] | <i>Bradyrhizobium</i> sp. | 61 | 295 | 3e-78 | PD002376, PD030418, Pfam01116, Pfam07876, COG191 |
| Operon <i>cbb2</i> | | | | | | | |
| ACK80366.1 | <i>cbbL2</i> | Ribulose biphosphate carboxylase/oxygenase large subunit 2 [4.1.1.39] | <i>Thiobacillus denitrificans</i> | 97 | 920 | 0 | PD417314, PD000044, Pfam00016, Pfam02788, COG1850 |
| ACK79774.1 | <i>cbbS2</i> | Ribulose biphosphate carboxylase/oxygenase small subunit 2 [4.1.1.39] | <i>Thiobacillus denitrificans</i> | 88 | 203 | 3e-51 | PD000290, Pfam00101, COG4451 |
| ACK80953.1 | <i>cbbQ2</i> | Rubisco activation protein | <i>Nitrosomonas europaea</i> | 92 | 483 | 6e-135 | PD490543, PD372819; Pfam08406, Pfam07728, COG0714 |
| ACK78928.1 | <i>cbbO2</i> | Rubisco activation protein | <i>Thiobacillus denitrificans</i> | 76 | 965 | 0 | PD140693, PD025507, COG4548 |
| Operon <i>cbb3</i> | | | | | | | |
| ACK80740.1 | <i>hyp3</i> | Hypothetical protein | <i>Thiobacillus denitrificans</i> | 49 | 149 | 8e-9 | PD796582 |
| ACK78212.1 | <i>suhB</i> | Inositol-phosphate phosphatase [3.1.3.25] | <i>Methylococcus capsulatus</i> | 66 | 646 | 8e-66 | PD001491, PD013702, pfam00459, pfam00316, COG0483, COG1218 |
| ACK80404.1 | <i>cbbF</i> | Fructose-1,6-bisphosphatase [3.1.3.11] | <i>Mariprofundus ferrooxydans</i> | 71 | 823 | 3e-86 | PD007014, PD863173, pfam03320, COG1494 |
| ACK79091.1 | <i>cbbT</i> | Transketolase [2.2.1.1] | <i>Methylococcus capsulatus</i> | 75 | 2264 | 0.0 | PD308336, pfam00456, pfam02779, COG3959, COG0021 |
| ACK78716.1 | <i>cbbG</i> | Glyceraldehyde-3-phosphate dehydrogenase type I [1.2.1.-] | <i>Burkholderia thailandensis</i> | 82 | 1189 | 1e-128 | PD959395, PD859695, pfam02800, pfam00044, COG0057 |

Table 3 Predicted genes of *cbb* operons (Continued)

| | | | | | | | |
|---------------------------|-------------|---|--------------------------------------|----|------|--------|--|
| ACK79414.1 | <i>cbbK</i> | Phosphoglycerate kinase [2.7.2.3] | <i>Alcanivorax borkumensis</i> | 80 | 1296 | 6e-141 | PD000619, PDA014E1, pfam00162, COG0126 |
| ACK78522.1 | <i>pykA</i> | Pyruvate kinase II [2.7.1.40] | <i>Thiobacillus denitrificans</i> | 79 | 1491 | 2e-163 | PD983049, PD745602, pfam00224, pfam02887, COG0469 |
| ACK79923.1 | <i>cbbA</i> | Fructose-bisphosphate aldolase [4.1.2.13] | <i>Nitrosococcus oceani</i> | 90 | 1474 | 1e-161 | PD875785, PD002376, pfam01116, COG0191 |
| ACK80630.1 | <i>cbbE</i> | Ribulose-5-phosphate 3-epimerase [5.1.3.1] | <i>Herminiimonas arsenicoxydans</i> | 80 | 753 | 2e-78 | PD003683, PD591639, pfam00834, COG0036 |
| ACK80633.1 | <i>cbbZ</i> | Phosphoglycolate phosphatase [3.1.3.18] | <i>Thiobacillus denitrificans</i> | 64 | 484 | 4e-47 | PD946755, PDA11895, pfam00702, COG0546, COG0637 |
| ACK78314.1 | <i>trpE</i> | Anthranilate synthase component I [4.1.3.27] | <i>Methylococcus capsulatus</i> | 77 | 1569 | 2e-172 | PD005777, PD105823, pfam00425, pfam04715, COG0147, COG1169 |
| ACK78895.1 | <i>trpG</i> | Anthranilate synthase component II [4.1.3.27] | <i>Nitrosomonas europaea</i> | 86 | 770 | 2e-80 | PD806135, PD976090, pfam00117, pfam07722, COG0512, COG0518 |
| Operon <i>cbb4</i> | | | | | | | |
| ACK79981.1 | <i>metK</i> | S-adenosylmethionine synthetase [2.5.1.6] | <i>Ralstonia eutropha</i> | 86 | 591 | 2e-167 | PD499406, PD606972, pfam02773, pfam02772, COG0192 |
| ACK78713.1 | <i>sahA</i> | S-adenosyl-L-homocysteine hydrolase [3.3.1.1] | <i>Pseudomonas stutzeri</i> | 88 | 748 | 0 | PD730548, PD551162, pfam05221, pfam00670, COG0499 |
| ACK78001.1 | <i>metF</i> | 5,10-methylenetetrahydrofolate reductase [1.7.99.5] | <i>Methylococcus capsulatus</i> | 69 | 306 | 1e-81 | PD756524, PD763008, pfam02219, COG0685 |
| ACK78673.1 | <i>cbbP</i> | Phosphoribulokinase [2.7.1.19] | <i>Nitrosococcus oceani</i> | 78 | 402 | 2e-110 | PD739884, PD015803, pfam00485, COG3954 |
| ACK79243.1 | <i>ynbD</i> | Phosphosterase, PA-phosphatase | <i>Polaromonas naphthalenivorans</i> | 81 | 759 | 1e-81 | PD589889, pfam 01569, COG0474, CD03386, CD00127 |

* The sequence and annotation of the complete *A. ferrooxidans* strain ATCC 23270 genome is available at the Comprehensive Microbial Resource (CMR) (J. Craig Venter Institute, <http://www.jcvi.org>) and in GenBank/EMBL/DBJ accession number CP001219.

^a Proposed gene name.

^b Proposed enzyme activity with EC number if available


^c Organism with the best BlastP hit to the candidate gene.

^d Percentage of similarity (% Sim) of candidate gene to that found in the organism listed in row (c).

^e Score of BlastP match.

^f E value of BlastP match.

^g Motif and domains identified in the candidate proteins: CD, Conserved Domains; COG, Clusters of Orthologous Groups of Proteins; Pfam, protein families; PD, Prodom (protein domains); PS, Prosite tat signal peptide



γ -proteobacteria divisions (Figure 4, Table 4), whereas only 4 out of 14 facultative heterotrophs were detected with this clustering. These four exceptions are found only in the β - or γ -proteobacteria and none in the α -proteobacterial division (Figure 4, Table 4). This suggests a previously unreported linkage between genes encoding CBB cycle associated enzymes and *trpEG* or *trpE* that is most conserved in obligate autotrophs of the β - and γ -proteobacteria.

We hypothesize that in *A. ferrooxidans* production of pyruvate via anthranilate synthase activity provides a novel network connection between the CBB cycle on the one hand and general central carbon metabolism including the incomplete ("horseshoe"-like) TCA [2] on the other hand. Consistent with this idea is the presence of a predicted *pykA* upstream of *trpEG* in the *cbb3* operon. *PykA* is predicted to encode pyruvate kinase that catalyzes the conversion of phosphoenol pyruvate (PEP) to pyruvate. In addition to supplying pyruvate, *PykA* could also reduce the level of intracellular PEP. PEP has been shown to be a ligand of *CbbR* in *Ralstonia eutropha* H16, promoting its binding to target DNA sites and consequently effecting the regulation of *cbb* genes [40]. If PEP carries out a similar function in *A. ferrooxidans*, the depletion of PEP via *PykA* activity could provide a means for feedback control of operons that are regulated by *CbbR*, including the auto-regulation of operon *cbb3*.

The organization of *cbb* genes in *A. ferrooxidans* exhibits similarities with obligate autotrophs that distinguish this group from facultative autotrophs. For example, *A. ferrooxidans*, contains three or more gene clusters dedicated to carbon assimilation. This is similar to other obligate autotrophic γ -proteobacteria including *A. caldus*, *A. thiooxidans*, *Hydrogenovibrio marinus*, *Nitrosococcus oceani* and *Thiomicrospira crunogena*, and obligate autotrophic β -proteobacteria such as *Nitrosomonas europaea*, *Nitrosomonas eutropha*, and *Nitrospira multiformis* and *Thiobacillus denitrificans*. This contrasts with facultative autotrophs that contain only one or two *cbb* clusters (Figure 4, Table 4), with some exceptions, e.g. the α -proteobacteria *Bradyrhizobium sp.*, *N. hamburgensis*, *N. winogradski*, *R. sphaeroides* and *R. palustris* and the β -proteobacterium *R. eutropha*, which contain unique, but duplicated, *cbb* clusters). Multiple *cbb* clusters could provide obligate autotrophs with a greater flexibility in regulating CO_2 fixation compared to facultative autotrophs. For example, this flexibility may be necessary to adjust carbon assimilation in response to changing environmental concentrations of CO_2 [18], whereas facultative autotrophs might be able to circumvent this need by exploiting organic carbon sources in times of low CO_2 concentrations.

Another characteristic of *cbb* gene organization in *A. ferrooxidans* is the lack of linkage of the phosphoribulokinase gene, *cbbP*, with other *cbb* genes (Figure 4, Table 4) as has previously been reported for the deep-sea vent obligate chemolithoautotroph *T. crunogena* XCL-2 and for several other obligate autotrophs [20,41]; we now extend this list to include *A. ferrooxidans* ATCC 23270 and ATCC 53993, *A. caldus*, *A. thiooxidans* *H. marinus*, *N. europaea* and *Thiomicrospira crunogena* (Figure 4, Table 4). In contrast, in all sequenced facultative autotrophs *cbbP* is associated with other *cbb* genes (Figure 4, Table 4).

In obligate autotrophs, the contextual disconnection of *cbbP* from *cbbLS* could provide greater flexibility for CO_2 fixation by allowing *RubisCO* to be differentially expressed according to environmental and/or metabolic requirements without simultaneously expressing the remaining CBB cycle genes, many of which carry out functions in addition to carbon fixation. This is in sharp contrast to the organization found in most facultative autotrophs, where *cbbP* is usually juxtaposed to *cbbLS* and other genes of the CBB cycle facilitating their coordinate repression during heterotrophic growth [13,20,34,36,41].

Model for predicted enzymes and pathways involved in CO_2 fixation

A model is proposed for C_1 fixation in *A. ferrooxidans* based on the predicted roles of the genes encoded in the *cbb* operons (Figure 5). In contrast to most facultative autotrophs, the main focus of regulation of the CBB cycle in *A. ferrooxidans* may be the CO_2 fixation reaction itself catalyzed by *RubisCO*, rather than at the level of the other CBB cycle enzymes. This hypothesis is supported by the observation that the genes encoding *RubisCO* and *RubisCo* accessory proteins, are clustered in operons that do not contain *cbbP* nor *cbb* that encode the main CBB enzymes. *cbbP* is also separated from the rest of the *cbb* genes in the *cbb4* operon, with an apparent absence of *CbbR* binding to its promoter. We suggest that the promoters for the *cbb1*, *cbb2* and *cbb3* operons have different affinities for *CbbR* and may thus exhibit different regulation patterns, possibly associated with the environmental availability of CO_2 . The *cbb1* operon, containing *cbbLS-cso*, is predicted to serve at low CO_2 concentrations because carboxysomes have been shown to improve *RubisCO* catalytic efficiency by concentrating CO_2 [6,13]. In contrast, the *cbb2* operon, containing *cbbLSQO*, is predicted to be used when higher concentrations of CO_2 are available since carboxysome synthesis is energetically and materially expensive [18].

The *cbb3* operon, containing genes for most CBB cycle enzymes and pyruvate kinase, is proposed to be




Figure 4 Organization of gene clusters involved in the CBB cycle of facultative and obligate autotrophic α -, β - and γ -proteobacteria presented as a phylogenetic cladogram based on 16 S RNA. Numbers refer to bootstrapping results from 1000 trees. Organism names are provided in the text. The asterisk indicates that the respective organism is an obligate autotroph.

Table 4 Characteristics of *cbb* gene clusters in facultative and obligate, autotrophic bacteria


| Organism | Autotrophy status | Phylogenetic classification -proteo-bacteria | No. copies <i>cbbR</i> | Presence of <i>cso</i> genes? | <i>trpE/G</i> associated with <i>cbb</i> ? | <i>cbb</i> gene cluster associated with <i>cbbP</i> ? | No. <i>cbb</i> gene clusters |
|---|-----------------------|--|------------------------|-------------------------------|--|---|------------------------------|
| <i>Acidithiobacillus ferrooxidans</i> ATCC 23270 and ATCC 53993 | obligate | Gamma- | 2 | Yes | Yes | No | 5* |
| <i>Acidithiobacillus thiooxidans</i> ATCC 19377 | obligate | Gamma- | 2 | Yes | Yes | No | 5 |
| <i>Acidithiobacillus caldus</i> ATCC 51756 | obligate | Gamma- | 2 | Yes | Yes | No | 5 |
| <i>Nitrosomonas europaea</i> ATCC 19718 | obligate | Beta- | 1 | No | Yes | No | 4 |
| <i>Nitrosomonas eutropha</i> C71 | obligate | Beta- | 1 | Yes | Yes | No | 4 |
| <i>Nitrosococcus oceani</i> ATCC 19707 | obligate | Beta- | 1 | No | Yes | No | 4 |
| <i>Thiomicrospira crunogena</i> XCL-2 | obligate | Gamma- | 3 | Yes | Yes | No | 5 |
| ⁵ <i>Hydrogenovibrio marinus</i> MH-110 | obligate | Gamma- | 2 | Yes | N/D | N/D | 3 |
| <i>Thiobacillus denitrificans</i> ATCC 25259 | obligate | Beta- | 2 | Yes | Yes | No | 5 |
| <i>Nitrosospira multiformis</i> ATCC 25196 | obligate | Beta- | 1 | No | Yes | No | 4 |
| <i>Methylococcus capsulatus</i> Bath | obligate methanotroph | Gamma- | 1 | No | Yes | Yes | 3 |
| ¹ <i>Nitrobacter hamburgensis</i> X14 | facultative | Alpha- | 3 | Yes | No | Yes | 3 |
| <i>Nitrobacter winogradskyi</i> Nb-255 | facultative | Alpha- | 3 | Yes | No | Yes | 3 |
| <i>Halorhodospira halophila</i> SL1 | facultative | Gamma- | 1 | No | Yes ³ | Yes | 2 |
| <i>Alkalilimnicola ehrlichii</i> MLHE-1 | facultative | Gamma- | 1 | No | Yes ³ | Yes | 2 |
| <i>Bradyrhizobium</i> sp. BTAi1 | facultative | Alpha- | 2 | Yes | No | Yes | 3 |
| <i>Bradyrhizobium japonicum</i> USDA 110 | facultative | Alpha- | 1 | No | No | Yes | 1 |
| <i>Ralstonia eutropha</i> H16 | facultative | Beta- | 1 | No | No | Yes | 2 ⁴ |
| <i>Dechloromonas aromatica</i> RCB | facultative | Alpha- | 1 | No | No | Yes | 2 |
| ² <i>Magnetospirillum magneticum</i> AMB-1 | facultative | Alpha- | ? | No | No | Yes | 2 |
| <i>Paracoccus denitrificans</i> PD1222 | facultative | Alpha- | 1 | No | No | Yes | 1 |
| <i>Rhodobacter sphaeroides</i> 2.4.1 | facultative | Alpha- | 1 | No | No | Yes | 2 |
| <i>Rhodoferrax ferrireducens</i> T118 | facultative | Beta- | 1 | No | No | Yes | 1 |
| <i>Rhodopseudomonas palustris</i> CGA009 | facultative | Alpha- | 2 | No | No | Yes | 3 |
| <i>Rhodospirillum rubrum</i> ATCC 11170 | facultative | Alpha- | 1 | No | No | Yes | 1 |
| <i>Sinorhizobium meliloti</i> 1021 | facultative | Alpha- | 1 | No | No | Yes | 1 |

*in addition to the four *cbb* operons described in this paper, a fifth gene cluster containing *cbb* genes (including a form II RubisCO gene) has recently been detected in *A. ferrooxidans* (43). ¹Two copies of *cbbR* and two *cbb* gene clusters are present on two plasmids; ²two highly similar operons present in the genome; ³in these organisms, *trpE* gene is neighbor to *cbbP* but not *cbbE*. ⁴*R. eutropha* H16 possesses a duplicated *cbb* operon, with similar copies in the chromosome and in a megaplasmid. ⁵Data derived from cloned sequences (18). N/D = no data.

responsible for connecting CO₂ fixation with the rest of central carbon metabolism. Except for *cbbG* and *cbbK* encoding glyceraldehyde-3-phosphate dehydrogenase, type I and phosphoglycerate kinase respectively, genes of the *cbb3* operon have duplicated copies in the genome (data not shown), potentially allowing regulation of the CBB cycle independently of the remaining pathways of central carbon metabolism. For example, some CBB cycle intermediates also form part of gluconeogenesis and glycolysis resulting in the production of pyruvate that is channeled, via the pyruvate dehydrogenase complex, into the incomplete TCA "horseshoe" where the flux of intermediates serves for amino acid biosynthesis (e.g. glutamate). The pyruvate dehydrogenase also

provides acetyl-CoA used in fatty acid biosynthesis. In addition, the presence of *cbbZ* in the *cbb3* operon is associated with phosphoglycolate phosphatase activity, responsible for removal of phosphoglycolate, an undesirable product of the oxygenase activity of RubisCO, that must be detoxified preferentially by rechanneling to 3-phosphoglycerate [13,36].

The co-transcriptional connection between the *cbb*, *pykA* and *trpEG* genes in the *cbb3* operon may reflect the substrate requirement of anthranilate phosphoribosyltransferase for an activated pentose (5-phosphoribosyl 1-pyrophosphate) in order to proceed to the next step of tryptophan biosynthesis [42]. The production of the activated pentose would be stimulated by the activity of



the operon. An alternate hypothesis is that the co-transcriptional connection represents a means for pyruvate regeneration since both *pykA* and *trpE/G* produce pyruvate.

In addition to the four *cbb* operons described herein, a fifth gene cluster has recently been detected in *A. ferrooxidans* that includes genes *cbbM*, *cbbQ3* and *cbbO3* predicted to encode form II of RubisCO and its associated chaperons, respectively [43]. The cluster also contains another putative *cbbR* divergently transcribed from *cbbMQO*. Future work will evaluate the role of this cluster in CO₂ fixation.

Additional material

Additional file 1: Prediction of secondary structure elements in CbbR of *Acidithiobacillus ferrooxidans*. Above: secondary structure predictions of alpha-helix, beta-sheet, HTH DNA binding domain, oligomerization domain and LysR-substrate like domain. Below: alignment of amino acid sequences from the HTH domain from several bacteria (abbreviations used can be found in Additional File 2) with the pfam domain00126.

Additional file 2: Alignment and conservation of DNA sequences in the intergenic regions between *cbbR* and *cbbl1* in autotrophic bacteria. The DNA sequences contain the *cbb* control elements including the operator, the operon promoter (*pcbbL*) and the promoter *cbbR* (*pcbbR*). The CbbR regulator binds to region R (recognition site) and the region A (activation site) of the *cbb* operator. The nucleotides conserved (TNA-N_{7/8}-TNA, T-N₁₁-A) for to bind CbbR are located in intergenic regions RI-1, RI-2 and RI-3. The prediction of the promoter and the sites for to bind $\sigma 70$ are in the columns (sequences -35 and -10). The names of bacteria are: *Acidithiobacillus ferrooxidans* (Af), *Hydrogenophilus thermoluteolus* (Ht), *Xanthobacter flavus* (Xf), *Nitrosomonas europaea* (Ne), *Rhodobacter capsulatus* (Rc), *Rhodobacter sphaeroides* (Rs), *Ralstonia eutropha* H16 (Ral), *Ralstonia metallidurans* CH34 (Rm), *Rhodospirillum rubrum* (Rr), *Nitrococcus oceani* (No), *Nitrobacter winogradskyi* (Nw), *Halorhodospira halophila* (Hh), *Xanthobacter autotrophicus* (Xa), *Thiomicrospira crunogena* (Tc), *Methylobium petroleophilum* (Mp), *Thiobacillus denitrificans* (Td), *Nitrosospora multiformes* (Nm), *Alkalilimnicola ehrlichii* (Ae). I and II indicated *cbbl* and *cbll* operons. Af23270 type strain from *A. ferrooxidans*. Af Fe1 strain from Kusano and Sugawara (1993)[4].

Additional file 3: Sequences used to generate LOGOS of the intergenic region between *cbbR* and *cbbl1*.

Acknowledgements

This work was supported by a grant from Fondecyt 1090451, a Microsoft Sponsored Research Award, a Deutscher Akademischer Austausch Dienst (DAAD) scholarship to ME, a CONICYT graduate student grant to J-PC and grant from the Deutsche Forschungsgemeinschaft to BB.

Author details

¹Center for Bioinformatics and Genome Biology, MIFAB, Fundación Ciencia para la Vida and Depto. de Ciencias Biológicas, Facultad de Ciencias Biológicas, Universidad Andres Bello, Santiago, Chile. ²ICBM, Faculty of Medicine, University of Chile, Santiago, Chile. ³Institut für Mikrobiologie und Genetik, Georg-August-Universität Göttingen, Göttingen, Germany.

⁴Departamento de Acuicultura, Facultad de Recursos del Mar, Universidad de Antofagasta, Antofagasta, Chile.

Authors' contributions

DH, EJ and ME conceived the study. ME carried out the experiments. BB and J-PC contributed significantly to the analysis and interpretation of results. DH

drafted the manuscript. All authors contributed to the draft and approved the manuscript.

Received: 4 March 2010 Accepted: 27 August 2010

Published: 27 August 2010

References

1. Holmes D, Bonnefoy V: Genetic and bioinformatic insights into iron and sulfur oxidation mechanisms of bioleaching organisms. In *Biomining*. Edited by: Rawlings DE, Johnson B. D: Springer Berlin Heidelberg; 2007:281-307.
2. Valdes J, Pedrosa I, Quatrini R, Dodson RJ, Tettelin H, Blake R, Eisen JA, Holmes DS: *Acidithiobacillus ferrooxidans* metabolism: from genome sequence to industrial applications. *BMC Genomics* 2008, **9**:597.
3. Ask A Scientist. Carbon dioxide and water. [http://www.newton.dep.anl.gov/askasci/chem03/chem03573.htm].
4. Kusano T, Sugawara K: Specific binding of *Thiobacillus ferrooxidans* RbcR to the intergenic sequence between the *rbc* operon and the *rbcR* gene. *J Bacteriol* 1993, **175**:1019-1025.
5. Tabita FR: Molecular and cellular regulation of autotrophic carbon dioxide fixation in microorganisms. *Microbiol Rev* 1988, **52**:155-189.
6. Price GD, Badger MR, Woodger FJ, Long BM: Advances in understanding the cyanobacterial CO₂-concentrating-mechanism (CCM): functional components, Ci transporters, diversity, genetic regulation and prospects for engineering into plants. *J Exp Bot* 2008, **59**:1441-1461.
7. Cannon GC, Baker SH, Soyer F, Johnson DR, Bradburne CE, Mehlman JL, Davies PS, Jiang QL, Heinhorst S, Shively JM: Organization of carboxysome genes in the thiobacilli. *Curr Microbiol* 2003, **46**:115-119.
8. Appia-Ayme C QR, Denis Y, Denizot F, Silver S, Roberto F, Veloso F, Valdes J, Cárdenas JP, Esparza M, Orellana O, Jedlicki E, Bonnefoy V, Holmes D: Microarray and bioinformatic analyses suggest models for carbon metabolism in the autotroph *Acidithiobacillus ferrooxidans*. *Hydrometallurgy* 2006, **83**:273-280.
9. van den Bergh ER, Dijkhuizen L, Meijer WG: CbbR, a LysR-type transcriptional activator, is required for expression of the autotrophic CO₂ fixation enzymes of *Xanthobacter flavus*. *J Bacteriol* 1993, **175**:6097-6104.
10. Windhovel U, Bowien B: Identification of *cfxR*, an activator gene of autotrophic CO₂ fixation in *Alcaligenes eutrophus*. *Mol Microbiol* 1991, **5**:2695-2705.
11. Viale AM, Kobayashi H, Akazawa T, Henikoff S: *rbcR* [correction of *rcbR*], a gene coding for a member of the LysR family of transcriptional regulators, is located upstream of the expressed set of ribulose 1,5-bisphosphate carboxylase/oxygenase genes in the photosynthetic bacterium *Chromatium vinosum*. *J Bacteriol* 1991, **173**:5224-5229.
12. Strecker M, Sickinger E, English RS, Shively JM: Calvin cycle genes in *Nitrobacter vulgaris* T3. *FEMS Microbiology Letters* 1994, **120**:45-50.
13. Shively JM, van Keulen G, Meijer WG: Something from almost nothing: carbon dioxide fixation in chemoautotrophs. *Annu Rev Microbiol* 1998, **52**:191-230.
14. Beller HR, Chain PS, Letain TE, Chakicherla A, Larimer FW, Richardson PM, Coleman MA, Wood AP, Kelly DP: The genome sequence of the obligately chemolithoautotrophic, facultatively anaerobic bacterium *Thiobacillus denitrificans*. *J Bacteriol* 2006, **188**:1473-1488.
15. Gibson JL, Tabita FR: Nucleotide sequence and functional analysis of *cbbR*, a positive regulator of the Calvin cycle operons of *Rhodobacter sphaeroides*. *J Bacteriol* 1993, **175**:5778-5784.
16. Paoli GC, Soyer F, Shively J, Tabita FR: *Rhodobacter capsulatus* genes encoding form I ribulose-1,5-bisphosphate carboxylase/oxygenase (*cbblS*) and neighbouring genes were acquired by a horizontal gene transfer. *Microbiology* 1998, **144**(Pt 1):219-227.
17. Falcone DL, Tabita FR: Complementation analysis and regulation of CO₂ fixation gene expression in a ribulose 1,5-bisphosphate carboxylase-oxygenase deletion strain of *Rhodospirillum rubrum*. *J Bacteriol* 1993, **175**:5066-5077.
18. Toyoda K, Yoshizawa Y, Arai H, Ishii M, Igarashi Y: The role of two CbbRs in the transcriptional regulation of three ribulose-1,5-bisphosphate carboxylase/oxygenase genes in *Hydrogenovibrio marinus* strain MH-110. *Microbiology* 2005, **151**:3615-3625.
19. Wei X, Sayavedra-Soto LA, Arp DJ: The transcription of the *cbb* operon in *Nitrosomonas europaea*. *Microbiology* 2004, **150**:1869-1879.

20. Scott KM, Sievert SM, Abril FN, Ball LA, Barrett CJ, Blake RA, Boller AJ, Chain PS, Clark JA, Davis CR, Detter C, Do KF, Dobrinski KP, Faza BI, Fitzpatrick KA, Freyermuth SK, Harmer TL, Hauser LJ, Hügler M, Kerfeld CA, Klotz MG, Kong WW, Land M, Lapidus A, Larimer FW, Longo DL, Lucas S, Malfatti SA, Massey SE, Martin DD, McCuiddin Z, Meyer F, Moore JL, Ocampo LH Jr, Paul JH, Paulsen IT, Reep DK, Ren Q, Ross RL, Sato PY, Thomas P, Tinkham LE, Zeruth GT: **The genome of deep-sea vent chemolithoautotroph *Thiomicrospira crunogena* XCL-2.** *PLoS Biol* 2006, **4**:383.
21. Quatrini R, Lemfil C, Veloso FA, Pedroso I, Holmes DS, Jedlicki E: **Bioinformatic prediction and experimental verification of Fur-regulated genes in the extreme acidophile *Acidithiobacillus ferrooxidans*.** *Nucleic Acids Res* 2007, **35**:2153-2166.
22. Maniatis T: *Molecular cloning: a laboratory manual/T. Maniatis, E.F. Fritsch, J. Sambrook* Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory 1982.
23. Guacucano M, Levican G, S Holmes D, Jedlicki E: **An RT-PCR artifact in the characterization of bacterial operons.** *Electronic Journal of Biotechnology* 2000, **3**:12-13.
24. Bradford MM: **A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding.** *Anal Biochem* 1976, **72**:248-254.
25. Vaitukaitis J, Robbins JB, Nieschlag E, Ross GT: **A method for producing specific antisera with small doses of immunogen.** *J Clin Endocrinol Metab* 1971, **33**:988-991.
26. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG: **Clustal W and Clustal X version 2.0.** *Bioinformatics* 2007, **23**:2947-2948.
27. Bryson K, McGuffin LJ, Marsden RL, Ward JJ, Sodhi JS, Jones DT: **Protein structure prediction servers at University College London.** *Nucleic Acids Res* 2005, **33**:W36-38.
28. Hulo N, Bairoch A, Bulliard V, Cerutti L, De Castro E, Langendijk-Genevaux PS, Pagni M, Sigrist CJ: **The PROSITE database.** *Nucleic Acids Res* 2006, **34**:D227-230.
29. Bru C, Courcelle E, Carrere S, Beausse Y, Dalmar S, Kahn D: **The ProDom database of protein domain families: more emphasis on 3D.** *Nucleic Acids Res* 2005, **33**:D212-215.
30. Finn RD, Tate J, Mistry J, Coghill PC, Sammut SJ, Hotz HR, Ceric G, Forslund K, Eddy SR, Sonnhammer EL, Bateman A: **The Pfam protein families database.** *Nucleic Acids Res* 2008, **36**:D281-288.
31. Jacobs GH, Chen A, Stevens SG, Stockwell PA, Black MA, Tate WP, Brown CM: **TransTerm: a database to aid the analysis of regulatory sequences in mRNAs.** *Nucleic Acids Res* 2009, **37**:D72-76.
32. Kumar S, Nei M, Dudley J, Tamura K: **MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences.** *Brief Bioinform* 2008, **9**:299-306.
33. Terazono K, Hayashi NR, Igarashi Y: **CbbR, a LysR-type transcriptional regulator from *Hydrogenophilus thermoluteolus*, binds two cbb promoter regions.** *FEMS Microbiol Lett* 2001, **198**:151-157.
34. Dubbs JM, Bird TH, Bauer CE, Tabita FR: **Interaction of CbbR and RegA* transcription regulators with the *Rhodobacter sphaeroides* cbbI promoter-operator region.** *J Biol Chem* 2000, **275**:19224-19230.
35. Dubbs P, Dubbs JM, Tabita FR: **Effector-mediated interaction of CbbRI and CbbRII regulators with target sequences in *Rhodobacter capsulatus*.** *J Bacteriol* 2004, **186**:8026-8035.
36. Bowien B, Kusian B: **Genetics and control of CO₂ assimilation in the chemoautotroph *Ralstonia eutropha*.** *Arch Microbiol* 2002, **178**:85-93.
37. Schell MA: **Molecular biology of the LysR family of transcriptional regulators.** *Annu Rev Microbiol* 1993, **47**:597-626.
38. Knochel T, Ivens A, Hester G, Gonzalez A, Bauerle R, Wilmanns M, Kirschner K, Jansonius JN: **The crystal structure of anthranilate synthase from *Sulfolobus solfataricus*: functional implications.** *Proc Natl Acad Sci USA* 1999, **96**:9479-9484.
39. Lin X, Xu S, Yang Y, Wu J, Wang H, Shen H: **Purification and characterization of anthranilate synthase component I (TrpE) from *Mycobacterium tuberculosis* H37Rv.** *Protein Expr Purif* 2009, **64**:8-15.
40. Grzeszlik C, Jeffke T, Schaferjohann J, Kusian B, Bowien B: **Phosphoenolpyruvate is a signal metabolite in transcriptional control of the cbb CO₂ fixation operons in *Ralstonia eutropha*.** *J Mol Microbiol Biotechnol* 2000, **2**:311-320.
41. Kusian B, Bowien B: **Organization and regulation of cbb CO₂ assimilation genes in autotrophic bacteria.** *FEMS Microbiol Rev* 1997, **21**:135-155.
42. Ivens A, Mayans O, Szadkowski H, Wilmanns M, Kirschner K: **Purification, characterization and crystallization of thermostable anthranilate phosphoribosyltransferase from *Sulfolobus solfataricus*.** *Eur J Biochem* 2001, **268**:2246-2252.
43. Esparza M, Bowien B, Holmes DS, Jedlicki E: **Gene organization and CO₂-responsive expression of four cbb operons in the biomining bacterium *Acidithiobacillus ferrooxidans*.** *Advanced Materials Research* 2009, **71-73**:207-210.

doi:10.1186/1471-2180-10-229

Cite this article as: Esparza et al.: Genes and pathways for CO₂ fixation in the obligate, chemolithoautotrophic acidophile, *Acidithiobacillus ferrooxidans*, Carbon fixation in *A. ferrooxidans*. *BMC Microbiology* 2010 10:229.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

