



# Alternative splicing of HER2: a novel mediator of EGFR TKI resistance

Zachary A. Yochum<sup>1,2^</sup>, Liza C. Villaruz<sup>1^</sup>

<sup>1</sup>Division of Hematology-Oncology, Department of Medicine, UPMC Hillman Cancer Center, Pittsburgh, PA, USA; <sup>2</sup>Medical Scientist Training Program, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

Correspondence to: Liza C. Villaruz, MD. UPMC Hillman Cancer Center, 5150 Centre Avenue, Pittsburgh, PA 15232, USA. Email: villaruz@upmc.edu.

Provenance and Peer Review: This article was commissioned by the editorial office, *Translational Lung Cancer Research*. The article did not undergo external peer review.

Comment on: Hsu CC, Liao BC, Liao WY, *et al.* Exon 16-Skipping HER2 as a Novel Mechanism of Osimertinib Resistance in EGFR L858R/T790M-Positive Non-Small Cell Lung Cancer. *J Thorac Oncol* 2020;15:50-61.

Submitted Apr 30, 2020. Accepted for publication May 14, 2020.

doi: 10.21037/tlcr-20-618

View this article at: <http://dx.doi.org/10.21037/tlcr-20-618>

Over the past three decades, lung cancer has been the leading cause of cancer deaths for both men and women (1). Lung cancer mortality remains high as it is usually diagnosed at a late stage and therapies in the metastatic setting have limited efficacy (1). However, there have been marked advances in the treatment of non-small cell lung cancer (NSCLC), the largest subset of lung cancer, since the introduction of immunotherapy and targeted therapy. When patients with advanced NSCLC are initially diagnosed, they are subtyped based on molecular drivers. While subsets of NSCLC patients with a molecular driver, such as those with *KRAS* mutations, do not as of yet have an approved targeted therapy, approximately 25% of NSCLC patients have a commercially available targeted agent (2,3). The largest group with a targetable molecular driver is the approximately 15% of patients with sensitizing *EGFR* mutations (2,3).

Multiple Phase III trials have demonstrated that first- and second-generation *EGFR* TKIs lead to improved objective response rates (ORR) and progression-free survival (PFS) compared with platinum-based chemotherapy (4,5). However, the efficacy of first- and second-generation *EGFR* TKIs in *EGFR*-mutant NSCLC is limited by the frequency of *de-novo* resistance (~20–30%) and the inevitability of acquired resistance (5–7). Attempts to establish mechanisms of resistance to first- and second-generation *EGFR* TKIs

have demonstrated that a significant proportion of patients develop secondary *EGFR* mutations (6). Approximately 50% of patients acquire the “gatekeeper” T790M *EGFR* mutation, which renders first- and second-generation *EGFR* TKIs ineffective as they can no longer access the ATP binding cleft of *EGFR* (6,8). T790M mutant *EGFR* remains sensitive to third-generation *EGFR* TKIs, such as osimertinib, which binds covalently to a cysteine residue (C797) in the ATP binding cleft (6,8). Osimertinib was initially approved as a treatment for patients with acquired T790M *EGFR* mutations following progression on first-line *EGFR* TKIs as multiple trials demonstrated osimertinib was superior to chemotherapy in this setting (7). Recently, osimertinib became the recommended first-line therapy for patients with *EGFR*-mutant NSCLC. The FLAURA trial demonstrated that using osimertinib as a first-line agent resulted in superior PFS and overall survival when compared to first-line erlotinib or gefitinib (9).

Despite improved clinical outcomes, response to first-line osimertinib remains limited by acquired resistance (9). The mechanisms of resistance to first-line osimertinib are not well-established. Early studies have demonstrated that resistance mechanisms appear to be less *EGFR*-dependent, as tertiary *EGFR* mutations and amplification of wild-type (WT) *EGFR* only account for resistance in 5–10% of patients (10–12). The predominant *EGFR*-

<sup>^</sup> ORCID: Zachary A. Yochum: 0000-0001-6483-8089; Liza C. Villaruz: 0000-0001-7456-523X.

independent mechanism is *MET* amplification, accounting for resistance in approximately 15% of patients (10-12). Unfortunately, the mechanism of resistance in approximately 50% of patients remains unknown (10-12). Additionally, there are no targeted therapies approved for patients following progression on first-line osimertinib. Immunotherapy in combination with platinum-based chemotherapy may be a viable therapeutic option in this setting as the recent IMpower 150 trial demonstrated that chemo-immunotherapy plus bevacizumab improved PFS and OS when compared to chemotherapy alone in *EGFR*-mutant and *ALK*-positive patients who had progressed on a targeted therapy (13). The IMpower 130 trial failed to demonstrate a clinical benefit for the addition of immunotherapy to chemotherapy without bevacizumab in *EGFR*-mutant NSCLCs (14). It is important to note that in the large registrational trials evaluating pembrolizumab alone or in combination with chemotherapy in patients with non-squamous NSCLC did not enroll patients with *EGFR* mutations or *ALK* translocations (15-17). An ongoing phase III trial, KEYNOTE 789, is specifically aimed at addressing if chemo-immunotherapy is a viable strategy in patients with *EGFR*-mutant NSCLC resistant to EGFR TKIs.

More recently, investigation into osimertinib resistance mechanisms in NSCLC has identified the HER2 signaling pathway as a driver of resistance. *HER2* amplification and mutations are molecular drivers in 5–6% of all NSCLC patients (3,18). *HER2* amplification has also been established as a driver of first- and second-generation EGFR TKI resistance in tumors that lack T790M *EGFR* mutations (19). Initial studies have identified *HER2* amplification or mutations in approximately 5% of patients with resistance to first-line osimertinib (10,11). While the HER2 pathway has been implicated in osimertinib resistance, it remains unclear whether additional HER2 aberrations mediate resistance to first-line osimertinib and the validity of HER2 pathways alterations as a therapeutic target in osimertinib resistance.


In a recent report, Hsu and colleagues demonstrated that the exon 16-skipping splice variant of HER2 (HER2D16) is a mediator of osimertinib resistance. Initially, the authors examine the expression of HER2D16 in a patient with metastatic *EGFR*-mutant NSCLC (20). The patient in the study had known lung lesions, liver metastases, and a malignant pleural effusion following progression on two separate EGFR TKIs (gefitinib and erlotinib) and two different lines of chemotherapy (20). Analysis of the malignant pleural effusion revealed L858R and T790M

*EGFR* mutations. Osimertinib was subsequently initiated and follow-up revealed that the lung lesion responded to therapy but the liver metastasis progressed. Analysis of cell-free DNA on progression revealed increase in the allelic fraction of HER2D16, *HER2* amplification, and loss of T790M *EGFR* expression (20).

After observing HER2D16 expression in a patient following the development of osimertinib resistance, the authors then examined the role of HER2D16 in mediating osimertinib resistance *in vitro*. Given previous studies demonstrated that HER2D16 primarily functions as a homodimer in breast cancer (21-23), Hsu *et al.* investigated whether HER2D16 functions as a homo- or heterodimer in NSCLC cells (20). Interestingly, both in cells with low HER family expression and in cells expressing L858R/T790M mutant *EGFR*, HER2D16 has the ability to homodimerize and form heterodimers with EGFR (Figure 1) (20). HER2 and HER2D16 overexpression in cells lacking HER family expression was sufficient to cause ERK activation, however, it was not sufficient to cause increased ERK or AKT signaling in cells expressing L858R/T790M mutant *EGFR* (20). The inability of HER2 to increase ERK and AKT activation in *EGFR*-mutant NSCLC cells likely reflects maximal activation of these pathways through constitutively active EGFR.

Interestingly, HER2 and HER2D16 overexpression was sufficient to cause osimertinib resistance in H1975 cells, an L858R/T790M *EGFR*-mutant NSCLC cell line (20). Additionally, in H1975 cells, HER2D16 overexpression led to a significantly higher osimertinib IC<sub>50</sub> when compared to HER2 overexpressing and control cells (20). Consistent with previous studies (8), osimertinib suppressed EGFR, ERK, and AKT activation in H1975 cells (20). In HER2 overexpressing cells, osimertinib treatment resulted in suppressed ERK activation and partially suppressed EGFR activation, however, downstream AKT signaling was maintained (20). In HER2D16 overexpressing cells, EGFR, ERK, and AKT activation was maintained in the presence of osimertinib (Figure 1) (20). Additionally, osimertinib did not prevent HER2D16 from forming hetero- or homodimers (20). These data suggest that HER2D16 likely causes osimertinib resistance through persistent activation of EGFR and canonical EGFR signaling pathways.

Previous studies have demonstrated that HER2D16 promotes breast cancer tumorigenesis through activation of SRC (23,24). However, HER2D16 overexpression did not increase total SRC or phospho-SRC expression in H1975 cells (20). Additionally, genetic or pharmacologic inhibition



**Figure 1** Overexpression of HER2D16, a HER2 exon-16 splice variant, is a novel mediator of osimertinib resistance in *EGFR*-mutant non-small cell lung cancer (NSCLC) and is associated with reactivation of EGFR pathway signaling. In the discussed report, the authors demonstrate that HER2D16 can homodimerize and/or heterodimerize with mutant EGFR and WT EGFR. Additionally, as HER2D16 expression frequently occurs with *HER2* amplification, it may form heterodimers with WT HER2. HER2D16 overexpression was sufficient to cause osimertinib resistance in *EGFR*-mutant NSCLC cell with a T790M mutation and was associated with reactivation of EGFR and downstream mitogenic/proliferative pathways such as the MAPK and PI3K/AKT pathways. The combination of afatinib, a pan-HER TKI, and osimertinib overcame HER2D16-mediated resistance. Future studies are needed to determine if HER2D16 drives resistance solely through reactivation of MAPK and/or AKT or if other known functions of HER2D16 such as suppression of apoptosis and induction of epithelial-mesenchymal transition also play a role. EMT, epithelial-mesenchymal transition, EMT-TF, epithelial-mesenchymal transition-transcription factors.

(with dasatinib) of SRC was unable to overcome resistance associated with HER2 or HER2D16 overexpression (20). Interestingly, the use of an irreversible pan-HER inhibitor, afatinib, was able to overcome resistance mediated both by HER2 and HER2D16 overexpression (20). In H1975 cells overexpressing HER2 or HER2D16, the combination of osimertinib and afatinib had synergistic activity that was accompanied by suppression of EGFR and downstream ERK/AKT pathway activation (Figure 1) (20). These results suggest that combining osimertinib with a pan-HER inhibitor may be a strategy to overcome HER2-mediated osimertinib resistance.

The recent report by Hsu *et al.* is the first study to implicate the Exon-16 skipping HER2 splice variant

as a mediator of osimertinib resistance. A strength of this study is that the authors demonstrate HER2D16 expression in an *EGFR*-mutant NSCLC patient after developing osimertinib resistance. This clinical evidence is accompanied by clear *in vitro* evidence that HER2D16 overexpression in *EGFR*-mutant NSCLC cells is sufficient to cause osimertinib resistance (20). The authors also elucidate novel functions of the HER2D16 splice variant in NSCLC cells by demonstrating HER2D16 functions both as a homo- and heterodimer with EGFR and functions to promote therapeutic resistance independently of SRC signaling (20). This differs from the function of HER2D16 in breast cancer. Previous studies demonstrate that in breast cancer HER2D16 promotes tumorigenesis primarily

as a homodimer and the ability of HER2D16 to mediate therapeutic resistance requires SRC activity (20-24). Studies investigating the factors that influence HER2D12 dimerization and the structure of the HER2 exon-16 skipping splice variant are lacking. Future studies are needed to delineate the structural changes in HER2 caused by exon-16 skipping, structural changes in HER2D16 induced by hetero- versus homodimer formation, and the similarity in kinase domain structure between HER2D16, WT HER2, and WT EGFR. These structural studies are important as they may have therapeutic significance as mutant or aberrant receptor tyrosine kinases (RTKs) can share significant structural overlap with wild-type RTKs. For example, the kinase domain of Exon-20 insertion mutant *EGFR* shares significant structural similarity to WT *EGFR* (25). Afatinib, which irreversibly inhibits wild-type *EGFR* as well as HER2 and HER4, has limited efficacy in the *EGFR* exon-20 insertion mutant NSCLC population as the structural similarity between the exon-20 insertion mutant *EGFR* and WT *EGFR* significantly narrows the therapeutic index of afatinib (25). The dose *in vivo* required to effectively inhibit the exon-20 insertion mutant *EGFR* is limited by the side-effects of inhibiting *EGFR* WT (25).

While the authors provide a possible therapeutic strategy of combining afatinib with osimertinib, to overcome HER2D16-mediated therapeutic resistance, afatinib may have additional side-effects in combination with osimertinib associated with inhibition of WT *EGFR*. Consequently, paralleling structural studies of HER2D16, *in vivo* studies are needed to ensure that afatinib in combination with osimertinib is not only efficacious in the setting of HER2D16-driven osimertinib resistance but also can adequately inhibit HER2D16 without significant toxicity or side-effects associated with WT *EGFR* inhibition. This combination was utilized in a *EGFR*-mutant NSCLC patient who developed a T790M *EGFR* mutation, *HER2* amplification, and a *HER2* Exon 20 mutation following progression on icotinib, a first-generation *EGFR* TKI (26). The combination of osimertinib and afatinib was well-tolerated (grade 2 diarrhea and grade 1 rash) and led to a partial response after 4 weeks of treatment (26). Conversely, a recent study demonstrated that the combination of osimertinib and erlotinib in a transgenic model of *EGFR*-mutant lung cancer was associated with kidney fibrosis and atrophy (27). Future studies should also evaluate osimertinib in combination with other HER2-specific inhibitors with well-established toxicity profiles like trastuzumab. While the evidence is conflicting, HER2D16 may be a marker

of HER2 dependence and a predictor of response to trastuzumab (18,22,23). Interestingly, a recent case report identified an activating *HER2* mutation, *HER2* amplification and a T790M *EGFR* mutation in a patient with *EGFR*-mutant NSCLC following progression on gefitinib. The patient received osimertinib and trastuzumab, which resulted in a clinical and molecular response, without overt toxicity (28).

An unanswered question remains as to the relevance of HER2D16 as a mechanism of resistance to first-line osimertinib, and the role of HER2D16 as a mediator of resistance in *EGFR*-mutant NSCLC that does not harbor T790M mutations. While *HER2* amplification and mutations have been observed as a resistance mechanism to first-line osimertinib (11), HER2 exon-16 splice variant expression has not been evaluated in this setting. Initial studies suggest that mechanisms of resistance to first-line osimertinib appear to be more *EGFR*-independent when compared to second-line osimertinib (10-12). For example, a recent study demonstrated that histological transformation to squamous NSCLC or small cell lung cancer (SCLC) occur in a higher frequency in the setting of first-line osimertinib (12). Additionally, given the presence of HER2D16 expression before the initiation of osimertinib and sequencing data only being available from plasma, it is unclear whether HER2D16 expression was present in the aforementioned patient's lung lesion or liver metastasis before osimertinib treatment. Importantly, the patient's lung lesion responded to osimertinib treatment while the patient's liver metastasis did not (20). The presence of HER2D16 expression in circulating tumor DNA (ctDNA) before osimertinib and the lack of response of the liver lesion suggests that HER2D16 was likely a mechanism of *de-novo*, rather than acquired, osimertinib resistance in this patient's liver lesion. The heterogeneity that was present in the aforementioned patient's liver and lung lesions stresses the importance of using paired biopsies in evaluating mechanisms of osimertinib resistance. Utilizing paired biopsies from cohorts of patients who have progressed on first-line osimertinib would allow investigators to establish the relevance of HER2D16 expression as mechanism of *de-novo* and/or acquired resistance. Additionally, obtaining ctDNA pre- and post-treatment would allow for the validation of using ctDNA as a means detecting HER2D16.

While the authors demonstrate that HER2D16 overexpression was associated with persistent *EGFR*, AKT, and ERK activation, it is unclear if this is the mechanism(s) by which HER2D16 promotes resistance.

Furthermore, it is unclear if HER2D16 homodimerization and/or heterodimerization with WT HER2, mutant EGFR, and/or WT EGFR is required for osimertinib resistance (Figure 1). Future studies should be aimed at determining whether HER2D16 promotes osimertinib resistance solely through activation of EGFR and/or downstream mitogenic/proliferative signaling pathways or whether HER2D16 promotes resistance through alternative mechanisms (Figure 1). Previous work in breast cancer has revealed that HER2D16 expression promotes epithelial-mesenchymal transition (EMT) and stemness potentially through increased expression of AXL and EMT-transcription factors, such as SNAIL1 and TWIST1 (Figure 1) (21,23,29). Interestingly, EMT is associated with resistance to EGFR TKIs, and overexpression of EMT-transcription factors is both sufficient and required for resistance to first- and third-generation EGFR TKIs in EGFR-mutant NSCLC (30,31). Numerous studies have also demonstrated that suppression of apoptosis, specifically through downregulation of the pro-apoptotic protein BIM, mediates resistance to EGFR TKIs (32). Future studies should evaluate whether HER2D16 suppresses osimertinib-induced apoptosis given previous studies have demonstrated that HER2D16 can lead to tamoxifen resistance in breast cancer cells through upregulation of anti-apoptotic BCL-2 proteins (Figure 1) (33). Determining the mechanism(s) why HER2D16 promotes resistance is important as it may render additional therapeutic targets. For example, tumors that undergo EMT as a mechanism of resistance to EGFR TKIs are more sensitive to BCL-2/BCL-xL inhibitors (30,31).

In summary, Hsu and colleagues demonstrated HER2D16, a HER2 splice variant, is a novel mediator of osimertinib resistance in EGFR-mutant NSCLC. This is the first study to demonstrate HER2D expression in a EGFR-mutant NSCLC patient with osimertinib resistance. They also demonstrated that HER2D16 overexpression is sufficient to mediate osimertinib resistance in EGFR-mutant NSCLC cells and that combining afatinib and osimertinib may be a novel therapeutic strategy to overcome HER2D16-mediated resistance. Future studies are needed to validate afatinib and osimertinib as a strategy to overcome HER2D16-mediated resistance *in vivo* and to determine the frequency of this mechanism in patients with resistance to first-line osimertinib. While osimertinib has benefited patients with EGFR-mutant NSCLC, therapeutic resistance remains inevitable. Chemotherapy-based regimens with the possible addition of immunotherapy are the only currently available options for patients following

progression on osimertinib in the absence of an actionable resistance mechanisms. Studies like Hsu and colleagues' recent work aimed at identifying and characterizing novel, potentially actionable, mediators of osimertinib resistance are critical to improving outcomes and sparing toxicity associated with chemotherapy in these patients.

## Acknowledgments

*Funding:* None.

## Footnote

*Conflicts of Interest:* Both authors have completed the ICMJE uniform disclosure form (available at <http://dx.doi.org/10.21037/tlcr-20-618>). LCV receives commercial research grants from Genentech, AstraZenca, Exelixis, Incyte, Merck, Rain, and GlaxoSmithKline and is a paid consultant for Achilles. ZAY has no conflicts of interest to disclose.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin* 2020;70:7-30.
2. Kris MG, Johnson BE, Berry LD, et al. Using Multiplexed Assays of Oncogenic Drivers in Lung Cancers to Select Targeted Drugs. *JAMA* 2014;311:1998-2006.
3. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature* 2014;511:543-50.
4. Fukuoka M, Wu YL, Thongprasert S, et al. Biomarker analyses and final overall survival results from a phase III,

- randomized, open-label, first-line study of gefitinib versus carboplatin/paclitaxel in clinically selected patients with advanced non-small-cell lung cancer in Asia (IPASS). *J Clin Oncol* 2011;29:2866-74.
5. Wu YL, Zhou C, Liang CK, et al. First-line erlotinib versus gemcitabine/cisplatin in patients with advanced EGFR mutation-positive non-small-cell lung cancer: analyses from the phase III, randomized, open-label, ENSURE study. *Ann Oncol* 2015;26:1883-9.
  6. Sequist LV, Waltman BA, Dias-Santagata D, et al. Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors. *Sci Transl Med* 2011;3:75ra26.
  7. Mok TS, Wu YL, Ahn MJ, et al. Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer. *N Engl J Med* 2017;376:629-40.
  8. Cross DA, Ashton SE, Ghiorghiu S, et al. AZD9291, an irreversible EGFR TKI, overcomes T790M-mediated resistance to EGFR inhibitors in lung cancer. *Cancer Discov* 2014;4:1046-61.
  9. Soria JC, Ohe Y, Vansteenkiste J, et al. Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *N Engl J Med* 2018;378:113-25.
  10. Ramalingam SS, Cheng Y, Zhou C, et al. Mechanisms of acquired resistance to first-line osimertinib: Preliminary data from the phase III FLAURA study. *Ann Oncol* 2018;29:viii740.
  11. Leonetti A, Sharma S, Minari R, et al. Resistance mechanisms to osimertinib in EGFR-mutated non-small cell lung cancer. *Br J Cancer* 2019;121:725-37.
  12. Schoenfeld AJ, Chan JM, Kubota D, et al. Tumor Analyses Reveal Squamous Transformation and Off-Target Alterations As Early Resistance Mechanisms to First-line Osimertinib in EGFR-Mutant Lung Cancer. *Clin Cancer Res* 2020;26:2654-63.
  13. Socinski MA, Jotte RM, Cappuzzo F, et al. Atezolizumab for First-Line Treatment of Metastatic Nonsquamous NSCLC. *N Engl J Med* 2018;378:2288-301.
  14. Cappuzzo F, McCleod M, Hussein M, et al. IMpower130: Progression-free survival (PFS) and safety analysis from a randomised phase III study of carboplatin + nab-paclitaxel (CnP) with or without atezolizumab (atezo) as first-line (1L) therapy in advanced non-squamous NSCLC. *Ann Oncol* 2018;29:viii742-3.
  15. Reck M, Rodriguez-Abreu D, Robinson AG, et al. Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell Lung Cancer. *N Engl J Med* 2016;375:1823-33.
  16. Mok TSK, Wu YL, Kudaba I, et al. Pembrolizumab versus chemotherapy for previously untreated, PD-L1-expressing, locally advanced or metastatic non-small-cell lung cancer (KEYNOTE-042): a randomised, open-label, controlled, phase 3 trial. *Lancet* 2019;393:1819-30.
  17. Gandhi L, Rodriguez-Abreu D, Gadgeel S, et al. Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell Lung Cancer. *N Engl J Med* 2018;378:2078-92.
  18. Li BT, Ross DS, Aisner DL, et al. HER2 Amplification and HER2 Mutation Are Distinct Molecular Targets in Lung Cancers. *J Thorac Oncol* 2016;11:414-9.
  19. Takezawa K, Pirazzoli V, Arcila ME, et al. HER2 amplification: a potential mechanism of acquired resistance to EGFR inhibition in EGFR-mutant lung cancers that lack the second-site EGFR T790M mutation. *Cancer Discov* 2012;2:922-33.
  20. Hsu CC, Liao BC, Liao WY, et al. Exon 16-Skipping HER2 as a Novel Mechanism of Osimertinib Resistance in EGFR L858R/T790M-Positive Non-Small Cell Lung Cancer. *J Thorac Oncol* 2020;15:50-61.
  21. Alajati A, Sausgruber N, Aceto N, et al. Mammary tumor formation and metastasis evoked by a HER2 splice variant. *Cancer Res* 2013;73:5320-7.
  22. Marchini C, Gabrielli F, Iezzi M, et al. The human splice variant Delta16HER2 induces rapid tumor onset in a reporter transgenic mouse. *PLoS One* 2011;6:e18727.
  23. Mitra D, Brumlik MJ, Okamgba SU, et al. An oncogenic isoform of HER2 associated with locally disseminated breast cancer and trastuzumab resistance. *Mol Cancer Ther* 2009;8:2152-62.
  24. Castagnoli L, Iezzi M, Ghedini GC, et al. Activated d16HER2 homodimers and SRC kinase mediate optimal efficacy for trastuzumab. *Cancer Res* 2014;74:6248-59.
  25. Yasuda H, Park E, Yun CH, et al. Structural, biochemical, and clinical characterization of epidermal growth factor receptor (EGFR) exon 20 insertion mutations in lung cancer. *Sci Transl Med* 2013;5:216ra177.
  26. Zhang P, Nie X, Wang B, et al. Combined therapy with osimertinib and afatinib in a lung adenocarcinoma patient with EGFR T790M mutation and multiple HER2 alterations after resistance to icotinib: A case report. *Thorac Cancer* 2018;9:1774-7.
  27. Starrett JH, Guernet AA, Cuomo ME, et al. Drug sensitivity and allele-specificity of first-line osimertinib resistance EGFR mutations. *Cancer Res* 2020;80:2017-30.
  28. Ralki M, Maes B, Pat K, et al. Triple Trouble: A Case of Multiple Resistance Mechanisms after First Generation EGFR-TKI in NSCLC. *Case Rep Oncol* 2019;12:625-30.

29. Goyette M-A, Duhamel S, Aubert L, et al. The Receptor Tyrosine Kinase AXL Is Required at Multiple Steps of the Metastatic Cascade during HER2-Positive Breast Cancer Progression. *Cell Reports* 2018;23:1476-90.
30. Song KA, Niederst MJ, Lochmann TL, et al. Epithelial-to-Mesenchymal Transition Antagonizes Response to Targeted Therapies in Lung Cancer by Suppressing BIM. *Clin Cancer Res* 2018;24:197.
31. Yochum ZA, Cades J, Wang H, et al. Targeting the EMT transcription factor TWIST1 overcomes resistance to EGFR inhibitors in EGFR-mutant non-small-cell lung cancer. *Oncogene* 2019;38:656-70.
32. Costa DB, Halmos B, Kumar A, et al. BIM mediates EGFR tyrosine kinase inhibitor-induced apoptosis in lung cancers with oncogenic EGFR mutations. *PLoS Med* 2007;4:1669-79; discussion 1680.
33. Cittelly DM, Das PM, Salvo VA, et al. Oncogenic HER2{Delta}16 suppresses miR-15a/16 and deregulates BCL-2 to promote endocrine resistance of breast tumors. *Carcinogenesis* 2010;31:2049-57.

**Cite this article as:** Yochum ZA, Villaruz LC. Alternative splicing of HER2: a novel mediator of EGFR TKI resistance. *Transl Lung Cancer Res* 2020;9(4):1606-1612. doi: 10.21037/tlcr-20-618