



OPEN

Characterization of outdoor air pollution from solid fuel combustion in Xuanwei and Fuyuan, a rural region of China

Wei Hu^{1✉}, George Downward², Jason Y. Y. Wong¹, Boris Reiss³, Nathaniel Rothman¹, Lützen Portengen², Jihua Li⁴, Rena R. Jones¹, Yunchao Huang⁵, Kaiyun Yang⁵, Ying Chen⁵, Jun Xu⁶, Jun He⁴, Bryan Bassig¹, Wei Jie Seow⁷, H. Dean Hosgood⁸, Linlin Zhang⁹, Guoping Wu⁹, Fusheng Wei^{9,10}, Roel Vermeulen^{2,10} & Qing Lan^{1,10}

Outdoor air pollution is a growing public health concern, particularly in urban settings. However, there are limited epidemiological data on outdoor air pollution in rural areas with substantial levels of air pollution attributed to solid fuel burning for household cooking and heating. Xuanwei and Fuyuan are rural counties in China where the domestic combustion of locally sourced bituminous (“smoky”) coal has been associated with the highest lung cancer rates in China. We previously assessed indoor and personal air pollution exposures in this area; however, the influence of indoor coal combustion and household ventilation on outdoor air pollution has not been assessed. Therefore, we measured outdoor fine particulate matter (PM_{2.5}), species of polycyclic aromatic hydrocarbons (PAHs) including naphthalene (NAP) and the known carcinogen benzo(a)pyrene (BaP), sulfur dioxide (SO₂), and nitrogen dioxide (NO₂) over two consecutive 24-h sampling periods in 29 villages. Just over half of the villages were revisited two to nine months after the initial sampling period to repeat all measurements. The overall geometric mean (GM) of outdoor PM_{2.5}, BaP, NAP, and NO₂ were 45.3 µg/m³, 9.7 ng/m³, 707.7 ng/m³, and 91.5 µg/m³, respectively. Using linear mixed effects models, we found that burning smoky coal was associated with higher outdoor BaP concentrations [GM ratio (GMR) = 2.79] and lower outdoor SO₂ detection rates (GMR = 0.43), compared to areas burning smokeless coal. Areas with predominantly ventilated stoves (> 50% of stoves) had higher outdoor BaP (GMR = 1.49) compared to areas with fewer ventilated stoves. These results show that outdoor air pollution in a rural region of China was associated with the type of coal used for cooking and heating indoors and the presence of stove ventilation. Our findings suggest that efforts of household stove improvement to reduce indoor air pollution have resulted in higher outdoor air pollution levels. Further reducing adverse health effects in rural villages from household coal combustion will require the use of cleaner fuel types.

Outdoor air pollution is a major environmental health concern that was linked to 3.7 million deaths worldwide in 2012 and 4.2 million deaths in 2016 (six percent due to lung cancer)^{1–3}. An upward trend in attributable deaths

¹Division of Cancer Epidemiology and Genetics, Occupational and Environmental Epidemiology Branch, National Cancer Institute, National Institutes of Health, 9609 Medical Center Drive, Rockville, MD 20850, USA. ²Institute for Risk Assessment Sciences, Division of Environmental Epidemiology, Utrecht University, Utrecht 3508 TD, The Netherlands. ³Mel and Enid Zuckerman College of Public Health, University of Arizona, Tucson, AZ 85724, USA. ⁴Qujing Center for Disease Control and Prevention, Qujing 655000, Yunnan, China. ⁵Department of Cardiothoracic Surgery, The Third Affiliated Hospital of Kunming Medical University (Yunnan Cancer Hospital, Yunnan Cancer Center), Kunming 650118, China. ⁶School of Public Health, Li Ka Shing, Faculty of Medicine, The University of Hong Kong, Hong Kong, SAR, China. ⁷Saw Swee Hock School of Public Health, National University of Singapore, Singapore 119077, Singapore. ⁸Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY 10461-1900, USA. ⁹China National Environmental Monitoring Center, Beijing 100012, China. ¹⁰These authors jointly supervised: Fusheng Wei, Roel Vermeulen and Qing Lan. ✉email: wei.hu@nih.gov

from 1990 to 2015 was partially due to increasing outdoor air pollution in low- and middle-income countries⁴. Outdoor air pollution is considered as a leading environmental cause of lung cancer by the International Agency for Research on Cancer (IARC) which recently classified outdoor air pollution and particulate matter (PM) as a Group 1 carcinogen (carcinogenic to humans)^{5–7}.

While automobile traffic is the predominant source of outdoor air pollution in developed urban areas, this is not the case in rural, underdeveloped areas where local sources such as cooking and heating stoves contribute significantly to air pollution levels⁸. Whereas many epidemiological studies focus on the health impacts of outdoor air pollution in urban areas, outdoor air pollution and its adverse health effects in rural areas are often overlooked⁸. More than 60% of the Chinese population lives in rural regions. In this population, household air pollution (HAP) due to domestic combustion of solid fuels contributes significantly to the total burden of disease⁹; however, epidemiological data on outdoor air pollution exposure in rural China is limited. Xuanwei and Fuyuan are rural counties located in Southwestern China that have elevated rates of nonmalignant and malignant lung diseases including the highest lung cancer rate in China that is directly associated with HAP from bituminous (“smoky”) coal^{10,11}. We previously reported indoor and personal exposure levels to PM_{2.5}, polycyclic aromatic hydrocarbons (PAHs), black carbon, NO₂, and SO₂ from burning solid fuels in Xuanwei and Fuyuan^{12–15}; and reported outdoor measurements of black carbon were positively correlated with the level of indoor measurements¹⁵. Additionally, we previously reported that there was variation in lung cancer risk for specific subtypes of smoky coal mined from different Xuanwei geological coal deposits^{16,17}, while improving home ventilation by installing stoves with chimneys and converting to portable stoves was associated with both a reduction in lung cancer rates and specific HAP constituents in this region^{18,19}. However, the contribution to neighborhood-level outdoor air pollution from both household ventilation and the indoor burning of coal mined from various coal deposits has not been assessed in this region. Given the clear need for outdoor air pollution research in rural settings, especially within China, we evaluated outdoor air pollution concentrations and the potential factors associated with those levels in Xuanwei and Fuyuan. As a part of a comprehensive evaluation of air quality study in Xuanwei and Fuyuan, this paper characterizes the outdoor air pollution in a rural area with higher incidence of lung cancer in China and provides quantitative evidence concerning the necessity of changing to clean fuel energy.

Results

Figure 1 shows the distribution of the 29 villages (16 of them were visited at both Phases I and II) and coal mine regions in Xuanwei and Fuyuan counties. Most villages were located in smoky coal mine areas in the center, east, north, and south parts of Xuanwei. Several villages in southwestern Xuanwei were in a historically smokeless coal mine region (27), where farmers have changed to wood. In Fuyuan, smokeless coal mines are predominantly located in the south, while smoky coal mines are in the north.

Background information of villages. Background measurement information about the villages and meteorological factors are summarized in Table 1. Less than 14% of the villages were within 5 km of a known factory, mine, or power station. The average village population size was 840 (750 SD). In the past, smoky coal was used in areas of Xuanwei and Fuyuan that had undergone stove improvement interventions, including the installation of chimneys, in an effort to reduce HAP^{18,19}. As a result, more than half of all households in 22 of the 29 villages (75.9%) included stoves ventilated with chimneys.

Outdoor pollution in coal deposits. Table 2 shows outdoor measurements by coal deposit. The overall GM of outdoor PM_{2.5}, BaP, NAP, and NO₂ were 45.3 µg/m³, 9.7 ng/m³, 708 ng/m³, and 91.5 µg/m³ respectively. The overall detection rate of SO₂ was 26.1%. Outdoor pollutant concentrations or detection rates (%Detect) were similar between the two counties [GM(GSD): 51.6(1.7) and 40.2(2.0) µg/m³ for PM_{2.5}; 10.5(1.7) and 8.9(1.6) ng/m³ for BaP; 730(2.3) and 678(2.6) ng/m³ for NAP; 93.2(1.4) and 89.7(1.5) µg/m³ for NO₂, and %Detect: 34.8 and 34.1% for SO₂, in Xuanwei and Fuyuan, respectively]. Based on an ANOVA test there is significant variation within each coal deposit for PM_{2.5} absorbance (PM_{abs}) ($p < 0.05$). A Tukey HSD test further reveals that PM_{abs} in villages located in several smoky coal deposits [GM(GSD): 5.8(1.7)—7.7(1.3) × 10⁻⁵/m] were significantly higher compared with villages located in smokeless coal deposits [GM(GSD): 3.3(1.4) × 10⁻⁵/m] ($p < 0.05$).

Outdoor pollution associated with coal type, ventilation, and season. Table 3 shows the concentrations of outdoor pollutants associated with coal type, stove ventilation, and season. Outdoor BaP and NAP concentrations in villages using smoky coal were significantly higher than those using smokeless coal [smoky coal vs smokeless coal, GM(GSD): 10.2(1.7) vs 5.7(1.4) ng/m³ for BaP, 795(2.4) vs 303(1.6) ng/m³ for NAP]; and $p < 0.05$ via Tukey HSD test. Further, PM_{2.5} concentrations were not significantly higher in these villages [smoky coal vs smokeless coal, GM(GSD): 47.7(1.7) vs 35.0(2.4) µg/m³]. However, using smoky coal was associated with significantly lower outdoor SO₂%Detect compared with using smokeless coal (50.0% vs 21.6%; $p < 0.05$). Overall, villages with greater than 50% of the households using chimneys generally had higher outdoor PM_{2.5}, BaP, NAP, and NO₂ compared to those with ≤ 50% chimney use, although these differences were not statistically significant.

ANOVA testing revealed significant seasonal variation in concentrations of the four outdoor pollutants (Table 3, $p < 0.05$). Concentrations of PM_{2.5}, NAP, BaP, and NO₂ were lowest in autumn. PM_{2.5} levels were 1.8 times as high in winter than in autumn (GM: 62.2 vs 35.3 µg/m³, $p < 0.05$). NAP was significantly higher in winter compared with either autumn or spring. Outdoor BaP in spring was the highest (GM: 11.9 ng/m³, $p < 0.05$) compared with levels in the autumn (GM: 7.4 ng/m³). Compared with levels in autumn (GM: 77.0 µg/m³), NO₂ levels were significantly higher during the spring and winter (GM: 97.9 and 107.1 µg/m³, respectively). There was no significant difference for SO₂ detection rates across seasons.




Figure 1. Map of geological coal deposits and study villages in Xuanwei and Fuyuan counties. Classification of coal regions based on the State Standard of China Coal Classification (GB5751-86); 1/3 coking, coking, gas fat, and meager lean coals are subtypes of smoky coal. *Historic smokeless coal deposit. Figure adapted from Fig. 1 in “Lung cancer risk by geologic coal deposits: A case-control study of female never-smokers from Xuanwei and Fuyuan, China” by Wong et al.¹⁷. Adapted with permission.

Correlation between outdoor and indoor air pollution. Outdoor $PM_{2.5}$ concentrations were moderately correlated with median indoor $PM_{2.5}$ concentrations that were measured in four to five households in each village (Spearman $r_s = 0.41$, $p < 0.0001$). Outdoor NO_2 concentrations were also moderately correlated with median indoor NO_2 concentrations ($r_s = 0.43$, $p < 0.0001$). However, a relatively weaker but non-significant correlation was found between outdoor and indoor BaP concentrations ($r_s = 0.21$, $p = 0.06$).

Characteristics	N (k) ^a	AM (SD)	Median (25%,75% percentile)
Daily average outside temperature, °C	88 (29)	10.7 (5.8)	11.2 (5.6,16.4)
Daily average outside humidity, %	88 (29)	78 (16.8)	82.2 (69.9,90.0)
Population	29 (29)	840 (750)	570 (420,920)
Season^b			
Autumn	33 (17)		
Spring	25 (13)		
Summer	2 (1)		
Winter	28 (15)		
Proportion of chimney stoves in a village, k (%)			
> 50%	22 (75.9)		
≤ 50%	7 (24.1)		
Factory within 5 km, k (%)			
Yes	4 (13.8)		
No	21 (72.4)		
Unknown	4 (13.8)		
Coal mine within 5 km, k (%)			
Yes	4 (13.8)		
No	21 (72.4)		
Unknown	4 (13.8)		
Power plants within 5 km, k (%)			
Yes	3 (10.3)		
No	22 (75.9)		
Unknown	4 (13.8)		

Table 1. Environmental and meteorological characteristics of the villages. *AM* arithmetic mean, *SD* standard deviation, *km* kilometer. ^aN = the number of measurements; k = the number of villages. ^bMeasurements in 16 villages were performed across multiple seasons.

Coal deposit in region	PM _{2.5} (µg/m ³)			BaP (ng/m ³)			NAP (ng/m ³)			SO ₂		NO ₂ (µg/m ³)			PM _{abs} (10 ⁻⁵ /m) [†]		
	N	AM ^a	GM (GSD) ^b	N	AM	GM (GSD)	N	AM	GM (GSD)	N	%Detect	N	AM	GM (GSD)	N	AM	GM (GSD)
Overall	85	53.1	45.3 (1.8)	51	11.1	9.7 (1.7)	58	1,045.8	707.7 (2.4)	88	26.1	87	97.6	91.5 (1.4)	66	5.5	5.0 (1.6)
Xuanwei	41	58.1	51.6 (1.7)	28	12.1	10.5 (1.7)	34	1,078.8	729.9 (2.3)	44	34.8	44	97.5	93.2 (1.4)	34	6.6	6.1 (1.6)
1, 2, 4, 7, 8 [%]	20	56.8	47.4 (1.8)	14	12.7	10.8 (1.8)	20	1,058.4	813.6 (2.2)	22	18.2	22	101.7	97.8 (1.3)	18	6.4	5.8 (1.7) ^c
9 [%]	4	70.1	67.9 (1.3)	4	13.7	12.0 (1.8)	4	629.0	620.0 (1.2)	4	25.0	4	118.5	116.8 (1.2)	4	7.2	6.9 (1.4) ^c
10 [%]	4	59.1	57.9 (1.3)	2	19.8	19.8 (1.0)	0	–	–	4	50.0	4	113.0	107.5 (1.4)	4	7.9	7.7 (1.3) ^c
LS ^{*,%}	4	49.2	47.8 (1.3)	3	6.2	6.1 (1.2)	4	2,282.6	682.8 (5.3)	4	25.0	4	85.1	83.7 (1.2)	2	5.9	5.8 (1.2)
Fuyuan	44	48.5	40.2 (2.0)	23	9.9	8.9 (1.6)	24	999.2	677.5 (2.6)	44	34.1	43	97.7	89.7 (1.5)	32	4.2	4.0 (1.4)
12, 13, 14, 38 [%]	10	58.8	50.8 (1.8)	8	9.2	8.7 (1.5)	8	1,272.8	1,093.9 (1.9)	10	20.0	9	88.9	87.6 (1.2)	4	7.2	6.6 (1.6) ^c
16, 17, 19 [%]	22	48.5	42.1 (1.8)	13	11.0	10.1 (1.6)	9	1,276.0	827.1 (3.2)	22	27.3	22	97.5	92.0 (1.5)	18	4.9	4.7 (1.3)
20, 27 ^{**,%}	21	48.3	39.0 (2.2)	7	8.4	7.4 (1.7)	13	475.2	413.0 (1.7)	22	31.8	22	92.8	81.7 (1.6)	16	3.5	3.3 (1.4)

Table 2. Concentrations of outdoor air pollutants in villages of Xuanwei and Fuyuan by coal deposit. ^aAM arithmetic mean; ^bGM geometric mean, *GSD* geometric standard deviation; ^cp < 0.05 when compared with deposit 20, 27 deposits via Tukey HSD test. *LS is located in deposit 27 in Xuanwei; **Deposit 27 is located in Xuanwei; [%]Deposit numbers refer to the map locations of each coal source shown in Fig. 1. *NAP* naphthalene, *BaP* Benzo(a)pyrene, *SO*₂ sulfur dioxide, *NO*₂ nitrogen dioxide; [†]p < 0.05 among coal deposits via ANOVA test, data were published in reference¹⁵.

Determinants of outdoor pollutants. Of all of the variables considered in the mixed models, coal type from the local coal mine, the percentage of homes in each village with chimney stoves, and season were identified as common factors contributing to outdoor PM_{2.5}, BaP, and NAP (Table 4). For NO₂ only, the average indoor concentration was identified as a contributing factor to its corresponding outdoor concentration (GMR = 1.01, p < 0.05). Coal type and the use of fire pits for cooking were associated with outdoor SO₂ detection rates.

Burning smoky coal and having > 50% of households with chimneys was associated with higher levels of outdoor BaP (GMR = 2.79, p < 0.05 for smoky coal compared to smokeless coal; GMR = 1.49, p < 0.05 for villages having > 50% of households with chimneys compared to those having ≤ 50% of households with chimneys) when adjusted for the variables shown in Table 4. However, using smoky coal was associated with a lower outdoor SO₂ detection rate (OR = 0.43, p < 0.05). Outdoor PM_{2.5} (GMR = 1.86, p < 0.05), BaP (GMR = 1.46, p < 0.05), and NAP

	PM _{2.5} (µg/m ³)			BaP (ng/m ³)			NAP (ng/m ³)			SO ₂		NO ₂ (µg/m ³)		
	N	AM ^a	GM (GSD) ^b	N	AM ^a	GM (GSD) ^b	N	AM ^a	GM (GSD) ^b	N	%Detect	N	AM ^a	GM (GSD) ^b
Coal type from coalmines														
Smokeless coal	14	46.1	35.0 (2.4)	4	5.9	5.7 (1.4)	7	331	303 (1.6)	14	21.6	14	106.3	90.8 (1.7)
Smoky coal	71	54.5	47.7 (1.7)	47	11.6	10.2 (1.7) ^c	51	1,144	795 (2.4) ^c	74	50.0 ^d	73	95.9	91.6 (1.4)
Proportion of ventilated stoves in a village														
≤ 50%	19	52.1	44.4 (2.0)	9	8.7	8.1 (1.5)	15	666	543 (1.9)	20	30.0	20	89.9	84.5 (1.4)
> 50%	66	53.4	45.6 (1.8)	42	11.6	10.1 (1.7)	43	1,178	776 (2.5)	68	25.0	67	99.9	93.7 (1.4)
Season														
Autumn	32	42.3	35.3 (1.9)	16	7.9	7.4 (1.4)	28	763.3	547.4 (2.3)	33	30.3	33	80.7	77.0 (1.4)
Spring	25	51.2	43.9 (1.9)	21	13.4	11.9 (1.7) ^e	17	813.9	629.2 (2.1)	24	24.0	24	105.4	97.9 (1.4) ^e
Summer	0	–	–	0	–	–	0	–	–	2	0.0	2	81.1	76.3 (1.6)
Winter	28	67.2	62.2 (1.5) ^e	14	11.3	9.9 (1.7)	13	1957.7	1,435.5 (2.3) ^{e,f}	28	25.0	28	112.0	107.1 (1.4) ^e

Table 3. Concentrations of outdoor air pollutants in villages of Xuanwei and Fuyuan, by coal type, stove type, and season. *NAP*, naphthalene, *BaP* Benzo-a-pyrene, *SO*₂ sulfur dioxide, *NO*₂ nitrogen dioxide. ^aAM arithmetic mean. ^bGM geometric mean, *GSD* geometric standard deviation. ^c*p* < 0.05 when compared with smokeless coal via Tukey HSD test. ^d*p* < 0.05 when compared with smokeless coal via Fisher's exact test. ^e*p* < 0.05 when compared with autumn via Tukey HSD test. ^f*p* < 0.05 when compared with spring via Tukey HSD test.

(GMR = 3.17, *p* < 0.05) concentrations were significantly higher in winter, and BaP concentrations (GMR = 1.69, *p* < 0.05) were higher in spring and summer compared to the corresponding pollutant levels in autumn. Outdoor PM_{2.5} and NAP did not vary significantly by coal deposit, but BaP levels in LS site (GMR = 0.44, *p* < 0.05) were significantly lower than at other sites. For coal deposits located in northeast Xuanwei, NO₂ levels for the smoky coal deposits (i.e. 1,2,4,7,8 and 9) were significantly higher compared to the smokeless coal deposits (i.e. 20, 27).

In villages with reference entry levels, the log-transformed values for outdoor PM_{2.5}, BaP, NAP, and NO₂ were 3.24 ln-µg/m³, 0.97 ln-ng/m³, 5.38 ln-ng/m³, and 3.67 ln-µg/m³, respectively.

Discussion

The Global Burden of Disease project (GBD) found that household solid fuel use accounted for 12% of ambient PM_{2.5} globally in 2010²⁰, with higher contribution in China (19% in 2013) and India (24% in 2015)^{21,22}. Exposure assessment for the GBD also showed substantial exposures occurring in rural areas²³. Potential sources of rural ambient air pollution may be from households using solid fuels for cooking and heating, from nearby urban and rural sources, and from secondary pollutants at intercontinental scales⁸.

Only a handful of studies have reported ambient concentrations in rural areas, as health-damaging air pollution has been considered a largely urban phenomenon⁸. Our study revealed high levels of outdoor air pollution in a rural area of China with a high incidence of lung cancer. The 24-h geometric mean outdoor PM_{2.5} concentrations in villages of the two rural counties, 51.6 µg/m³ in Xuanwei and 40.2 µg/m³ in Fuyuan, are almost two-fold that of the outdoor PM_{2.5} guideline value set by the World Health Organization (25 µg/m³)³. Further, measured PM_{2.5} concentrations were similar to the population-weighted average exposure to PM_{2.5} (52 µg/m³) in China²⁴, higher than those in rural areas in Hong Kong (24.9–30.0 µg/m³)²⁵, and comparable to levels at suburban sites reported in an earlier study in Guangdong, Wuhan, Lanzhou and Chongqing, China (GM: 39–94 µg/m³)²⁶.

The geometric means of outdoor BaP were 10.5 and 8.9 ng/m³ in Xuanwei and Fuyuan, respectively, which were up to 105 times as high as the background levels in rural areas of the United States reported by the Agency for Toxic Substances and Disease Registry²⁷. Both outdoor BaP and NAP in this rural area of China were much higher than the levels found in the rural Cantabria region in Spain (0.15 ng/m³ for BaP and < 0.01 ng/m³ for NAP)²⁸. In Xuanwei and Fuyuan, the combined outdoor BaP and NAP levels were five-fold and two-fold higher than rural air levels in central Taiwan, respectively (1.5 ng/m³ for BaP and 223 ng/m³ for NAP)²⁹. Outdoor NAP levels in our data were even higher than the highest NAP exposure category among occupationally exposed U.S. Air Force personnel³⁰. Average BaP levels exceeded the 24-h criterion of the Chinese national standard (2.5 ng/m³) by a factor of two³¹.

Varying from 85.1 to 118.5 across all coal deposits, the arithmetic mean outdoor NO₂ concentrations in our study area were ~98 µg/m³, two to ten times as high as those found in a rural area in India (4.7–9.6 µg/m³)³² and an industrial city in Turkey (14.0–57.5 µg/m³)³³.

Outdoor SO₂ detectable concentrations in our study were negligible compared with the levels measured in a rural area in India (2.5–5.2 µg/m³)³² and an industrial city in Turkey (5.9–41.2 µg/m³)^{32,33}. This is consistent with the finding that the rural residents used low sulfur coals in this study area³⁴. However, using smoky coal was associated with a significantly lower outdoor SO₂ detection rate than using smokeless coal (%Detect: 21.6 vs 50%), which is also consistent with both the observations of indoor SO₂¹⁴ and in a coal composition analysis³⁴.

In South Asia, regional concentrations of ambient PM_{2.5} derived from household cooking with solid fuels in 2010 was 8.6 µg/m³, which contributed 12% to ambient PM_{2.5} levels globally²⁰. In India, HAP contributes 22–52% to ambient air pollution and 24.2% to ambient PM_{2.5} in rural India^{22,35}. Although our study was not designed to measure emissions at the outlet of household chimneys, we calculated the ratio of outdoor levels

	ln-PM _{2.5}			ln-BaP			ln-NAP			SO ₂			ln-NO ₂			
	β	95% CI	GMR ^a	β	95% CI	GMR	β	95% CI	GMR	β	95% CI	OR	β	95% CI	GMR	
Coal type																
Smokeless coal	Ref		1.00	Ref		1.00	Ref		1.00	Ref		1.00				
Smoky coal	0.31	-0.31, 0.93	1.36	1.02	0.29, 1.76	2.79	0.89	0.04, 1.75	2.44	-0.85	-1.58, -0.12	0.43	Not included			
Proportion of chimney stoves in a village																
≤ 50%	Ref		1.00	Ref		1.00	Ref		1.00							
> 50%	-0.02	-0.39, 0.35	0.98	0.40	0.01, 0.79	1.49	0.06	-0.53, 0.64	1.06		Not included		Not included			
Season																
Autumn	Ref			Ref		1.00	Ref		1.00				Ref		1.00	
Spring/summer ^b	0.19	-0.18, 0.56	1.21	0.52	0.21, 0.84	1.69	0.29	-0.14, 0.71	1.33		Not included		-0.04	-0.21, 0.14	0.96	
Winter	0.62	0.25, 0.99	1.86	0.38	0.04, 0.72	1.46	1.15	0.65, 1.65	3.17				0.17	0.00, 0.33	1.18	
Population, per 1,000																
	0.46	-0.04, 0.95	1.58	Not included			Not included			Not included			Not included			
Use firepit to cook, 10%																
Not included				Not included			Not included			0.20	-0.31, 0.71		1.22	Not included		
Averaged indoor NO₂, μg/m³																
Not applicable				Not applicable			Not applicable			Not applicable			0.010	0.006, 0.014	1.01	
Coal deposit																
20,27	Ref			Ref		1.00	Ref		1.00	Not included			Ref		1.00	
1, 2, 4, 7, 8	-0.17	-0.79, 0.44	0.84	-0.22	-0.85, 0.41	0.80	-0.16	-0.95, 0.63	0.85				0.18	0.01, 0.35	1.20	
9	-0.74	-2.13, 0.66	0.48	-0.16	-0.91, 0.59	0.85	-0.04	-1.17, 1.08	0.96				0.51	0.21, 0.81	1.66	
10	-0.11	-0.98, 0.77	0.90	0.08	-0.81, 0.96	1.08	-	-	-				0.31	-0.01, 0.62	1.36	
LS	-1.40	-2.88, 0.09	0.25	-0.81	-1.61, -0.01	0.44	-0.38	-1.53, 0.76	0.68				0.15	-0.16, 0.46	1.16	
12, 13, 14, 38	-0.18	-0.84, 0.48	0.83	-0.30	-0.94, 0.34	0.74	0.59	-0.31, 1.50	1.81				0.21	-0.01, 0.44	1.24	
16, 17, 19	-0.32	-0.95, 0.30	0.72	-0.48	-1.12, 0.17	0.62	0.04	-0.79, 0.86	1.04				0.08	-0.10, 0.25	1.08	
Intraclass correlation																
Between villages		1			1			0.86			1			1		
Reference value ^c	3.24 ln-μg/m ³			0.97 ln-ng/m ³			5.38 ln-ng/m ³			-			3.67 ln-μg/m ³			

Table 4. Significant determinants of outdoor air concentrations from mixed modeling. *NAP* naphthalene, *BaP* Benzo(a)pyrene, *SO₂* sulfur dioxide, *NO₂* nitrogen dioxide. ^a*GMR* = geometric mean ratio = GM (estimate)/GM (reference) = Exp (β), number is bold if β is significantly different from zero (p < 0.05). ^bMerged due to few outdoor measurements in summer. ^cReference value represents log transformed value for the reference model entry.

over the average indoor concentrations measured in each village visited in Xuanwei and Fuyuan. The outdoor/indoor (O/I) ratios were ~ 10 to 30% for PM_{2.5}¹², ~ 20 to ~ 60% for BaP¹³, and ~ 70 to 80% for NO₂¹⁴ depending on the various household solid fuels that were used. It is not surprising that NO₂ presents a higher O/I ratio due to the higher penetration rates of gaseous pollutants. The higher O/I ratio for BaP compared to PM_{2.5} may imply other outdoor BaP sources, such as from emissions from other households in the village. Taken together, these results suggest that indoor air pollution due to coal combustion in households is a significant source of outdoor air pollution in rural villages.

Ventilated stove and fuel use in rural settings may have different effects on indoor, outdoor air pollution and personal exposures. A previous study of Xuanwei smoky coal use assessed the long-term health benefits of converting from unvented stoves to either stoves with chimneys or portable stoves (which were intended to be lit outdoors before being carried inside for use), and observed reduced indoor air exposures and lung cancer risks^{18,19}. This study is the first to evaluate outdoor air pollution in this rural area that has a high incidence of lung cancer. Installing chimneys to simply discharge pollutants from inside homes to the outdoors may increase outdoor concentrations and as a result not sufficiently reduce the overall exposure to carcinogens; therefore,

removal of the pollution source by moving populations up the “energy ladder” towards the use of cleaner fuels (e.g. biogas and electricity) should be strived for.

Meteorological factors may affect the spreading of primary pollutants and the formation of secondary pollutants in rural villages³⁶. A study conducted in an urban area of Hong Kong found that temperature, humidity, and solar irradiation played a vital role in the variation of the O/I ratio, which increased with upward changes of these weather parameters³⁷. However, in the current study, seasonal category was found to more accurately predicate the outdoor exposure than meteorological factors.

Limitations of this study include small sample size and availability of a background monitoring spot in this rural area.

This study showed that outdoor air pollution in a rural region of China with a high incidence of lung cancer was associated with coal type and the proportion of ventilated stoves in a village. These findings suggest that the further reduction of adverse health effects in rural villages from the indoor burning of coal, will most likely require the use of stoves that reduce environmental exhaust, and ultimately the replacement of coal with cleaner fuel types.

Methods

Study design and air pollution measurements. The exposure assessment study design and population have been described in detail elsewhere¹². We reported here the part of outdoor air measurement. Briefly, outdoor air measurements were taken between August 2008 and June 2009 from 29 selected villages in Xuanwei and Fuyuan as part of a large case–control study of lung cancer, and a cross-sectional molecular epidemiological study. Villages were selected to represent all major geological coal deposit areas based on a local geologic analysis of coal-type in Xuanwei and Fuyuan counties. In each selected village, four or five households were selected to conduct personal and indoor exposure measurements, while outdoor air measurements were conducted in a central location within each village in phase I. In phase II, approximately half of the villages ($n = 16$) and households were visited for a second round of repeated measurements 2–9 months later.

Village background information. Background demographic information for each village was collected through an in-person interview with either a local doctor or the village head. Global Positioning System (GPS) coordinates were recorded on-site. Survey information included: house types and ventilation, fuels used for cooking and heating, stove types, main coal mines providing coal to the village, village altitude, total population, land area in km², percentage of households having a television set, average household income, and presence of nearby industries. Distribution of some variables were shown in Table 1.

Outdoor sample collection and analysis. An ambient air monitoring station was installed at a central location within each village away from any direct sources of emissions (e.g. chimneys). The air monitoring methods that were used were similar to those used to measure indoor air pollution and are described in detail^{12,13,15}. In brief, samples of fine PM and associated particle phase PAHs were collected on 37 mm Teflon filters using a cyclone with an aerodynamic cut-off of 2.5 μm (model BGI, GK 2.05SH) at a flow rate of 3.5 L/min ($\pm 20\%$). XAD-2 sorbent tubes were used to measure gas phase PAHs at an air flow rate of 100 mL/min. Particulate mass was measured by pre- and post-weighing of the filters in an environmentally-controlled weighing room using a microbalance at 1 μg accuracy. PAH extraction from the Teflon filters and the XAD-2 sorbent tubes was performed using the US EPA 3540C-1996 (Soxhlet extraction) method to determine concentrations of 16 PAHs by a gas chromatograph connected to a mass spectrometer (Shimadzu QP2010 Plus). Passively diffusing OGAWA badges were deployed to collect measurements of NO₂ and SO₂. Flow injection analysis and ion chromatography were used to determine the concentrations of NO₂ and SO₂, respectively. In addition, a weather station (WeatherLink Wireless Vantage Pro2) was deployed to record meteorological factors (e.g. temperature, wind speed, humidity, rainfall amount). Two sequential 24-h outdoor air measurements were conducted in each village in parallel with indoor and personal measurements. For quality control purposes, field blank and 13 duplicate filter samples as well as duplicate OGAWA badges (~10%) were collected. More than 97% of field blank filters reported non-detect PAHs. The coefficient of variation of the duplicate samples was 13% for PM_{2.5}, 25% for BaP, 38% for NAP, and 27% for NO₂, respectively. The percentage agreement in detect vs non-detect of the duplicate SO₂ samples was 90%.

Statistical analyses. Normal probability plots indicated that the measured values could be best described by a log-normal distribution; therefore, outdoor air pollution concentrations were natural log-transformed to approximate normal distributions for the statistical analyses that were conducted. Outdoor measurements were summarized as arithmetic means (AM), geometric means (GM), and geometric standard deviations (GSD) by coal deposit. Meteorological factors are summarized as AM, standard deviations (SD), and medians. Spearman correlations were calculated between concentrations of outdoor and indoor pollutants. Analysis of variance (ANOVA) and Tukey Honestly Significant Difference (HSD) testing was performed on log-transformed values to assess differences between coal deposits, fuel types, stove ventilation, and season. Due to the large proportion of undetectable values of SO₂ measurements (73.9%), detection rate (%Detect) was calculated for each coal deposit and overall region. Linear mixed effect models were used to identify variables which may be associated with outdoor PM_{2.5}, BaP, NAP, and NO₂, while a mixed effects logistic model was used for SO₂. Villages were assigned random effects with a variance-components covariance structure. Multiple variables were considered for inclusion as fixed effects including coal deposits, season (winter, spring, summer, autumn), proportion of ventilated stoves (i.e. chimneys) in a village, meteorological factors, proportion of solid fuel types in a village to heat rooms or cook, proportion of stove types used in a village, type of coal mines (i.e. smoky coal or smokeless

coal), altitude, village area, population size, nearby industry, average indoor concentration of the pollutant measured in a village, average income and other surrogates of socioeconomic status such as proportion of households having a TV set. Inclusion of variables in the final model was based on the combination of their influence on the Akaike information criterion (AIC) score. The linear mixed effect model can be expressed with the following equation:

$$y_{ijf} = \mu + \beta_1 x_1 + \beta_2 x_2 \dots \beta_n x_n + b_1 I_i + b_2 J_{ij} + \varepsilon_{ijf}$$

where y_{ijf} represents the natural log-transformed value of outdoor pollutant concentration being modeled for coal deposit i in village j on day f ; μ represents the intercept value (i.e., the log-transformed value for the reference group); β_1 through β_n represent the fixed effect variable coefficients for variables x_1 through x_n ; $b_1 I_i$ represents the coefficient for coal deposit i ; $b_2 J_{ij}$ represents the random effect coefficient for village j from coal deposit i ; and ε_{ijf} represents the error for village j in coal deposit i on day f .

All analyses were carried out using SAS/STAT software, version 9.4 of the SAS System for Windows. Copyright © 2016 SAS Institute Inc. SAS and all other SAS Institute Inc. product or service names are registered trademarks or trademarks of SAS Institute Inc., Cary, NC, USA.

The study protocol was approved by the institutional review boards of the National Cancer Institute and China National Environmental Monitoring Center. All participants provided written informed consent prior to participating in the study. This study was conducted in accordance to the World Medical Association Declaration of Helsinki's recommendations for human subject protection.

Consent for publication. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does the mentioning of trade names, commercial products, or organizations imply endorsement by the U.S. Government.

Data availability

All data generated or analyzed during this study are included in this published article.

Received: 30 January 2020; Accepted: 16 June 2020

Published online: 09 July 2020

References

- World Health Organization. *7 Million Premature Deaths Annually Linked to Air Pollution*. <https://www.who.int/mediacentre/news/releases/2014/air-pollution/en/> (2014).
- World Health Organization. *Ambient Air Pollution: A Global Assessment of Exposure and Burden of Disease* (WHO Document Production Services, Geneva, 2016).
- World Health Organization. *Ambient (Outdoor) Air Quality and Health*. <https://www.who.int/mediacentre/factsheets/fs313/en/> (2016).
- Cohen, A. J. *et al.* Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: An analysis of data from the Global Burden of Diseases Study 2015. *Lancet* **389**, 1907–1918. [https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6) (2017).
- Loomis, D. *et al.* The carcinogenicity of outdoor air pollution. *Lancet Oncol.* **14**, 1262–1263. [https://doi.org/10.1016/s1470-2045\(13\)70487-x](https://doi.org/10.1016/s1470-2045(13)70487-x) (2013).
- Loomis, D., Huang, W. & Chen, G. The International Agency for Research on Cancer (IARC) evaluation of the carcinogenicity of outdoor air pollution: Focus on China. *Chin. J. Cancer* **33**, 189–196. <https://doi.org/10.5732/cjc.014.10028> (2014).
- International Agency for Research on Cancer. *Outdoor air Pollution/IARC Monographs on the Evaluation of Carcinogenic Risks to Humans*, vol. 109 (World Health Organization, Geneva, 2016).
- Smith, K. R. in *Testimony at the Commission on Sustainable Development Session on Integrated Approaches to Addressing Air Pollution* (United Nations, New York, 2006).
- Zhang, J. J. & Smith, K. R. Household air pollution from coal and biomass fuels in China: Measurements, health impacts, and interventions. *Environ. Health Perspect.* **115**, 848–855. <https://doi.org/10.1289/ehp.9479> (2007).
- Mumford, J. L. *et al.* Lung cancer and indoor air pollution in Xuan Wei, China. *Science* **235**, 217–220. <https://doi.org/10.1126/science.3798109> (1987).
- Barone-Adesi, F. *et al.* Risk of lung cancer associated with domestic use of coal in Xuanwei, China: Retrospective cohort study. *BMJ* **345**, e5414 (2012).
- Hu, W. *et al.* Personal and indoor PM_{2.5} exposure from burning solid fuels in vented and unvented stoves in a rural region of China with a high incidence of lung cancer. *Environ. Sci. Technol.* **48**, 8456–8464. <https://doi.org/10.1021/es502201s> (2014).
- Downward, G. S. *et al.* Polycyclic aromatic hydrocarbon exposure in household air pollution from solid fuel combustion among the female population of Xuanwei and Fuyuan Counties, China. *Environ. Sci. Technol.* **48**, 14632–14641. <https://doi.org/10.1021/es504102z> (2014).
- Seow, W. J. *et al.* Indoor concentrations of nitrogen dioxide and sulfur dioxide from burning solid fuels for cooking and heating in Yunnan Province, China. *Indoor Air* **26**, 776–783. <https://doi.org/10.1111/ina.12251> (2016).
- Downward, G. S. *et al.* Outdoor, indoor, and personal black carbon exposure from cookstoves burning solid fuels. *Indoor Air* **26**, 784–795. <https://doi.org/10.1111/ina.12255> (2016).
- Lan, Q. *et al.* Variation in lung cancer risk by smoky coal subtype in Xuanwei, China. *Int. J. Cancer* **123**, 2164–2169. <https://doi.org/10.1002/ijc.23748> (2008).
- Wong, J. Y. Y. *et al.* Lung cancer risk by geologic coal deposits: A case-control study of female never-smokers from Xuanwei and Fuyuan, China. *Int. J. Cancer* **144**, 2918–2927. <https://doi.org/10.1002/ijc.32034> (2019).
- Lan, Q., Chapman, R. S., Schreinemachers, D. M., Tian, L. & He, X. Household stove improvement and risk of lung cancer in Xuanwei, China. *J. Natl. Cancer Inst.* **94**, 826–835. <https://doi.org/10.1093/jnci/94.11.826> (2002).
- Hosgood, H. D. *et al.* Portable stove use is associated with lower lung cancer mortality risk in lifetime smoky coal users. *Br. J. Cancer* **99**, 1934–1939. <https://doi.org/10.1038/sj.bjc.6604744> (2008).
- Chafe, Z. A. *et al.* Household cooking with solid fuels contributes to ambient PM_{2.5} air pollution and the burden of disease. *Environ. Health Perspect.* **122**, 1314–1320. <https://doi.org/10.1289/ehp.1206340> (2014).

21. GBD MAPS Working Group. *Burden of Disease Attributable to Coal-Burning and Other Air Pollution Sources in China*. <https://www.healtheffects.org/publication/burden-disease-attributable-coal-burning-and-other-air-pollution-sources-china> (2016).
22. GBD MAPS Working Group. *Burden of Disease Attributable to Major Air Pollution Sources in India*. <https://www.healtheffects.org/publication/gbd-air-pollution-india> (2018).
23. Brauer, M. *et al.* Exposure assessment for estimation of the global burden of disease attributable to outdoor air pollution. *Environ. Sci. Technol.* **46**, 652–660. <https://doi.org/10.1021/es2025752> (2012).
24. Rohde, R. A. & Muller, R. A. Air pollution in China: Mapping of concentrations and sources. *PLoS One* **10**, e0135749. <https://doi.org/10.1371/journal.pone.0135749> (2015).
25. Cao, J. J. *et al.* Indoor/outdoor relationships for PM_{2.5} and associated carbonaceous pollutants at residential homes in Hong Kong—case study. *Indoor Air* **15**, 197–204. <https://doi.org/10.1111/j.1600-0668.2005.00336.x> (2005).
26. Wei, F. *et al.* Ambient concentrations and elemental compositions of PM₁₀ and PM_{2.5} in four Chinese cities. *Environ. Sci. Technol.* **33**, 4188–4193. <https://doi.org/10.1021/es9904944> (1999).
27. ATSDR (Agency for Toxic Substances and Disease Registry). *Toxicological Profile for Polycyclic Aromatic Hydrocarbons* (ATSDR, Atlanta, 1995).
28. Arruti, A., Fernandez-Olmo, I. & Irabien, A. Evaluation of the urban/rural particle-bound PAH and PCB levels in the northern Spain (Cantabria region). *Environ. Monit. Assess.* **184**, 6513–6526. <https://doi.org/10.1007/s10661-011-2437-4> (2012).
29. Fang, G. C., Chang, K. F., Lu, C. & Bai, H. Estimation of PAHs dry deposition and BaP toxic equivalency factors (TEFs) study at Urban, Industry Park and rural sampling sites in central Taiwan, Taichung. *Chemosphere* **55**, 787–796. <https://doi.org/10.1016/j.chemosphere.2003.12.012> (2004).
30. Egeghy, P. P., Hauf-Cabalo, L., Gibson, R. & Rappaport, S. M. Benzene and naphthalene in air and breath as indicators of exposure to jet fuel. *Occup. Environ. Med.* **60**, 969–976. <https://doi.org/10.1136/oem.60.12.969> (2003).
31. Ministry of Environmental Protection of the People's Republic of China. *Ambient Air Quality Standards (GB 3095–2012)*. <https://kjs.mep.gov.cn/hjbhbz/bzwb/dqjhbh/dqjzlbz/201203/W020120410330232398521.pdf> (2016).
32. Gupta, A., Kumar, R., Kumari, K. M. & Srivastava, S. S. Measurement of NO₂, HNO₃, NH₃ and SO₂ and related particulate matter at a rural site in Rampur, India. *Atmos. Environ.* **37**, 4837–4846. <https://doi.org/10.1016/j.atmosenv.2003.07.008> (2003).
33. Bozkurt, Z. *et al.* Determination of the personal, indoor and outdoor exposure levels of inorganic gaseous pollutants in different microenvironments in an industrial city. *Environ. Monit. Assess.* **187**, 590. <https://doi.org/10.1007/s10661-015-4816-8> (2015).
34. Downward, G. S. *et al.* Heterogeneity in coal composition and implications for lung cancer risk in Xuanwei and Fuyuan counties, China. *Environ. Int.* **68**, 94–104. <https://doi.org/10.1016/j.envint.2014.03.019> (2014).
35. Chowdhury, S. *et al.* Policy Brief: *The Contribution of Household Fuels to Ambient Air Pollution in India—A Comparison of Recent Estimates*. <https://ccapc.org.in/policy-briefs/2019/5/30/the-contribution-of-household-fuels-to-ambient-air-pollution-in-india-a-comparison-of-recent-estimates> (2019).
36. Kamens, R. M., Guo, Z., Fulcher, J. N. & Bell, D. A. The influence of humidity, sunlight, and temperature on the daytime decay of polyaromatic hydrocarbons on atmospheric soot particles. *Environ. Sci. Technol.* **22**, 103–108. <https://doi.org/10.1021/es00166a012> (1988).
37. Chan, A. T. Indoor–outdoor relationships of particulate matter and nitrogen oxides under different outdoor meteorological conditions. *Atmos. Environ.* **36**, 1543–1551. [https://doi.org/10.1016/S1352-2310\(01\)00471-X](https://doi.org/10.1016/S1352-2310(01)00471-X) (2002).

Acknowledgements

The authors would like to acknowledge Jackie King from BioReliance, and Peter Hui and Nathan Appel from Information Management Systems, Inc. for their support. This project was supported by the National Institutes of Health Intramural Research Program, National Natural Science Foundation of China (81702274) and Yunnan Health Training Project of High-Level Talents (H-2018025).

Author contributions

Q.L., R.V., N.R. and F.W. conceived of and designed the study. B.R., W.H., J.L., J.X., J.H., L.Z., G.W., Y.H., K.Y. and Y.C. participated in data collection. W.H. and G.D. conducted the analyses and were primarily responsible for drafting the paper. J.W., B.B., L.P., R.J., W.J.S., and D.H. took part in interpretation of results and manuscript preparation. All authors reviewed and approved the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to W.H.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2020